QK
431
M93

SECOND SYSTEMATIC CENSUS

OF

AUSTRALIAN PLANTS,

WITH

Chronologic, Literary and Geographic Annotations;

BY

BARON FERDINAND VON MUELLER,
K.C.M.G., M.D., Ph.D., F.R.S.
F.L.S., F.G.S., F.C.S., F.R.G.S., C.M.Z.S., H.F.R.H.S., &c.
GOVERNMENT BOTANIST FOR THE COLONY OF VICTORIA.

PART I.—VASCULARES.

"LEVATE OCULOS IN EXCELSUM ET VIDETE QUIS CREAVIT HAEC OMNIA. QUIS EDUCIT MILITIAM EORUM. —*Isaiah* xl.

MELBOURNE:
PRINTED FOR THE VICTORIAN GOVERNMENT
BY McCARRON, BIRD & CO., 479 COLLINS STREET.
1889.

THE LIBRARY
OF
THE UNIVERSITY
OF CALIFORNIA
DAVIS

Ernst Dörfler &c.

zur freundlichen Erinnerung

und für gelegentlichen Gebrauch

von Ferd. von Mueller.

SECOND SYSTEMATIC CENSUS

OF

AUSTRALIAN PLANTS,

WITH

Chronologic, Literary and Geographic Annotations;

BY

BARON FERDINAND VON MUELLER

K.C.M.G., M.D., Ph.D., F.R.S.

F.L.S., F.G.S., F.C.S., F.R.G.S., C.M.Z.S., H.F.R.H.S., &c.

GOVERNMENT BOTANIST FOR THE COLONY OF VICTORIA.

PART I.—VASCULARES.

LEVATE OCULOS IN EXCELSUM ET VIDETE QUIS CREAVIT HAEC OMNIA, QUIS EDUCIT MULTITUDINEM EORUM."—*Isaiah* xl.

MELBOURNE:

PRINTED FOR THE VICTORIAN GOVERNMENT

BY McCARRON, BIRD & CO., 479 COLLINS STREET.

1889.

LIBRARY

UNIVERSITY OF CALIFORNIA

Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

http://www.archive.org/details/secondsystematic00muelrich

To

SIR JOSEPH HOOKER, K.C.S.I., C.B., F.R.S., V.P.L.S.

AND TO

M. ALPHONSE DE CANDOLLE, F.M.R.S., F.M.L.S.,

WHO AS HEIRS OF GREAT NAMES WORTHILY SUSTAIN WORLD-WIDE ANCESTRAL FAME,

AND WHO AS LEADERS IN PHYTOGRAPHY DURING HALF A CENTURY

WILL BE PRE-EMINENT IN BIOMORPHIC SCIENCE THROUGH ALL AGES,

THESE PAGES

ARE OFFERED AS AN APPRECIATIVE TRIBUTE

FROM A YOUNG COLONY IN ANTIPODAL REMOTENESS.

DICOTYLEDONEAE.

Ray, Method. Plant. Nova 2 (1682).

CHORIPETALEAE HYPOGYNAE.

F. v. Mueller, native plants of Victoria I, 1 (1879).

DILLENIACEAE.

Salisbury, Parad. Lond. I, 73 (1801).—Genera (here) 4, Species 95.

WORMIA; Rottboell, Nye Saml. Vid. Selsk. Skrivt. II, 532 (1783). (Dilleniae subgenus).

W. alata; R. Brown in De Cand. syst. veg. I, 434 (1818) ...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,16	M.fr.VIII,124. ...	P.

HIBBERTIA; Andrews, Bot. Reposit. t. 126 (1800). (Candollea, Pleurandra, Hemistemma, Adrastaea, Ochrolasia, Hemistephus, Huttia, Warburtonia.)

H. Banksii; F. v. Mueller, pl. of Vict. I, 14, implied (1860) ...	...	...	—	—	—	—	—	Q.	—	B.fl.I,21	M.fr,II.1; XI,92.	
H. Holtzei; F. v. M. in Wing's S. Sc. Rec. new ser. I, Nov. (1885) ...	...	—	—	—	—	—	—	N.A.		—		
H. Brownii; Bentham, Fl. Austr. I, 21 (1863) ...	...	...	...	...	—	—	—	—	Q.	—	B.fl.I,21	M.fr,II.1.
H. dealbata; F. v. M., pl. of Vict. I, 14, implied (1860) ...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,21	M.fr.II,1.
H. candicans; Bentham, Fl. Austr. I, 21 (1863) ...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.I,21	
H. Benthami; F. v. M., syst. Cens. of Austr. pl. 1 (1882)..	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,21	M.fr.II,1.
H. Muelleri; Bentham, Fl. Austr. I, 21 (1863) ..	..	...	...	...	—	—	—	—	—	N.A.	B.fl.I,21	
H. ledifolia; Bentham, Fl. Austr. I, 22 (1863) ...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.I,22	
H. verrucosa; F. v. M., pl. of Vict. I, 14, implied (1860) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,22	M.fr.XI,92.
H. spicata; F. v. M., fragm. phytogr. Austral. II, 1 (1860) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,22	M.fr.XI,20,92.
H. polystachya; Bentham, Fl. Austr. I, 22 (1863) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,22	M.fr.XI,92.
H. rhadinopoda; F. v. M., fragm. XI, 91 (1880) ...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.XI,91.
H. furfuracea; Bentham, Fl. Austr. I, 23 (1863) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,23	M.fr.I,161; XI,92.
H. hypericoides; F. v. M., pl. of Vict. I, 13, implied (1860) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,23	
H. microphylla; Steudel in Lehmann, pl. Preiss. I, 273 (1844) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,23	M.fr.I,217.
H. recurvifolia; Bentham, Fl. Austr. I, 24 (1863) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,24	
H. lineata; Steudel in Lehmann, pl. Preiss. I, 272 (1844) ...	..	...	...	W.A.	—	—	—	—	—	—	B.fl.I,24	M.fr.XI,92.
H. acerosa; Bentham, Fl. Austr. I, 74 (1863) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,24	M.fr.XI,92.
H. aurea; Steudel in Lehmann, pl. Preiss. I, 272 (1844) ..	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,25	M.fr.XI,92.
H. crassifolia; Bentham, Fl. Austr. I, 25 (1863) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.I,25	
H. nitida; F. v. M., pl. of Vict. I, 15 (1860) ...	...	...	..	...	—	—	—	N.S.W.	—	—	B.fl.I,25	
H. bracteata; Bentham, Fl. Austr. I, 26 (1863) ...	...	...	...	...	—	—	—	N.S.W.	—	—	B.fl.I,25	
H. densiflora; F. v. M., pl. of Vict. I, 15 (1860) ...	...	...	...	...	—	S.A.	T. V.	N.S.W.	—	—	B.fl.I,26	M.fr.VII,125; XI,92.
H. hirsuta; Bentham, Fl. Austr. 21 (1863) ...	...	..	...	...	—	S.A.	T. —	—	—	—	B.fl.1,26	
H. synandra; F. v. M., fragm. IV, 151 (1864) ...	...	...	...	...	—	—	— —	—	Q.	—	—	M.fr.IV,151.
H. stricta; R. Brown in De Candolle, syst. veg. I, 422 (1818) ...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,27	
H. humifusa; F. v. M., pl. of Vict. I, 16, t. Suppl. I (1860) ...	...	...	...	—	—	—	V.	—	—	—	B.fl.I,27	
H. Billardieri; F. v. M., pl. of Vict. I, 14 (1860) ...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	—	B.fl.I,28	M.fr.IV,116.
H. gracilipes; Bentham, Fl. Austr. I, 28 (1863) ...	...	...	..	...	W.A.	—	— —	—	—	—	B.fl.I,28	M.fr.XI,92.
H. acicularis; F. v. M., pl. of Vict. I, 17 (1860) ...	...	...	...	...	—	S.A.	T. V.	N.S.W.	Q.	—	B.fl.I,29	M.fr.XI,93.
H. mucronata; F. v. M., pl. of Vict. I, 17 (1860) ...	...	...	...	...	W.A.	—	— —	—	—	—	B.fl.I,29	M.fr.XI,92.

H. hermannifolia ; De Candolle, syst. veg. I, 431 (1818)	—	—	—	—	N.S.W.	—	—	B.fl.I,30	
H. velutina ; R. Brown in Bentham, Fl. Austr. I, 30 (1863)	—	—	—	—	—	Q.	—	B.fl.I,30	
H. oblongata ; R. Brown in De Candolle, syst. veg. I, 431 (1818)...	—	—	—	—	—	—	N.A.	B.fl.I,30	M.fr.XI,94.
H. melhanoides ; F. v. M., fragm. IV, 116 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,116.
H. tomentosa ; R. Brown in De Candolle, syst. veg. I, 432 (1818)...	—	—	—	—	—	—	N.A.	B.fl.I,30	
H. cistifolia ; R. Brown in Ds Candolle, syst. veg. I, 431 (1818)	—	—	—	—	—	—	N.A.	B.fl.I,30	
H. echiifolia ; R. Brown in Bentham, Fl. Austr. I. 31 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,31	
H. scabra ; R. Brown in Bentham, Fl. Austr. I, 31 (1863)...	—	—	—	—	—	—	N.A.	B.fl.I,31	
H. lepidota ; R. Brown in De Candolle, syst. veg. I, 432 (1818)	—	—	—	—	—	Q.	N.A.	B.fl.I,31	
H. vestita ; Cunningham in Bentham, Fl. Austr. I, 31 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,31	
H. serpyllifolia ; R. Brown in De Cand. syst. veg. I, 430 (1818)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,32	
H. pedunculata ; R. Brown in De Cand. syst. veg. 1, 430 (1818)	—	—	—	V.	N.S.W.	—	—	B.fl.I,32	
H. ochrolasia ; Bentham, Fl. Austr. I, 32 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,32	M.fr.VII,125 ; XI,93.
H. angustifolia ; Salisbury, parad. Lond. 73 (1807)	—	—	T.	V.	—	—	—	B fl.I,33	
H. fasciculata ; R. Brown in De Candolle, syst. veg. I, 428 (1818)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,33	
H. virgata ; R. Brown in De Candolle, syst. veg. I, 428 (1818)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,34	M.fr.XI,94.
H. inclusa ; Bentham, Fl. Austr. I, 34 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,3	
H. rostellata ; Turczaninow in Bull. Soc. Mosc. XXII, part III., 8 (1849) ...	W.A.	—	—	—	—	—	—	B.fl.I,35	M.fr.XI,93.
H. glomerata ; Bentham, Fl. Aust. I, 34 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,35	
H. argentea ; Steudel in Lehmann, pl. Preiss. I, 268 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,35	
H. pilosa ; Steudel in Lehmann, pl. Preiss. I, 272 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,35	
H. montana ; Steudel in Lehmann, pl. Preiss. I, 270 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,35	M.fr.XI,93.
H. obtusifolia ; De Candolle, syst. veg. I, 428 (1818)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,36	
H. diffusa ; R. Brown in De Candolle, syst. veg. I, 424 (1818)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,36	
H. saligna ; R. Brown in De Candolle, syst. veg. I, 427 (1818)	—	—	—	—	N.S.W.	—	—	B.fl.I,37	M.fr.XI,94.
H. volubilis ; Andrews in Bot. Reposit. t. 126 (1800)	—	—	—	—	N.S.W.	Q.	—	B.fl.1,37	M.fr.IV,116 ; VII,125 ; XI,94.
H. grossularifolia ; Salisbury, parad. Lond. t. 73 (1806)	W.A.	—	—	—	—	—	—	B.fl.I,37	M.fr.I,230 ; II,182 ; XI,94.
H. dentata ; R. Brown in De Candolle, syst. veg. I, 426 (1818)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,38	M.fr.VII,37,125 ; XI,94.
H. Cunninghamii ; Aiton in Hook. Bot. Mag. t. 318 (1832)...	W.A.	—	—	—	—	—	—	B.fl.I,39	M.fr.X1,93.
H. perfoliata ; Huegel, enum. pl. Nov. Holl. austr. occ. 3 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,38	M.fr.XI,93.
H. bracteosa ; Turczaninow, in Bull. Soc. Mosc. XXV, 140 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,38	M.fr.XI,94.
H. longifolia ; F. v. M., fragm. IV, 115 (1814)	—	—	—	—	—	Q.	—	—	M.fr.IV,115.
H. oenotheroides ; F. v. M., fragm. VII, 37 (1870)	—	—	—	—	—	Q.	—	—	M.fr.VII,37.
H. glaberrima ; F. v. M., fragm. III, 1 (1862)	—	S.A.	—	—	—	Q.	—	B.fl.I,39	M.fr.XI,94.
H. Milnei ; Bentham, Fl. Austr. I, 39 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,39	
H. lasiopus ; Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	M.fr.XI,94.
H. potentilliflora ; F. v. M. in Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	M.fr.XI,94.
H. pungens ; Bentham, Fl. Austr. I, 40 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,40	
H. nutans ; Bentham, Fl. Austr. I, 40 (1863)...	W.A.	—	—	—	—	—	—	B.fl.1,40	
H. leptopus ; Bentham, Fl. Austr. I, 41 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,41	
H. stellaris ; Endlicher in Hueg. enum. 3 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,41	M.fr.XI,93.
H. glomerosa ; F. v. M., syst. Cens. of Austr. plants 2 (1882)	W.A.	—	—	—	—	—	—	B.fl.I,43	M.fr.XI,94.
H. obcuneata ; Salisbury, parad. Londin. t. 73 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,42	M.fr.XI,94.
H. teretifolia ; F. v. M., fragm. IV, 117 (1864)	W.A.	—	—	—	—	—	—	B.fl.I,43	M.fr.XI,95.
H. desmophylla ; F. v. M., fragm. XI, 95 (1880)	W.A.	—	—	—	—	—	—	B.fl.I,43	M.fr.XI,95.
H. helianthemoides ; F. v. M., syst. Cens. of Austr. plants 2 (1882)	W.A.	—	—	—	—	—	—	B.fl.I,43	
H. depressa ; Steudel in Lehm. pl. Preiss. I, 268 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,44	M.fr.XI,95.
H. Huegelii ; F. v. M., fragm. XI, 95 (1880)	W.A.	—	—	—	—	—	—	B.fl.I,44	M.fr.XI,95.
H. polygonoides ; F. v. M., fragm. IV, 116 (1864)	W.A.	—	—	—	—	—	—	B.fl.I,45	M.fr.XI,95.

H. vaginata; F. v. M., fragm. XI, 96 (1880) ...	W.A.	—	—	—	—	—	—	B.fl.I,45	M.fr.XI,96.		
H. subexcisa; Steudel in Lehmann, pl. Preiss. I. 269 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,46	M.fr.XI,95.		
H. squamosa; Turczaninow in Bull. Soc. Mosc. XXII, part III, 9 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,46			
H. unciuata; F. v. M., syst. Cens. of Austr. pl. 2 (1882) ...	W.A.	—	—	—	—	—	—	B.fl.I,46			
H. salicifolia; F. v. M., fragm. I, 161 (1859) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,46	M.fr.VII,124; XI,94.		
H. Huttii; F. v. M., fragm. VII, 123 (1870) ...	W.A.	—	—	—	—	—	—	B.fl.I,47	M.fr.XI,20.		
H. Goyderi; F. v. M., fragm. VII, 123 (1870) ...	—	—	—	—	—	—	N.A.	—	M.fr.VII,123.		
TETRACERA; Linné, gen. pl. 345 (1737).											
T. Nordtiana; F. v. M., fragm. V, 1 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr.V,1	..	P.
T. Daemeliana; F. v. M., fragm. V, 191 (1866) ...	—	—	—	—	—	Q.	—	—	M.fr.V,191.		
T. Wuthiana; F. v. M., fragm. X, 49 (1876) ...	—	—	—	—	—	Q.	—	—	M.fr.X,49.		
PACHYNEMA; R. Brown in De Candolle, syst. veg. I, 412 (1818).											
P. junceum; Bentham, Fl. Austr. I, 47 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,47	M.fr.X,124.		
P. complanatum; R. Brown in De Candolle, syst. veg. I, 412 (1818) ...	—	—	—	—	—	—	N.A.	B.fl.I.48	M.fr.X.124.		
P. dilatatum; Bentham, Fl. Austr. I, 48 (1863) ...	—	—	—	—	—	—	N.A.	B.fl.I,48	M.fr.X,124.		
P. sphenandrum; F. v. M. & Tate in Trans. R. S. of S.A. V, 79 (1882) ...	—	—	—	—	—	—	N.A.	—			

RANUNCULACEAE.

A. L. de Jussieu, Rec. de l'Acad. des sc. (1773) from B. de Jussieu (1759).—Genera 5, Species 17.

CLEMATIS; Linné, gen. pl. 163 (1737) from l'Ecluse (1576).															
C. aristata; R. Brown in De Candolle, syst. veg. I, 142 (1818) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,6	M.fr.X,2.		
C. pubescens; Huegel, enum. pl. Nov. Holl. austr. occ. 1 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.I,6			
C. glycinoides; De Candolle, syst. veg. I, 145 (1818) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,7	M.fr.IX,76; X,2.		P.
C. microphylla; De Candolle, syst. veg. I, 142, (1818) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,7	M.fr.X,2; XI,20,27		
C. Fawcetti; F. v. M., fragm. X, 1 (1876) ...	—	—	—	—	N.S.W.	Q.	—	—	M. fr. X, 1.		
ANEMONE; Tournefort, inst. rei herb. 275, t. 174 (1700) from Hippocrates, Theophrastos and Dioscorides.											
A. crassifolia; Hooker, icones pl. t. 257 (1840) ...	—	—	T.	—	—	—	—	B.fl.I,8	M.fr.X,1.		
MYOSURUS; Dillenius, catal. pl. Giess. 106, t. 4 (1749) from L'Obel (1576). (Myosuros.)											
M. minimus; Linné, sp. pl. 284 (1753) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,8	M.fr.X,2; XI,27		E., As., Afr., Am.
RANUNCULUS; Tournefort, inst. 285, t. 149 (1700) from Bock (1552) from Plinius.											
R. parviflorus; Linné, sp. pl. sec. ed. 780 (1762) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,14	M.fr.X,2; XI,27		E., As., Afr., N.Z.
R. hirtus; Banks & Solander in G. Forst. fl. ins. austr. prodr. 90 (1786) ...	W.A.	—	—	V.	N.S.W.	Q.	—	B.fl.I,13	M.fr.X,2	...	N.Z.
R. rivularis; Banks & Solander in G. Forst. fl. ins. austr. prodr. 90 (1786) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,13	M.fr.X,2	...	N.Z.
R. Muelleri; Bentham, Fl. Austr. I, 13 (1863) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,13			
R. lappaceus; Smith in Rees' Cyclopædia, XXIX (1815) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,12	M.fr.X,2	...	N.Z.
R. Gunnianus; Hooker, Jour. of Bot. I, 245, t. 233 (1834) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,11			
R. anemoneus; F. v. M. in Trans. Phil. Soc. Vict. I, 97 (1855) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,11			
R. Millani; F. v. M. in Hook. Kew misc. VII, 358 (1855) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,10	M.fr.X,2.		
R. acquatilis; Dodonæus, stirp. hist. pempt. 387 (1583) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,10	...	...	E., As., Afr., Am.
CALTHA; Ruppius, flora Jenensis 119 (1718) from Bock (1552). (Trollius.)											
C. ntroloba; F. v. M. in Trans. Phil. Soc. Vic. I, 98 (1855) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.I,15	M.fr.X,2; XI,136		N.Z.

CERATOPHYLLEAE.

J. E. Gray; Arrangm. of Brit. pl. II, 395 (1821).—Genus 1, Species I.

CERATOPHYLLUM, Linné, syst. nat. 9 (1735).											
C. demersum; Linné, spec. plant. 992 (1753) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,491	M.fr.VIII,103		E., As., Afr., Am., P.

NYMPHAEACEAE.

SALISBURY in Koenig & Sims' Ann. of Bot. II, 70 (1805).—Genera 3, Species 5.

CABOMBA; Aublet, Hist. pl. Guian. I, 321, t. 124 (1775). (Brasenia, Hydropeltis.)

C. peltata; F. v. M., Introd. to Bot. Teachings, 89 (1877)	—	—	—	V.	N.S.W.	Q	—	B.fl I,60	M.fr.X,77	...	As., Am.

NYMPHAEA; Tournefort, inst. 260, t. 137 8 (1700) from Theophrastos and Dioscorides.

N. tetragona; Georgi, Reise im Russ. Reich. I, 220 (1775)	—	—	—	—	—	Q.	—	—	...	...	...	As.
N. stellata; Willdenow, spec. pl. II, 1153 (1799)	—	—	—	—	—	Q.	N.A.	—	M.fr.II,142;	X,77		As., Afr.
N. gigantea, Hooker, Bot. Mag. t. 4647 (1852)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,61	M.fr.II,141;	X,78		P.

NELUMBO; Tournefort, inst. 261 (1700). (Nelumbium.)

N. nucifera; Gaertner, de fruct. I, 73 (1788)	—	—	—	—	—	Q.	N.A.	B.fl.I,62	M.fr.X,77	...	E., As., Afr.

PIPERACEAE.

L. CL. RICHARD in Humb. Bonpl. & Kunth, Nov. gen. I, 46 (1815).—Genera 2, Species 10.

PIPER; Linné, gen. pl. 333 (1737), from C. Bauhin (1623). (Macropiper.)

P. subpeltatum; Willdenow, spec. plant. V. 166 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,204	M.fr.V,198	...	As., Afr., Am., P.
P. excelsum; G. Forster, florul. ins. Austr. prodr. 5 (1786)	—	—	—	—	N.S.W.	—	—	B.fl.VI,204	M.fr.IX,77	...	N.Z., P.
P. Novae Hollandiae; Miquel in Medd. Akad. Vetensk. Amsterd. sec. ser. II, 5 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,204	M.fr.V,198.		
P. Banksii; Miquel in Medd. Akad. Amst. sec. ser. II, 9 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,205			
P. triandrum; F. v. M., fragm. V, 197 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,205	M.fr.V,197.		
P. hederaceum; Cunningham in De Cand. prodr. XVI, 366 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.VI,205			

PEPEROMIA; Ruiz & Pavon, fl. Peru. & Chil. prodr. 8, t. 2 (1794).

P. leptostachya; Hooker & Arnott, Bot. Beech. 96 (1841)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,206	M.fr.V,198;	IX,76	P.
P. reflexa; A. Dietrich, spec. plant. I, 180 (1831)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,207	M.fr.V,198;	IX,77	As., Afr., Am., P.
P. Urvilleana; Ach. Richard in d'Urville, voy. Bot. 356 (1832)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX.76	...	N.Z.
P. Baneriana; Miquel, syst. piperac. 120 (1843)	—	—	—	—	N.S.W.	—	—	—			

MAGNOLIACEAE.

J. DE ST. HILAIRE, Expos. Fam. nat. II, 74, t. 83-84 (1805).—Genus 1, Species 4.

DRIMYS; R. & G. Forster, char. gen. 83, t. 42 (1776). (Tasmannia.)

D. aromatica; F. v. M., pl. of Vict. I, 20 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,49	M.fr.VII,18.		
D. membranea; F. v. M., fragm. V, 175 (1866)	—	—	—	—	—	Q.	—	—	M.fr.VII,18.		
D. dipetala; F. v. M., pl. of Vict. I, 21 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,49	M.fr.VII,81.		
D. Howeana; F. v. M., fragm. VII, 17 (1870)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,76.		

ANONACEAE.

A. L. DE JUSSIEU, Gen. 283 (1789) from B. de Jussieu (1759).—Genera 11, Species 18.

UVARIA; Linné, Amoen. acad. 404 (1747).

U. membranacea; Bentham, Fl. Austr. I, 51 (1863)	—	—	—	—	—	Q.	—	B.fl.I,51			
U. Goezeana; F. v. M., fragm. VII, 125 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,125.		
U. Holtzei; F. v. M. in Wing's S. Sc. Rec., July 1883	—	—	—	—	—	—	N.A.	—			

FITZALANIA; F. v. M., fragm. IV, 33 (1863).

F. heteropetala; F. v. M., fragm. IV, 33 (1863)	—	—	—	—	—	Q.	—	B.fl.I,51	M.fr.III,1; VII,126.					

CANANGA; Rumphius, herb. Amboin. II, 195, t. 65 (1741). (Fitzgeraldia.)

C. odorata; J. Hooker & Thomson, fl. Indic. 130 (1855)	—	—	—	—	—	Q.	—	—	M.fr.VI,1; VIII,279.					

ANCANA; F. v. M., fragm. V, 27 (1865).

A. stenopetala; F. v. M., fragm. V, 27 (1865)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,189; VII,121.					

POLYALTHIA; Blume. Fl. Jav. Anon. 68 (1829).

P. nitidissima; Bentham Fl. Austr. I, 51 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,51	M.fr.III,2,167; VII,126.	P.				
P. Holtzeana; F. v. M. in Wing's S. Sc. Rec. II, 230 (1882)	—	—	—	—	—	—	N.A.	—						
P. Armitiana; F. v. M. in Austr. Journal of Pharmacy, Jan. 1887	—	—	—	—	—	Q.	—	—						

POPOWIA; Endlicher, Gen. 831 (1839).

P. Australis; Bentham, Fl. Austr. I, 52 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,52						

MELODORUM; Loureiro, Fl. Cochinch. I, 351 (1790).

M. Leichhardtii; Bentham, Fl. Austr. I, 52 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,52	M.fr.III,41,VI,2,176; VII,126.					
M. Uhrii; F. v. M., fragm. VI, 2 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VI,2.					
M. Maccreai; F. v. M., fragm. VI, 176 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,250.					

MITREPHORA; Spach, Veget. Phanerog. VII, 497 (1839).

M. Froggattii; F. v. M. in Austr. Journ. of Pharmacy II, 3 (1887)	—	—	—	—	—	Q.	—	—						

MILIUSIA; Leschenault in Mém. Genèv. V, 213 (1832).

Species undetermined	—	—	—	—	—	Q.	—	—						

SACCOPETALUM; Bennett in Horsf. pl. Jav. rar. 115, t. 35 (1838).

S. Bidwilli; Bentham, Fl. Austr. I, 53 (1863)	—	—	—	—	—	Q.	—	B.fl.I,53						
S. Brahei, F. v. M., fragm. VIII, 159 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,159.					

EUPOMATIA; R. Brown in App. Flind. voy. II, 597, t. 2 (1814).

E. Bennettii; F. v. M., fragm. 1, 45 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,54	M.fr.VI,177.					
E. laurina; R. Brown in App. Flind. voy. II, 597, t. 2 (1814)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,54	M.fr.I,45; VI,2; VII,126.					

MONIMIACEAE.

A. L. de Jussieu, in Ann. du Mus. XIV, 30 (1809).—Genera 7, Species 16.

DORYPHORA; Endlicher, Gen. 3015 (1837). (Learosa.)

Sassafras; Endlicher, iconogr. gen. pl., t. 10 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,283						

ATHEROSPERMA; Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806).

A. moschatum; Labillardière, Nov. Holl. pl. spec. II, 74, t. 224 (1806)... ...	—	—	T.	V.	N.S.W.	—	—	B.fl.V,284	M.fr.VIII,142; X,106.					

DAPHNANDRA; Bentham, Fl. Austr. V., 285 (1870).

D. micrantha; Bentham, Fl. Austr. V, 285 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,285						
D. repandula; F. v. M., fragm. X, 105 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,105.					
D. aromatica; Bailey, first suppl. to synop. Q.L. fl. 46 (1886)	—	—	—	—	—	Q.	—	—						

MOLLINEDIA; Ruiz & Pavon, prodr. 83, t. 15 (1794). (Kibara, Wilkiea.)

M. Huegeliana; Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1855)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,286						
M. Wardellii; F. v. M., fragm. V, 155 (1866)	—	—	—	—	—	Q.	—	B.fl.V,287	M.fr.VI,252.					
M. loxocarya; Bentham, Fl. Austr. V, 287 (1870)	—	—	—	—	—	Q.	—	B.fl.V,287						
M. acuminata; F. v. M., fragm. V, 155 (1866)	—	—	—	—	—	Q.	—	B.fl.V,287	M.fr.V,155.					
M. macrophylla; Tulasne in Ann. sc. nat. sér. quatr. III, 45 (1885)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,288	M.fr.V,3.					

M. longipes ; Bentham, Fl. Austr. V, 289 (1870)	—	—	—	—	—		Q.	—	B.fl.V,289						
M. laxiflora ; Bentham, Fl. Austr. V, 289 (1870)	—	—	—	—	—		Q.	—	B.fl.V,289						
M. pubescens ; Bentham, Fl. Austr. V, 290 (1870)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,290						
HEDYCARYA ; R. & G. Forster, char. gen. 127, t. 64 (1776).															
H. Cunninghami ; Tulasne in Archiv. du Mus. d'hist. nat. VIII, 408 (1855) ...	—	—	—	V.	N.S.W.		Q.	—	B.fl.V,291						
PALMERIA ; F. v. M., fragm. IV, 151 (1864).															
P. scandens ; F. v. M., fragm. IV, 152 (1864)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,291	M.fr.V,2.					
PIPTOCALYX ; Oliver in Bentham's Fl. Austr. V, 292 (1870).															
P. Moorei ; Oliver in Bentham's Fl. Austr. V, 292 (1870)	—	—	—	—	N.S.W.		—	—	B.fl.V,292	M.fr.X,106.					

MYRISTICEAE.

R. Brown, prodr. 399 (1810) from A. L. de Jussieu (1806).—Genus 1, Species 1.

MYRISTICA ; Linné, gen. pl. ed. II, 524 (1742).															
M. insipida ; R. Brown, prodr. 400 (1810)	—	—	—	—	—		Q.	N.A.	B.fl.V,281						

LAURACEAE.

Ventenat ; Tabl. I, 245 (1799).—Genera 7, Species 37.

CRYPTOCARYA ; R. Brown, prodr. 402 (1810). (Caryodaphne.)															
C. Murrayi ; F. v. M., fragm. V, 170 (1866)	—	—	—	—	—		Q.	—	B.fl.V,295	M.fr.X,253.					
C. Mackinnoniana ; F. v. M., fragm. V, 169 (1866)	—	—	—	—	—		Q.	—	B.fl.V,296	M.fr.V,169.					
C. patentinervis ; F. v. M. in De Candolle, prodr. XV, 508 (1864)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,296	M.fr.V,196.					
C. obovata ; R. Brown, prodr. 402 (1810)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,296	M.fr.V,166,170.					
C. glaucescens ; R. Brown, prodr. 402 (1810)...	—	—	—	—	N.S.W.		Q.	N.A.	B.fl.V,297	M.fr.V,165,166.					
C. triplinervis ; R. Brown, prodr. 402 (1810)...	—	—	—	—	N.S.W.		Q.	—	B.fl.V,297	M.fr.V,166.					
C. cinnamomifolia ; Bentham, Fl. Austr. V, 298 (1870)	—	—	—	—	—		Q.	—	B.fl.V,298						
C. Meissneri ; F. v. M., fragm. V, 170 (1866)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,298	M.fr.V,170.					
C. Australis ; Bentham, Fl. Austr. V, 299 (1870)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,299	M.fr.V,167.					
BEILSCHMIEDIA ; Nees in Wallich, pl. Asiat. rarior. II, 69 (1831). (Nesodaphne.)															
B. obtusifolia ; Bentham in B. & H. gen. III, 1521 (1880)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,299	M.fr.V,166.					
ENDIANDRA ; R. Brown, prodr. 402 (1810).															
E. glauca ; R. Brown, prodr. 402 (1810)	—	—	—	—	—		Q.	—	B.fl.V,300	M.fr.V,166.					
E. hypotephra ; F. v. M., fragm. V, 166 (1866)	—	—	—	—	—		Q.	—	B.fl.V,301	M.fr.V,166.					
E. discolor ; Bentham, Fl. Austr. V, 301 (1870)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,301						
E. Sieberi ; Nees, Syst. Laurin. 194 (1836)	—	—	—	—	N.S.W.		—	—	B.fl.V,301	M.fr.V,166.					
E. virens ; F. v. M., fragm. II, 90 (1860)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,302	M.fr.V,166.					
E. Muelleri ; Meissner in De Candolle, prodr. XV, 509 (1864)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,302						
E. pubens ; Meissner in De Candolle, prodr. XV, 509 (1864)	—	—	—	—	N.S.W.		Q.	—	B.fl.V,302						
CINNAMOMUM ; Burmann, fl. Zeil. 62 (1737).															
C. Tamala ; C. G. & T. F. Nees, Syst. Laurin. 56 (1836)	—	—	—	—	—		Q.	—	B.fl.V,303	M.fr.V,165	...	As.			
LITSEA ; Lamarck, Diction. III, 574 (1789). (Glabraria 1771, Tetranthera, Cylicodaphne, Litsaea).															
L. Chinensis ; Lamarck, Diction. III, 574 (1789)	—	—	—	—	—		Q.	N.A.	B.fl.V,305	...	...	As.			
L. Bindoniana ; F. v M., syst. Census of Austr. plants 4 (1882)	—	—	—	—	—		Q.	—	B.fl.V,305	M.fr.V,167.					
L. Hexanthus ; A. L. de Jussieu in Annal. du Mus. VI, 212 (1805)	—	—	—	—	—		Q.	—	B.fl.V,305						

L. reticulata ; Bentham in B. & J. H. gen. pl. III, part I, 161 (1880)	—	—	—	—	—	Q.	—	B.fl.V,306						
L. Zeylanica ; C. G. & T. F. Nees, in Amoen. Bonn. I, 58, t. 5 (1823)	—	—	—	—	—	Q.	—	B.fl.V,307			...	...	As.	
L. dealbata ; Nees, Syst. Laurin. 630 (1836)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,307	M.fr.V,166.					
CASSYTHA ; Osbeck in Linné, sp. pl. 35 (1753).														
C. nodiflora ; Meissner in De Candolle, prodr. XV, 252 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,309						
C. glabella ; R. Brown, prodr. 404 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,309	M.fr.V,167.					
C. flava ; Nees in Lehmann, pl. Preiss. I, 620 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,310						
C. pubescens ; R. Brown, prodr. 404 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,310	M.fr.V,167.					
C. phaeolasia ; F. v. M., fragm. V. 167 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.V,310						
C. paniculata ; R. Brown, prodr. 404 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,311						
C. filiformis ; Linné, sp. pl. 35 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.V,311	M.fr.V,167		...		As., Afr., Am.	
C. melantha ; R. Brown, prodr. 404 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,311	M.fr.V,167.					
C. micrantha ; Meissner in De Candolle, prodr. XV, 256 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,312						
C. racemosa ; Nees in Lehmann, pl. Preiss. I, 621 (1845)	W.A.	—	—	—	—	Q.	—	B.fl.V,312						
C. pomiformis ; Nees in Lehmann, pl. Preiss. I, 620 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,313						
HERNANDIA ; Plumier, Gen. 6, t. 40 (1703).														
H. peltata ; Meissner in De Candolle, prodr. XV, 263 (1864)	—	—	—	—	—	Q.	—	B.fl.V,314			...		As., Afr., P.	
H. bivalvis ; Bentham, Fl. Austr. V, 314 (1870)	—	—	—	—	—	Q.	—	B.fl.V,314						

MENISPERMEAE.

A. L. de Jussieu, Gen. 284 (1789).—Genera 15, Species 17.

TINOSPORA; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 35 (1851).														
T. smilacina ; Bentham in Journ. Linn. Soc. V, suppl. II, 2 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.I,55	M.fr.IX,82.					
T. Walcottii ; F. v. M. in Bentham's Fl. Austr. I, 56 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,56	M.fr.IX,83.					
FAWCETTIA ; F. v. M., fragm. X, 93 (1877).														
F. tinosporoides ; F. v. M., fragm. X, 93 (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,136.					
COCCULUS ; De Candolle, sys. veg. I, 515 (1818) from C. Bauhin (1623). (Pericampylus, Legnephora.)														
C. Moorei ; F. v. M., fragm. I, 162 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,56	M.fr.IX,81,197.					
HYPSERPA ; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 36 (1851). (Selwynia.)														
H. Selwyni ; F. v. M., fragm. IX, 82 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IV,153.					
ADELIOPSIS ; Bentham & J. Hooker, Gen. pl. I, 436 (1862).														
A. decumbens ; Bentham, Fl. Austr. I, 59 (1863)	—	—	—	—	—	Q.	—	B.fl.I,59	M.fr.IX,83					
TRISTICHOCALYX ; F. v. M., fragm. IV, 27 (1863).														
T. pubescens ; F. v. M., fragm. IV, 27 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,58	M.fr.IV,27.					
T. diffusus ; Miers, contrib. to Bot. III, 286 (1871)	—	—	—	—	—	Q.	—	—						
SARCOPETALUM ; F. v. M., pl. of Vict. I, 26 (1860).														
S. Harveyanum ; F. v. M., pl. of Vict. 27 and 221 (1860)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,57	M.fr.IX,83.					
LEICHHARDTIA ; F. v. M., fragm. X, 67 (1876).														
L. clamboides ; F. v. M., fragm. X, 68 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,68.					
STEPHANIA ; Loureiro, Fl. Cochinch. II, 608 (1790).														
S. hernandifolia ; Walpers, repert. bot. syst. I, 96 (1842)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.I,57	M.fr.IX,76 ; X,94				As., Afr.	
CISSAMPELOS ; Linné, gen. plant. 368 (1737).														
C. Pareira ; Linné, spec. plant. 1031 (1753)	—	—	—	—	—	Q.	—	—					As., Afr., Am.	

PACHYGONE; Miers in Ann. and Mag. nat. hist. sec. ser. VII, 37 (1851).
P. Hullsii; F. v. M., fragm. IX, 81 (1875) ... ... — — — — — Q. N.A. — M.fr.V,147.

PYCNARRHENA; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851).
P. Australiana; F. v. M. in Vict. Naturalist III, 61 (1886)... ... ... — — — — — Q. — —

PLEOGYNE; Miers in Taylor's Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851). (Microclisia.)
P. Cunninghami; Miers in Ann. and Mag. of nat. hist. sec. ser. VII, 37 (1851)... — — — — — Q. — B.fl.I,59 M.fr.IX,82.

HUSEMANNIA; F. v. M. in Wing's S. Sc. Rec. III, 187 (1883).
H. protensa; F. v. M. in Wing's S. Sc. Rec. III, 127 (1883) ... ... — — — — — Q. — —

CARRONIA; F. v. M., fragm. IX, 84 (1875).
C. multisepalea; F. v. M., fragm. IX, 171 (1875) ... ... — — — — N.S.W. — — — M.fr.IX,84.

PAPAVERACEAE.

A. L. de Jussieu; Gen. 235 (1789) from B. de Jussieu (1759).—Genus 1, Species 1.

PAPAVER; Tournefort, inst. 237, t. 119 (1700) from Plinius.
P. aculeatum; Thunberg, Fl. Capensis 431 (1813) ... ... — S.A. T. V. N.S.W. Q. — B.fl.I,63 ... ... Afr.

CAPPARIDEAE.

Ventenat; Tabl. III, 118 (799).—Genera 6, Species 29.

CLEOME; Linné, syst. nat. 9 (1735); Linné, gen. 200 (1737). (Polanisia.)
C. oxalidea; F. v. M., fragm, I, 69 (1858) ... ... — — — — — Q. N.A. B.fl.I,90 M.fr.I,69
C. tetrandra; Banks in De Candolle, prodr. I, 240 (1824) ... ... W.A. — — — — Q. N.A. B.fl.I,90 M.fr.I,70,VII,40; IX,174.
C. viscosa; Linné, sp. pl. 672 (1753) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.I,90 M.fr.I,70,IX,174. As., Afr.
C. grandiflora; F. v. M., ined. ... ... — — — — — Q. N.A. —

ROEPERIA; F. v. M. in Hooker's Kew Misc. IX, 15 (1857). (Tetratelia.)
R. cleomoides; F. v. M. in Hooker's Kew Misc. IX, 15 (1857) ... ... — — — — — Q. N.A. B.fl.I,91 M.fr.V,105; IX,174.

EMBLINGIA; F. v. M., fragm. II, 2 (1860).
E. calceoliflora; F. v. M., fragm. II, 3 (1860)... ... ... W.A. — — — — — — B.fl.I,92 M.fr.; XI,20.

CADABA; Forskal, Fl. Aeg. Arab. 67 (1775).
C. capparoides; De Candolle, prodr. I, 244 (1824) ... ... — — — — — — N.A. B.fl.I,92 ... ... As.

CAPPARIS; Tournefort, inst. 261, t. 139 (1700) from Theophrastos, Dioscorides and Plinius. (Busbeckea, Busbecquea.)
C. umbellata; R. Brown in De Candolle, prodr. I, 247 (1824) ... ... — — — — — Q. N.A. B.fl.I,93 M.fr.V,104; IX,172.
C. lasiantha; R. Brown in De Candolle, prodr. I, 247 (1824) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.I,94 M.fr.IX,173.
C. quiniflora; De Candolle, prodr. I, 247 (1824) ... ... — — — — — Q. N.A. B.fl.I,94 M.fr.IX,173. ... P.
C. spinosa; Linné, sp. pl. 503 (1753) ... ... W.A. S.A. — — — Q. N.A. B.fl.I,94 M.fr.I,143; V,104; IX,173. E., As., Afr., P.
C. sarmentosa; Cunningham in Bentham's Fl. Austr. I, 95 (1863) ... .. — — — — — Q. — B.fl.I,95 M.fr.V,104; IX,173.
C. ornans; F. v. M. in Bentham's Fl. Austr. I, 95 (1863) ... ... — — — — — Q. — B.fl.I,95 M.fr.V,104; IX,173.
C. nobilis; F. v. M. in Bentham's Fl. Austr. I, 95 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,95 M.fr.I,163; V,104.
C. canescens; Banks in De Candolle, prodr. I, 246 (1824) ... ... — — — — — Q. — B.fl.I,96 M.fr.V,104; IX,173.
C. Shanesiana; F. v. M., fragm. X, 94 (1877) ... ... — — — — — Q. — — M.fr.X,94.
C. lucida; R. Brown in Bentham's Fl. Austr. I, 96 (1863) ... ... — — — — — Q. — B.fl.I,96 M.fr.I,163; V,104; IX,173.
C. Mitchelli; Lindley in Mitchell's Three Exped. I, 315 (1838) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.I,96 M.fr.IX,173.
C. Ioranthifolia; Lindley in Mitchell's Trop. Austr. 220 (1848) ... ... — — — — N.S.W. Q. N.A. B.fl.I,97

C. umbonata; Lindley in Mitchell's Trop. Austr. 257 (1848) ... ... — — — — — Q. N.A. B.fl.I,97 M.fr.IX,173
C. uberiflora; F. v. M., fragm. IX, 172 (1875) ... ... — — — — — Q. — — M.fr.IX,172.
C. humistrata; F. v. M., fragm. V, 156 (1866) ... ... — — — — — Q. — — M.fr.V,156.
C. Thozetiana; F. v. M., frsgm. V., 104 (1866) ... ... — — — — — Q. — — M.fr.V,156.

APOPHYLLUM; F. v. M. in Hooker's Kew Misc. IX, 307 (1857).

A. anomalum; F. v. M. in Hooker's Kew Misc. IX. 307 (1857) ... ... — — — — N.S.W. Q. N.A. B.fl.I,97 M.fr.IX,173; XI,27.

CRUCIFERE.

A. L. DE JUSSIEU, Gen. 237 (1789) from B. de Jussieu (1759).—Genera 14, Species 54.

NASTURTIUM; R. Brown in W. T. Aiton's hortus Kewensis IV, 110 (1812) from Linné, syst. nat. 9 (1735).

N. terrestre; R. Brown in Aiton's hort. Kewens. IX, 110 (1812) ... ... — S.A. T. V. N.S.W. Q. — B.fl.I,66 M.fr.XI,60. ... E., As., Afr., Am., P., N.Z.

BARBARAEA; Beckmann, Lex. botan. 33 (1801). (Barbarea, from L'Obel 1576.)

B. vulgaris; R. Brown in De Candolle, prodr. I, 140 (1824) ... ... — — T. V. N.S.W. — — B.fl.I,66 M.fr.VIII,142; XI,6,60 E., As., Am., N.Z.

ARABIS; Linné, gen. pl. 198 (1737) from Dalechamps (1554). (Turritis.)

A. glabra; Crantz, stirp. Austriac. 36 (1767) ... ... — — — V. N.S.W. — — B.fl.I,67 ... ... E., As., Am.

CARDAMINE; Tournefort, inst. 224, t. 109 (1700) from l'Ecluse (1576).

C. stylosa; De Candolle, regn. veg. syst. nat. II, 24 (1821) ... ... — — T. V. N.S.W. Q. — B.fl.I,68 M.fr.VII,19; XI,60 ... N.Z.
C. dictyosperma; Hooker, Journ. of Bot. I, 246 (1834) ... ... — — T. V. N.S.W. — — B.fl.I,68 M.fr.VII,19; X,6,60.
C. radicata; J. Hooker in icones pl. t. 882 (1852) ... ... — — T. — — — — B.fl.I,69 M.fr.XI,6.
C. laciniata; F. v. M. in Transact. Phil. Soc. Vict. I, 34 (1854) ... ... — S.A. — V. N.S.W. — — B.fl.I,69 M.fr.XI,27,60.
C. hirsuta; Linné, sp. pl. 655 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.I,70 M.fr.VII,19; XI,60 E., As., Am., N.Z., P.
C. eustylis; F. v. M. in Transact. Vict. Inst. I, 114 (1854)... ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.I,71 M.fr.VII,19; XI,60.

WILCKIA; Scopoli, introd. hist. nat. 317 (1777). (Malcolmia.)

W. Africana; F. v. M., native plants of Vict. I, 33 (1879) ... ... — — — V. N.S.W. — — — M.fr.VII,18 ... E., As., Afr.

SISYMBRIUM; Tournefort, inst. 225, t. 109 (1700). (Blennodia.)

S. filifolium; F. v. M. in Transact. Phil. Soc. Vict. I, 34 (1854) ... ... — S.A. — — N.S.W. — — B.fl.I,73
S. trisectum; F. v. M. in Transact. Vict. Inst. I, 114 (1854) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.I,74 M.fr.VII,20; XI,60.
S. nasturtioides; F. v. M. in Transact. Vict. Inst. I, 115 (1854) ... ... — S.A. — V. N.S.W. Q. — B.fl.I,74 M.fr.X,53.
S. Richardsii; F. v. M., fragm. X, 105 (1877) ... ... W.A. S.A. — — — — — — M.fr.X,105.
S. eremigenum; F. v. M., fragm. II, 143 (1861) ... ... — — — — N.S.W. Q. — B.fl.I,74 M.fr.II,143.
S. Lucae; F. v. M., syst. Cens. of Austr. pl. 5 (1882) ... ... — — — V. N.S.W. — — — M.fr.XI,59.
S. cardaminoides; F. v. M. in Transact. Phil. Soc. Vict. I, 34 (1854) ... ... — S.A. — V. N.S.W. Q. — B.fl.I,75 M.fr.VII,19; VII,223; X,53; XI,6,27,59.

ERYSIMUM; Linné, gen. pl. 198 (1737). (Blennodia, Alyssopsis.)

E. curvipes; F. v. M. in Linnaea XXV, 368 (1852) ... ... — S.A. — V. N.S.W. — — B.fl.I,75 M.fr.VII,20.
E. brevipes; F. v. M. in Linnaea XXV, 367 (1852) ... ... W.A. S.A. — V. N.S.W. — — B.fl.I,75 M.fr.VII,20; X,53,119; XI,6.
E. blennodioides; F. v. M. in Linnaea XXV, 367 (1852) ... ... — S.A. — V. N.S.W. Q. — B.fl.I,76 M.fr.VII,20; XI, 27,60.
E. Cunninghamii; F. v. M. Cens. 5 (1882) from Bentham (1863)... ... ... — — — — N.S.W. Q. — B.fl.I,76
E. Blennodia; F. v. M., fragm. X, 78 (1877) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.I,76 M.fr.VII,20.
E. capsellinum; F. v. M., native pl. of Vict. I, 35 (1879) ... ... — — — V. N.S.W. — — B.fl.I,77

ALYSSUM; Tournefort, inst. 217 (1700) from l'Ecluse (1576). (Meniocus, Alysson.)

A. minimum; Pallas, Reise Prov. Russ. Reich. III, 746 (1776) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.I,71 M.fr.XI,27,60 ... E., As., Afr.

STENOPETALUM; R. Brown in De Candolle, Mém. du Mus. VII, 239 (1821.)

S. velutinum; F. v. M., pl. Vict. I, 49 (1860) ... ... — S.A. — V. N.S.W. Q. — B.fl.I,78 M.fr.III,27; VII,20; XI,27.
S. lineare; R. Brown in De Candolle, syst. veg. I, 513 (1821) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.I,78 M.fr.VII,20; XI,6.

S. sphaerocarpum ; F. v. M. in Transact. Phil. Soc. Vict. I, 35 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,78	M.fr.VII,20.					
S. nutans ; F. v. M., fragm. III, 27 (1862)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,79	M.fr.VII,20 ; XI,6.					
S. robustum ; Endlicher in Huegel enum. 4 (1837)	W.A.	S.A.	—	—	—	—	—	B.fl.I,79	M.fr.VII,20 ; XI,5,60.					
S. pedicellare ; F. v. M. in Bentham, Fl. Austr. I, 79 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,79	M.fr.VII,20.					
S. croceum ; Bunge in Lehmann, pl. Preiss. I, 258 (1844)	W.A.	—	—	—	—	—	—	—	M.fr.XI,6.					
MENKEA ; Lehmann, Ind. sem. hort. bot. Hamb. 8 (1843).														
M. Australis ; Lehmann in Ind. sem. hort. bot. Hamb. 8 (1843)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,80	M.fr.II,142.					
M. draboides ; J. Hooker in Bentham Fl. Austr. I, 80 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,80	M.fr.II,142 ; VIII,223 ; XII,19.					
M. sphaerocarpa ; F. v. M., fragm. VIII, 223 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,223.					
CAPSELLA ; Medicus, Pflanzen-Gattungen 85 (1792). (Hutchinsia partly, Thlaspi partly, Microlepidium).														
C. elliptica ; C. A. Meyer, Verz. der. Pfl. v. Caucas. 194 (1831)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,81	M.fr.VII,19 ; XI,60	E., As., Afr., Am.				
C. antipoda ; F. v. M., pl. of Vict. I, 44 (1860)	—	—	T.	V.	—	—	—	B.fl.I,81						
C. pilosula ; F. v. M., pl. of Vict. I, 44 (1860)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,82	M.fr.XI,60.					
C. Tasmanica ; F. v. M., fragm. XI, 26)1878)	—	—	T.	—	—	—	—	B.fl.I,87						
C. cochlearina ; F. v. M., pl. of Vict. I, 51 (1860)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,88	M.fr.XI,26.					
C. Drummondii ; F. v. M., fragm. XI, 21 (1878)	W.A.	S.A.	—	—	—	—	—	B.fl.I,88	M.fr.VII,90 ; XI,26.					
C. humistrata ; F. v. M., fragm. XI, 25 (1878)	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.XI,25.					
C. Andraeana ; F. v. M. in Wing's S. Sc. Rec. March p. 49 (1885)	—	—	—	—	N.S.W.	—	—	—						
SENEBIERA ; De Candolle in Mém. soc. d'hist. nat. Par. I, 140 (1799). Carara, Medicus (1792) from Caesalpini (1583).														
S. integrifolia ; De Candolle in Mém soc. d'hist. nat. Par. I, 144 (1799)	—	—	—	—	—	Q.	—	B.fl.I,82	...	...	As., Afr.			
LEPIDIUM ; Tournefort, inst. 215, t. 103 (1700) from Dioscorides and Plinius. (Iberis partly, Lepia, Monoploca).														
L. strongylophyllum ; F. v. M. in Benth. Fl. Austr. I, 84 (1863)...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,84	M.fr.XI,6,60.					
L. linifolium ; Bentham, Fl. Austr. I, 84 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,84	M.fr.VII,20 ; XI,6.					
L. leptopetalum ; F. v. M., pl. of Vict. I, 48 (1860)...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,84						
L. rotundum ; De Candolle, syst. veg. II, 537 (1818)	W.A.	S.A.	—	—	—	—	—	B.fl.I,85	M.fr.VII,20 ; XI,6.					
L. phlebopetalum ; F. v. M., pl. of Vict. I, 47 (1860)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,85	M.fr.VII,19 ; XI,60.					
L. pedicellosum ; F. v. M., fragm. XI, 27 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.XI,27.					
L. monoplocoides ; F. v. M. in Trans. Phil. Soc. Vict. 35 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,85	M.fr.XI,27.					
L. papillosum ; F. v. M. in Linnaea XXV, 370 (1852)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,86	M.fr.VII,19 ; XI,6,28.					
L. foliosum ; Desvaux, Journ. de Bot. III, 164 et 180 (1813)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,86	M.fr.VII,19 ; IX,76.					
L. ruderale ; Linné, sp. pl. 645 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,86	M.fr.VII,19 ; XI,27,61.					
CAKILE ; Tournefort, coroll. 49, t. 483 (1700) from L'Obel (1576).														
C. maritima ; Scopoli, Fl. Carniolica, 344 (1760)	—	S.A.	T.	V.	N.S.W.	—	—	—	M.fr.VII,19 ; XI,6,28,60	E.,As.,Afr.				

VIOLACEAE.

De Candolle ; Flor. Franç. IV, 801 (1805).—Genera 4, Species 13.

VIOLA ; Tournefort, inst. 419, t. 236 (1700) from Plinius. (Erpetion.)														
V. betonicifolia ; Smith in Rees's Cyclopedia XXXVII (1817)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,99	M.fr.X,82.					
V. hederacea ; Labillardière, Nov. Holl. pl. spec. I, 66, t. 91 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,99	M.fr.X,82.					
V. Cunninghamii ; J. Hooker, Fl. N. Zel. I, 16 (1853)	—	—	T.	—	—	—	—	B.fl.I,100	...	...	N.Z.			
V. Caleyana ; G. Don, gen. syst. dichlam. pl. I, 329 (1831)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,100	M.fr.X,82.					
HYBANTHUS ; Jacquin, Stirp. Amer. hist. 77, t. 75 (1763). (Ionidium, Pigea, Vlamingia).														
H. floribundus ; F. v. M., nat. pl. of Vict. I, 45 (1879)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,112	M.fr.X,81.					
H. Vernonii ; F. v. M., nat. pl. of Vict. I, 45 (1879)	—	—	—	V.	N.S.W.	—	—	B.fl.I,103	M.fr.X,81.					
H. filiformis ; F. v. M., nat. pl. of Vict. I, 44 (1879)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,103	M.fr.X,81.					

H. debilissimus ; F. v. M., fragm. XI, 4 (1878)	W.A.	—	—	—	—	—	—	—		M.fr.X,4.						
H. Tatei ; F. v. M. in Trans. R. Soc. of S. Austr. IX, 102 (1882)	—	S.A.	—	—	—	—	—	—								
H. calycinus ; F. v. M., fragm. X, 81 (1877)	W.A.	—	—	—	—	—	—	B.fl.I,104		M.fr.XI,4.						
H. ennesspermus ; F. v. M., fragm. X, 81 (1877)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,101		M.fr.X.81	...	As., Afr.				
HYMENANTHERA ; R. Brown in Tuck. Cong. 442 (1818).																
H. Banksii ; F. v. M., first gen. report 9 (1853)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,104		M.fr.IX,77 ; X.82		N.Z.				
MELICYTUS ; R. & G. Forster, char. gen. 123, t. 62 (1776).																
M. ramiflorus ; R. & G. Forst., char. gen. 123, t. 2 (1776)	—	—	—	—	N.S.W.	—	—	—		...	...	...	N.Z.			

FLACOURTIEAE.

Cl. Richard in Mém. du Mus. I, 366 (1815).—Genera 4, Species 7.

COCHLOSPERMUM ; Kunth, Malvac. 6 (1822).																
C. Fraseri ; Planchon in Hook. Lond. Journ. VI, 307 (1847)	—	—	—	—	—	—	N.A.	B.fl.I,106		M.fr.IX,83.						
C. Gillivraei ; Bentham, Fl. Austr. I, 106 (1863)	—	—	—	—	—	Q.	—	B.fl.I,106								
C. Gregorii ; F. v. M., fragm. I, 71 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,106		M.fr.IX,61.						
SCOLOPIA ; Schreber, Gen. 355 (1789). (Phoberos).																
S. Brownii ; F. v. M., fragm. III, 11 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,107		M.fr.IV,174 ; IX,61.						
XYLOSMA ; G. Forster, prodr. 380 (1786).																
X. ovata ; Bentham, Fl. Austr. I, 108 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,108		M.fr.IX,60.						
STREPTOTHAMNUS ; F. v. M., fragm. III, 28 (1862).																
S. Moorei ; F. v. M., fragm. III, 28 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,108		M.fr.III,28.						
S. Beckleri ; F. v. M., fragm. III, 28 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,108		M.fr.IX,61.						

SAMYDACEAE.

J. Gaertner ; de Fruct. III, 328 (1805).—Genera 2, Species 4.

CASEARIA ; Jacquin, Stirp. Amer. hist. 132, t. 85 (1763).																
C. esculenta ; Roxburgh. Fl. Ind. II, 422 (1832)	—	—	—	—	—	Q.	—	B.fl.VI,309	...	...	As.					
C. Dallachii ; F. v. M., fragm. V, 107 (1866)	—	—	—	—	—	Q.	—	B.fl.III,309	M.fr.V,214 ; VI,252.							
HOMALIUM ; Jacquin, Stirp. Amer. hist. 170, t. 183 (1763). (Blackwellia.)																
H. Vitiense ; Bentham in Journ. Linn. Soc. IV, 36 (1860)	—	—	—	—	—	Q.	—	B.fl.III,310	...	...	P.					
H. brachybotrys ; F. v. M., fragm. II, 127 (1861)	—	—	—	—	—	Q.	—	B.fl.III,310	M.fr.II,127.							

PITTOSPOREAE.

R. Brown in Flinders' voy. 542 (1814).—Genera 8, Species 40.

PITTOSPORUM ; Banks in Gaertner, de Fruct. I, 286, t. 59 (1788).																
P. rhombifolium ; Cunningham in Hook. icon. pl. t. 621 (1844)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,110		M.fr.XII,4.						
P. melanospermum ; F. v. M., fragm. I, 70 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,111		M.fr.VI,167 ; XII,4.						
P. undulatum ; Andrews, Bot. reposit. 383 (1800)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,111		M.fr.VI,167 ; XII,4.						
P. revolutum ; Aiton, Hort. Kew, sec. ed. II, 27 (1811)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,111		M.fr.VI,167,186 ; VII,140 ; XII,4.						
P. venulosum ; F. v. M., fragm. VI, 186 (1868)	—	—	—	—	—	Q.	—	—		M.fr.VII,140 ; XII,4.						
P. ferrugineum ; Aiton, Hort. Kew, sec. ed. II, 27 (1811)	—	—	—	—	—	Q.	—	B.fl.I,112		M.fr.II,78 ; III,165 ; VI,167 ; XII,4.						
P. bracteolatum ; Endlicher, prodr. fl. ins. Norf. 78 (1833)	—	—	—	—	N.S.W.	—	—	—	...	...	...	...	As., P.			

P. Wingii; F. v. M. in Wing's S. Sc. Rec. March (1885), p. 49	—	—	—	—	—	Q.	—	—		
P. rubiginosum; Cunningham in Ann. of Nat. Hist. IV, 108 (1842)	—	—	—	—	—	Q.	—	B.fl.I,112	M.fr.VI,167; VII,140; IX,190.	
P. phillyroides; De Candolle, prodr. I, 347 (1824)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,112	M.fr.VI,167; IX,75.	
P. bicolor; Hooker, Journ. Bot. I, 249 (1834)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,113	M.fr.VIII,142; XII,4.	
P. erioloma; C. Moore et F. v. M., fragm. VII, 139 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,139.	
HYMENOSPORUM; R. Brown in Hook. Bot. Magaz. 4799 (1854).										
H. flavum; F. v. M., fragm. II, 77 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I.114	M.fr.V,210; VI,168; VII,140; XII,4.	
BURSARIA; Cavanilles, icon. et descr. pl. IV, 30, t. 350 (1797).										
B. spinosa; Cavanilles, icon. pl. IV, 30, t. 350 (1797)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,115	M.fr.XII,4.	
MARIANTHUS; Huegel, enum. pl. Nov. Holl. 8 (1837). (Oncosporum, Rhytidosporum, Calopetalon, Spiranthera.)										
M. procumbens; Bentham, Fl. Austr. I, 117 (1863)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,117	M.fr.XII,2.	
M. rhytidosporeus; F. v. M., fragm. II, 145 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,117	M.fr.II,145.	
M. villosus; Bentham, Fl. Austr. I, 117 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,117		
M. parviflorus; F. v. M., fragm. II, 144 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,118	M.fr.II,144; XII,2.	
M. bignoniaceus; F. v. M. in Trans. Phil. Soc. Vict. 6 (1854)	—	S.A.	—	V.	—	—	—	B.fl.I,118	M.fr.XII,2.	
M. Drummondianus; Bentham, Fl. Austr. I, 119 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,119	M.fr.XII,2.	
M. coeruleo-punctatus; Klotzsch in Link and Otto, icon. pl. 28, t. 12 (1841) ...	W.A.	—	—	—	—	—	—	B.fl.I,119	M.fr.XII,2.	
M. candidus; Huegel, enum. pl. Nov. Holl. 8 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,120	M.fr.XII,2.	
M. erubescens; Putterlick in Endl. nov. stirp. dec. 60 (1839)	W.A.	—	—	—	—	—	—	B.fl.I,120	M.fr.I,219; XII,2.	
M. ringens; F. v. M., fragm. I, 218 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,120	M.fr.I,218; XII,2.	
M. pictus; Lindley in Bot. Regist. XXV, App. XXV (1839)	W.A.	—	—	—	—	—	—	B.fl.I,121	M.fr.I,217; II,78,182; XII,2.	
CITRIOBATUS; Cunningham in Loudon's hort. Brit. Suppl. I, 585 (1832). (Ixiosporum.)										
C. multiflorus; Cunningham in Loud. hort. Brit. I, 585 (1832)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,121	M.fr.II,77; VI,168; XII,4.	
C. pauciflorus; Cunningham in Loud. hort. Brit. 1, 585 (1832)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,122	M.fr.II,76; VI,168; VII,180.	
BILLARDIERA; Smith, Sp. Bot. New. Holl. I, t. 1 (1793). (Pronaya, Campylanthera.)										
B. longiflora; Labillardière, Nov. Holl. plant. spec. I, 64, t. 89 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,123		
B. scandens; Smith, Spec. bot. New Holl. I, t. 1 (1793)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,123	M.fr.XI,123.	
B. coriacea; Bentham, Fl. Austr. I, 124 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,124	M.fr.XII,1.	
B. cymosa; F. v. M. in Transact. Vict. Inst. I, 29 (1855)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,124	M.fr.VI,168; VII,140.	
B. elegans; F. v. M., pl. of Vict. I, 78 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,125	M.fr.XII,2.	
B. variifolia; De Candolle, prodr. I, 346 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,125	M.fr.II,145.	
B. Lehmanniana; F. v. M., pl. of Vict. I, 78 (1860)...	W.A.	—	—	—	—	—	—	B.fl.I,125	M.fr.XII,2.	
B. floribunda; F. v. M. in South. Sc. Record II, 1 (1882)	W.A.	—	—	—	—	—	—	B.fl.I,120	M.fr.XII,1.	
SOLLYA; Lindley, Bot. Regist. XVII, t. 1460 (1831). (Xerosollya.)										
S. heterophylla; Lindley, Bot. Regist. XVII, t. 1466 (1831)	W.A.	—	—	—	—	—	—	B.fl.I,126	M.fr.XII,3.	
S. parviflora; Turcz. in Bull. Soc. Mosc. XXVII, 361 (1854)	W.A.	—	—	—	—	—	—	B.fl.I,126	M.fr.XII,3.	
CHEIRANTHERA; Cunningham in Brongn. Bot. voy. Coquille, t. 77 (1829).										
C. linearis; Cunningham in Bot. Regist. 1719 (1834)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,127	M.fr.I,97; II,180; VI,168; IX,168; XII,4.	
C. filifolia; Turczaninow in Bull. Soc. Mosc. XXVII, 364 (1854)	W.A.	—	—	—	—	—	—	B.fl.I,127	M.fr.I,97; II,79; VI,168; IX,168; XII,4.	
C. parviflora; Bentham, Fl. Austr. I, 128 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,128	M.fr.XII,3.	

DROSERACEAE.

SALISBURY; Parad. Lond. 95 (1809).—Genera 3, Species 46.

ALDROVANDA; Monti in Act. Bonon. II, 404, t. 12 (1747).										
A. vesiculosa; Linné, sp. pl. 281 (1753)	—	—	—	—	—	Q.	—	—	M.fr.VI,104;X,79 ... E., As.	

DROSERA ; Linné, gen. pl. 89 (1773). (Sondera).

D. Indica ; Linné, sp. pl. 232 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,456	M.fr.X,79	...	As., Afr.
D. Adelae ; F. v. M., fragm. IV, 154, t. XXXIII (1864)	—	—	—	—	—	Q.	—	—	M.fr.X,80.		
D. Arcturi ; Hooker, Journ. of Bot. I, 247 (1834)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,456	...	...	N.Z.
D. glanduligera ; Lehmann, pugill. VIII, 37 (1844)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,457	M.fr.X,80.		
D. pygmaea ; De Candolle, prodr. I, 317 (1824)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,457	M.fr.X,80	...	N.Z.
D. platystigma ; Lehmann, pugill. VIII, 37 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,457	M.fr.X,80.		
D. pulchella ; Lehmann, pugill. VIII, 38 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,458	M.fr.X,80.		
D. leucoblasta ; Bentham, Fl. Austr. II, 458 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,458	M.fr.X,80.		
D. nitidula ; Planchon in Ann. des sc. nat. IX, 285 (1848)	W.A.	—	—	—	—	—	—	B.fl.II,458			
D. paleacea ; R. Brown in De Candolle, prodr. I, 318 (1824)	W.A.	—	—	—	—	—	—	B.fl.II,458			
D. parvula ; Planchon in Ann. des sc. nat. IX, 287 (1848)	W.A.	—	—	—	—	—	—	B.fl.II,459			
D. Burmanni ; Vahl, Symbol. III, 50 (1794)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,459	M.fr.X,80.	...	As.
D. spathulata ; Labillardière, Nov. Holl. pl. spec. I, 79, t. 106 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,459	...	...	As., N.Z.
D. Drummondii ; Lehmann, pl. Preiss. II. 235 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,460			
D. scorpioides ; Planchon in Ann. des sc. nat. IX, 288 (1848)	W.A.	—	—	—	—	—	—	B.fl.II,460	M.fr.VIII,185 ; X,85.		
D. petiolaris ; R. Brown in De Candolle, prodr. I, 318 (1824)	—	—	—	—	—	Q.	N.A.	B.fl.II,460	M.fr.X,80.		
D. binata ; Labillardière, Nov. Holl. pl. spec. I, 78, t. 105 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,461	M.fr.X,80	...	N.Z.
D. zonaria ; Planchon in Ann. des sc. nat. IX, 303 (1848)	W.A.	—	—	—	—	—	—	B.fl.II,462			
D. bulbosa ; Hooker, icon. pl. t. 375 (1841)	W.A.	—	—	—	—	—	—	B.fl.II,462			
D. rosulata ; Lehmann, pugill. VIII, 36 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,462	M.fr.X,80.		
D. Whittakerii ; Planchon in Ann. des sc. nat. IX, 302 (1848)	—	S.A.	—	V.	—	—	—	B.fl.II,462			
D. macrophylla ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A.	—	—	—	—	—	—	B.fl.II,463			
D. squamosa ; Bentham, Fl. Austr. II, 463 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,463			
D. erythrorrhiza ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A.	—	—	—	—	—	—	B.fl.II,463			
D. stolonifera ; Endlicher in Hueg. enum. 5 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,463			
D. humilis ; Planchon in Ann. des sc. nat. IX, 300 (1848)	W.A.	—	—	—	—	—	—	B.fl.II,464			
D. ramellosa ; Lehmann, pugill. VIII, 40 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,464			
D. platypoda ; Turczaninow in Bull. de l'Ac. de St. Petersb. 343 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.II,464	M.fr.VIII,185 ; X,80.		
D. auriculata ; Backhouse in Ann. des sc. nat. IX, 295 (1848)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,465	M.fr.X,80	...	N.Z.
D. lunata ; Buchanan in De Candolle, prodr. I, 319 (1824)	—	—	—	—	—	Q.	—	—	...	...	As.
D. peltata ; Smith in Willd. spec. pl. I, 1546 (1797)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,465	M.fr.X,80.		
D. Neesii ; Lehmann, pugill. VIII, 42 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,466			
D. gigantea ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A.	—	—	—	—	—	—	B.fl.II,466			
D. myriantha ; Planchon in Ann. des sc. nat. IX, 291 (1848)	W.A.	—	—	—	—	—	—	B.fl.II,466			
D. pallida ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A.	—	—	—	—	—	—	B.fl.II,467			
D. penicillaris ; Bentham, Fl. Austr. II, 467 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,467			
D. filicaulis ; Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,467			
D. Huegelii ; Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,467	M.fr.VIII,185;X,81.		
D. macrantha ; Endlicher in Hueg. enum. 6 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,468			
D. Menziesii ; R. Brown in De Candolle, prodr. I, 319 (1824)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,468			
D. calycina ; Planchon in Ann. des sc. nat. IX, 299 (1848)...	W.A.	—	—	—	—	—	—	B.fl.II,468			
D. heterophylla ; Lindley, Bot. Regist. XXV, App. XX (1839)	W.A.	—	—	—	—	—	—	B.fl.II,469			
D. Banksii ; R. Brown in De Candolle, prodr. I, 319 (1824)	—	—	—	—	—	Q.	—	B.fl.II,469			

BYBLIS ; Salisbury Parad. Lond. t. 95 (1808).

B. liniflora ; Salisbury, Parad. Lond. t. 95 (1808)	W.A.	—	—	—	—	Q.	N.A.	B.fl.II,470	M.fr.X,81.		
B. gigantea ; Lindley Bot. Regist. XXV, App. XXI (1839)	W.A.	—	—	—	—	...	...	B.fl.II,470			

ELATINEAE.

Cambessedes in Mém. du Mus. XVIII 225 (1829).—Genera 2, Species 4.

ELATINE; Linné, gen. pl., 18 (1737).											
E. Americana; Arnott in Edinb. journ. nat. sc. I, 431 (1830)	W.A.	S.A.	T.	V.	N.S.W.	Q.	...	B.fl.I,178	M.fr.XI,27	...	Am., P., N.Z.
BERGIA; Linné, mantiss. II, 152 (1771).											
B. ammannioides; Roxburgh, hort. Bengh. 34 (1814)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,180	M.fr.II,147; IV,173		As., Afr
B. pedicellaris; F. v. M. in Benth. Fl. Austr. I, 180 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,180	M.fr.II,145; IV,173.		
B. perennis; F. v. M. in Benth. Fl. Austr. I, 180 (1863)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,181	M.fr.II,146.		

HYPERICINAE.

J. de St. Hilaire, Expos. Fam. II, 23, t. 75 (1805).—Genus 1, Species 1.

HYPERICUM; Tournefort, inst. 254, t. 131 (1700), from Dioscorides and Plinius. (Ascyrum, Brathys).											
H. Japonicum; Thunberg, Fl. Jap. 295, t. 31 (1784)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,182	...	...	As., N.Z.

TERNSTROEMIACEAE.

Mirbel in Bullet. Soc. Philomat. Dec. p. 381 (1813).—Genus 1, Species 1.

SARAUIA; Willdenow in Schrift. nat. Freunde, Berlin III, 406, t. 4 (1801).											
S. Andraeana; Dyer & Oliver in Bailey, synops. Queensl. flora, suppl. (1885)	—	—	—	—	—	Q.	—	—	M.fr.V,175.		

GUTTIFERAE.

A. L. de Jussieu; Gen. 243 (1789).—Genera 2, Species 3.

CALOPHYLLUM; Linné, gen. 154 (1737).											
C. inophyllum; Linné, sp. pl. 513 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,183	M.fr.IX,175	...	As., Afr., P.
C. tomentosum; Wight, Illustrat. Pl. Ind. Orient. I, 128 (1840)	—	—	—	—	—	Q.	—	—	M.fr.IX,174	...	As.
KAYEA; Wallich, pl. As. rar. III, 4 (1832).											
K. Larnachiana; F. v. M. in Vict. Nat. III, 126 (1887)	—	—	—	—	—	Q.	—				

POLYGALEAE.

A. L. de Jussieu, in Ann. du Mus. XIV, 386 (1809).—Genera 4, Species 32.

SALOMONIA; Loureiro, Fl. Cochinch. I, 14 (1790)											
S. oblongifolia; De Candolle, prodr. I, 334 (1824)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,138	M.fr.XI,4	...	As., P.
POLYGALA; Tournefort, inst. 174, t. 79 (1700), from Dioscorides and Plinius.											
P. Sibirica; Linné, sp. pl. 702 (1753)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,139	M.fr.XI,4	...	As.
P. leptalea; De Candolle, prodr. I, 325 (1824)	—	—	—	—	—	Q.	N.A.	B.fl.I,139	M.fr.XI,4	...	As.
P. eriocephala; F. v. M. in Benth. Fl. Austr. I, 139 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,139			
P. persicarifolia; De Candolle, prodr. I, 326 (1818)	—	—	—	—	—	Q.	N.A.	—	M.fr.XI,4	...	As., Afr.
P. Chinensis; Linné, sp. pl. 704 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,140	M.fr.XI,4	...	As.
P. stenoclada; Bentham, Fl. Austr. I, 141 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,141	M.fr.XI,4.		

COMESPERMA; Labillardière, Nov. Holl. pl. spec. II, 21 (1806).

C. sphaerocarpum; Steetz in Lehm. pl. Preiss. II, 304 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,143	M.fr.I,145; XI,3.
C. scoparium; Steetz in Lehm. pl. Preiss. II, 314 (1847)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,143	M.fr.I,145.
C. aphyllum; R. Brown in Benth. Fl. Austr. I, 143 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,143.	
C. spinosum; F. v. M., fragm. I, 144 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,144	M.fr.I,144.
C. volubile; Labillardière, Nov. Holl. pl. spec. II, 24, t. 163 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,144	M.fr.XI,3.
C. ciliatum; Steetz in Lehm. pl. Preiss. II, 304 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,144	M.fr.X,3.
C. integerimum; Endlicher in Hueg. enum. (1837)	W.A.	—	—	—	—	—	—	B.fl.I,145	M.fr.XI,3.
C. secundum; Banks in De Candolle, prodr. I, 334 (1824)	—	—	—	—	—	Q.	N.A.	B.fl.I,145.	
C. Drummondii; Steetz in Lehm. pl. Preiss. II, 301 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,145	M.fr.X,5; XI,2.
C. pracelsum; F. v. M., fragm. XI, 2 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,2.
C. retusum; Labillardière, Nov. Holl. pl. spec. II, 22, t. 160 (1806)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,145	M.fr.XI,2.
C. sylvestre; Lindley in Mitch. Trop. Austr. 342 (1848)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,146	M.fr.I,49; X,5; XI,2.
C. viscidulum; F. v. M., fragm. X, 4 (1876)	W.A.	S.A.	—	—	—	—	—	—	M.fr.XI,2.
C. accrosum; Steetz in Lehm. pl. Preiss. II, 299 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,146	M.fr.XI,3.
C. ericinum; De Candolle, prodr. I, 334 (1824)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,146	M.fr.I,48; XI,2.
C. confertum; Labillardière, Nov. Holl. pl. spec. II, 23, t. 161 (1806)	W.A.	—	—	—	—	—	—	B.fl.I,147	M.fr.XI,3.
C. flavum; De Candolle, prodr. I, 334 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,147	M.fr.XI,3.
C. calymega; Labillardière, Nov. Holl. pl. spec. II, 23, t. 162 (1806)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,147	M.fr.XI,3.
C. rhadinocarpum; F. v. M., fragm. XI, 1 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,1.
C. lanceolatum; R. Brown in Benth. Fl. Austr. I, 148 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,148	M.fr.XI,1.
C. defoliatum; F. v. M., pl. of Vict. I, 189 (1862)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,148	M.fr.XI,3.
C. nudiusculum; De Candolle, prodr. I, 334 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,148	M.fr.XI,4.
C. virgatum; Labillardière, Nov. Holl. pl. spec. II, 21, t. 159 (1806)	W.A.	—	—	—	—	—	—	B.fl.I,149	M.fr.XI,2.
C. polygaloides; F. v. M. in Trans. Phil. Soc. Vict. I, 7 (1854)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,149	M.fr.XI,4.

XANTHOPHYLLUM; Roxburgh, Pl. Corom. III, 81, t. 284 (1819). (Macintyria.)

X. Macintyrii; F. v. M, fragm. V, 57 (1865)	—	—	—	—	—	—	—	—	M.fr.V,8; XI,4.

TREMANDREAE.

R. Brown in Flind. voy. II, 554 App. (1814).—Genera 3, Species 17.

PLATYTHECA; Steetz in Lehmann, pl. Preiss. I, 220 (1845).

P. galioides; Steetz in Lehm. pl. Preiss. I, 220 (1845)	W.A.	—	—	—	—	—	—	B.fl.I,136	M.fr.XII.

TETRATHECA; Smith, Spec. Bot. New Holl. I, t. 2 (1793).

T. Harperi; F. v. M., fragm. V, 49 (1865)	W.A.	—	—	—	—	—	—	—	M.fr.V,189.
T. ciliata; Lindley in Mitch. three Exped. II, 206 (1838)	—	S.A.	T.	V.	—	—	—	B.fl.I,130	M.fr.XII,7.
T. efoliata; F. v. M., fragm. X, 3 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,3.
T. aphylla; F. v. M. in South. Science Record, vol. II, May (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII,5.
T. ericifolia; Smith, Exot. Bot. I, 37, t. 20 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,131	M.fr.XII,6.
T. juncea; Smith, specim. of Bot. of New Holl. 5, t. 2 (1793)	—	—	—	—	N.S.W.	—	—	B.fl.I,132	M.fr.XII,6.
T. affinis; Endlicher in Hueg. enum. 7 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,132	M.fr.XII,5.
T. nuda; Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII,5.
T. virgata; Steetz in Lehm. pl. Preiss, I, 212 (1845)	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII,5.
T. confertifolia; Steetz in Lehm. pl. Preiss. I, 214 (1845)	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII,6.
T. setigera; Endlicher in Hueg. enum. 8 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,133	M.fr.XII,8.
T. hirsuta; Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	W.A.	—	—	—	—	—	—	B.fl,I,134	M.fr.XII,7.
T. pilifera; Lindley, Bot. Regist. XXV, App. XXXVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,135	M.fr.XII,8.
T. filiformis; Bentham, Fl. Austr. I, 135 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,135	M.fr.XII,8

TREMANDRA ; R. Brown in App. Flind. voy. II, 554 (1814).

T. stelligera ; R. Brown in De Candolle, prodr. I, 344 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,136	M.fr.XII,8.					
T. diffusa ; R. Brown in De Candolle, prodr. I, 344 (1824)	W.A.	—	—	—	—	—	—	B.fl.I,137	M.fr.XII,9.					

MELIACEAE.

VENTENAT ; Tabl. III, 159 (1799).—Genera 11, Species 36.

TURRAEA ; Linné mantiss. II, 150 (1771).

T. pubescens ; Hellenius in K. Sv. Vet. Acad. Handl. 26, t. 10 (1788)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,379	M.fr.V,144	...	As.			

MELIA ; Linné, gen. pl. 127 (1737).

M. Azedarach ; Linné, sp. pl. 384 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,380	M.fr.V,144 ; IX,134 ; XI,135	As., Afr.				

DYSOXYLUM ; Blume, Bijdrag. 172 (1825). (Dysoxylon, Epicharis, Hartigshea.)

D. oppositifolium ; F. v. M., fragm. V, 144 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,177 ; IX,61.					
D. latifolium ; Bentham, Fl. Austr. I, 381 (1863)	—	—	—	—	—	Q.	—	B.fl.I,381	M.fr.V,144 ; IX,61.					
D. Klanderi ; F. v. M., fragm. V, 176 (1866)...	—	—	—	—	—	Q.	—	—	M.fr.IX,61,97,134.					
D. Fraserianum ; Bentham, Fl. Austr. I, 381 (1863)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,381	M.fr.IX,61.					
D. arborescens ; Miquel in Annal. Mus. Bot. Lugd. IV, 24 (1868)	—	—	—	—	—	Q.	—	—	M.fr.IX,134 ...	As.				
D. Muelleri ; Bentham, Fl. Austr. I, 381 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,381	M.fr.V,145.					
D. Schiffneri ; F. v. M. in Melb. Chemist, 53 (1881)	—	—	—	—	—	Q.	—	—	M.fr.XI,138.					
D. Becklerianum ; C. de Candolle in Monogr. phan. I, 509 (1878)...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,138 ; XI,82.					
D. Lessertianum ; Bentham, Fl. Austr. I, 382 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,382	M.fr.V,145 ; IX,61.					
D. Schultzii ; C. de Candolle in Monogr. phanerog. I, 509 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.XI,138.					
D. rufum ; Bentham, Fl. Austr. I, 382 (1863)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,382	M.fr.V,145.					
D. Nernstii ; F. v. M., fragm. V, 176 (1866)	—	—	—	—	—	Q.	—	—	M.fr.V,176.					
D. Patersonianum ; Bentham & Hooker, gen. pl. I, 332 (1862) implied	—	—	—	—	N.S.W.	—	—	—						

AMOORA ; Roxburgh, pl. Corom. III, 54, t. 258 (1819).

A. nitidula ; Bentham, Fl. Austr. I, 383 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,383	M.fr.V,145 ...	P.				

SYNOUM ; Adr. de Jussieu in Mém. du Mus. XIX, 226, t. 15 (1830).

A. glandulosum ; A. de Jussieu in Mém. du Mus. XIX, 227, t. 15 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,384	M.fr.V,145 ; IX,134.					
A. Muelleri ; C. de Candolle in Monogr. phanerog. I, 593 (1878)	—	—	—	—	—	Q.	—	—	M.fr.V,145 ; XI,138.					

OWENIA ; F. v. M. in Hook. Kew Misc. IX, 304 (1857).

O. acidula ; F. v. M. in Hook. Kew Misc. IX, 303 (1857)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,385	M.fr.III,14 ; IX,134.					
O. vernicosa ; F. v. M., fragm. III, 15 (1862)...	—	—	—	—	—	—	N.A.	B.fl.I,385	M.fr.IX,134.					
O. venosa ; F. v. M. in Hook. Kew Misc. IX, 304 (1857)	—	—	—	—	—	Q.	—	B.fl.I,386	M.fr.III,14 ; IX,134 ; XI,82.					
O. cepiodora ; F. v. M., fragm. XI, 81 (1880)...	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,87 ...					
O. reticulata ; F. v. M. in Hook. Kew Misc. IX, 305 (1857)	W.A.	—	—	—	—	—	N.A.	B.fl.I,386	M.fr.III,13 ; IX,134.					

AGLAIA ; Loureiro, Fl. Cochinch. I, 173 (1790), non Allemand (1770). (Milnea, Nemedra.)

A. elaeagnoides ; Bentham, Fl. Austr. I, 383 (1863)...	—	—	—	—	—	Q.	N.A.	B.fl.I,383	M.fr.V,145 ; IX,134.					

HEARNIA ; F. v. M., fragm. V, 55 (1865). (Aglaiopsis).

H. sapindina ; F. v. M., fragm. V, 55 (1865)...	—	—	—	—	—	Q.	—	B.fl.I,387	M.fr.V,55.					

CARAPA ; Aublet, Hist. pl. Guian. II, Suppl. 32, t. 387 (1775). (Xylocarpus.)

C. Moluccensis ; Lamarck, Encycl. meth. Bot. I, 621 (1783)	—	—	—	—	—	Q.	N.A.	B.fl.I,387	M.f.V,145 ; IX,134.	As., Afr.				

CEDRELA ; P. Browne, Nat. hist. of Jamaica 158 (1756).

C. Australis ; F. v. M., fragm. I, 4 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,387	M.fr.IX,133 ; XI,82.					

FLINDERSIA; R. Brown in Flind. voy. II, 595 (1814). (Oxleya, Strzeleckia.)

F. Australis; R. Brown in Flind. voy. II. 595, t. 1 (1814)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,388	M.fr.I,65; II,174; III,26; IX,133.
F. Bourjotiana; F. v. M., fragm. IX, 133 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,133.
F. Schottiana; F. v. M., fragm. III, 25 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,388	M.fr.V,143; VI,143; IX,133.
F. Oxleyana; F. v. M., fragm. I, 65 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,389	M.fr.III,25; IX,133.
F. Pimenteliana; F. v. M., fragm. IX, 132 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,132.
F. Bennettiana; F. v. M. in Benth. Fl. Austr. I, 389 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,389	M.fr.IX,131; XI,135.
F. Ifflaiana; F. v. M., fragm. X, 94 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,94.
F. Brayleyana; F. v. M., fragm. V, 143 (1866)	—	—	—	—	—	Q.	—	—	M.fr.VI,252.
F. Strzeleckiana; F. v. M., fragm. I, 65 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,389	M.fr.IX,133.

OCHNACEAE.

De Candolle in Ann. du Mus. XVII, 398 (1811), Ochnae subgenus.—Genus 1, Species 1.

BRACKENRIDGEA; Asa Gray, Bot. Wilk. Expl. Exped. 361, t. 42 (1854).

B. Australiana; F. v. M., fragm. V, 29 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,213.

RUTACEAE.

A. L. de Jussieu, Gen. 296 (1789).—Genera 26, Species 190.

ZIERIA; Smith in Transact. Linn. Soc. IV, 216 (1798).

Z. laevigata; Smith in Rees's Cyclopaedia XXXIX (1819)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,304	M.fr.I,101; IX,116.
Z. pilosa; Rudge in Transact. Linn. Soc. X, 293, t. 17 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.I,305	M.fr.I,101; IX,116.
Z. obcordata; Cunningham in Field, N.S.Wales, 330 (1625)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,305	M.fr.100; IX,116.
Z. cytisoides; Smith in Rees's Cyclopaedia XXXIX (1819)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,306	M.fr.IX,116.
Z. Smithii; Andrews, Bot. Rep. t. 606 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,306	M.fr.I,100; IX,116.
Z. granulata; C. Moore in Benth. Fl. Austr. I, 307 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,307	M.fr.I,101; IX,116.
Z. veronicea; F. v. M. in Trans. phil. Soc. Vict. I, 11 (1854)	—	S.A.	T.	V.	—	—	—	B.fl.I,305	M.fr.IX,116.

BORONIA; Smith, Tracts relat. nat. hist. 285 (1798). (Cyanothamnus.)

B. grandisepala; F. v. M., fragm. I, 66 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,311	M.fr.II,179.
B. artemisifolia; F. v. M., fragm. I, 66 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,311	M.fr.IX,111.
B. affinis; R. Brown in Benth. Fl. Austr. I, 311 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,311	
B. filicifolia; Cunningham in Benth. Fl. Austr. I, 311 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,311	
B. alata; Smith in Transact. Linn. Soc. VIII, 283 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,312	M.fr.IX,111.
B. algida; F. v. M. in Transact. phil. Soc. Vict. I, 100 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.I,312	M.fr.IX,111.
B. Ewardsii; Bentham, Fl. Austr. I, 312 (1863)	—	S.A.	—	—	—	—	—	B.fl.I,312	
B. ternata; Endlicher, nov. stirp. dec. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.I,312	M.fr.IX,111.
B. ericifolia; Bentham, Fl. Austr. I, 313 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,313	M.fr.IX,111.
B. inconspicua; Bentham, Fl. Austr. I, 313 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,313	
B. eriantha; Lindley in Mitch. Trop. Austr. 298 (1848)	—	—	—	—	—	Q.	—	B.fl.I,313	
B. alulata; Solander in Benth. Fl. Austr. I, 313 (1863)	—	—	—	—	—	Q.	—	B.fl.I,313	
B. ledifolia; J. Gay, monogr. des Lasiopetalées 20 (1821)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,314	M.fr.I,67; IX,111.
B. lanceolata; F. v. M., fragm. I, 66 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,314	M.fr.VII,38; IX,112; XI,97.
B. Bowmanii; F. v. M., fragm. IV, 135 (1864)	—	—	—	—	—	Q.	—	—	M.fr.VII,38.
B. Fraseri; Hooker, Bot. Mag. t. 4052 (1844)	—	—	—	—	N.S.W.	—	—	B.fl.I,315	M.fr.IX,111.
B. mollis; Cunningham in Lindl. Bot. Reg. 47 (1841)	—	—	—	—	N.S.W.	—	—	B.fl.I,315	M.fr.IX,112.

B. megastigma; Nees in Lehm. Pl. Preiss. II, 227 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,315	M.fr.II,97 ; IV,172; IX,112.
B. heterophylla ; F. v. M., fragm. II, 98 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,315	M.fr.IX,112.
B. elatior ; Bartling in Lehm. pl. Preiss. I, 170 (1844)	W.A	—	—	—	—	—	—	B.fl.I,316	M.fr.II,98 ; IX,122 ; XI,97.
B. tetrandra ; Labillardière, Nov. Holl. pl. spec. I, 98, t. 125 (1804)	W.A.	—	—	—	—	—	—	B.fl.I,316	
B. crassifolia ; Bartling in Lehm. pl. Preiss. I, 169 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,316	M.fr.IX.112.
B. albiflora ; R. Brown in Benth. Fl. Austr. I, 317 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,317	M.fr.IX,114.
B. lanuginosa; Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,317	M.fr.IX,114.
B. pulchella ; Turczaninow in Bull. Soc. Mosc. XXV, 162 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,318	M.fr.IX,114.
B. gracilipes ; F. v. M., fragm. II, 99 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,318	M.fr.IX,114.
B. microphylla ; Sieber in Spreng. syst. cur. post. 148 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,318	
B. floribunda ; Sieber in Spreng. syst. veg. cur. post. 148 (1827)	—	—	—	—	N.S.W.	—	—	—	
B. pinnata ; Smith, Tracts relat. nat. hist. 290, t. 4 (1798)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,318	M fr.IX,114.
B. pilosa ; Labillardière, Nov. Holl. pl. spec. I, 97, t. 124 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,319	
B. spinescens ; Bentham, Fl. Austr. I, 319 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,319	
B. baeckeacea ; F. v. M., fragm. IV, 28 (1863)	W.A.	—	—	—	—	—	—	—	M.fr.IX,112.
B. coerulescens ; F. v. M. in Trans. phil. Soc. Vict. II (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,320	M.fr.VII,38 ; IX,112.
B. defoliata ; F. v. M., fragm. IX, 113 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX.113.
B. Busselliana ; F. v. M., fragm. IX, 113 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,113.
B. subcoerulea ; F. v. M., fragm. II, 100 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,320	M.fr.IX,112&198.
B polygalifolia ; Smith, Tracts relat. nat. hist. 297, t. 7 (1798)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,320	M.fr.IX,114.
B. falcifolia ; Cunningham in Lindl. Bot. Reg. 47 (1841)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,322	M.fr.IX,114.
B. penicillata ; Bentham, Fl. Austr. I, 322 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,322	
B. crassipes ; Bartling in Lehm. pl. Preiss. I, 168 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,322	M.fr.IX,116.
B. subsessilis ; Bentham, Fl. Austr. I, 322 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,322	
B. capitata ; Bentham, Fl. Austr. I, 323 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.IX,116.
B. nematophylla ; F. v. M., fragm. II, 100 (1861)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.II,100.
B. crenulata ; Smith in Transact. Linn. Soc. VIII, 284 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,323	M.fr.XI,97,115&198.
B. serrulata ; Smith, Tracts relat. nat. hist. 292 (1798)	—	—	—	—	N.S.W.	—	—	B.fl.I,323	M.fr.IX,115.
B. rhomboidea ; Hooker icon. pl. t. 722 (1848)	—	—	T.	—	N.S.W.	—	—	B.fl.I,324	
B. parviflora ; Smith, Tracts relat. nat. hist. 295. t. 5 (1798)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,324	M.fr.IX,115.
B. viminea ; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,324	
B. filifolia ; F. v. M., fragm. I, 3 (1858)	—	S.A.	—	V.	—	—	—	B.fl.I,325	M.fr.IX,114.
B. clavellifolia ; F. v. M. in Trans. phil. Soc. Vict. I, 12 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,325	M.fr.I,99 ; VII,38 ; IX,114&198.
B. scabra ; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,326	
B. thymifolia ; Turczaninow in Bull. Soc. Mosc. XXV, part II, 165 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.I,326	M.fr.I,99 ; II,99 ; IX,116.
B. ovata ; Lindley, Bot. Regist. 47 (1841)	W.A.	—	—	—	—	—	—	B.fl.I,326	M.fr.IX,116.
B. Barkeriana ; F. v. M., fragm. XI, 96 (1880)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,96.
B. denticulata ; Smith in Transact. Linn. Soc. VIII, 284 (1807)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115&198.
B. Machardiana ; F. v. M., fragm. IX, 115 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,115.
B. spathulata ; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115.
B. juncea ; Bartling in Lehm. pl. Preiss. I, 166 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,327	M.fr.IX,115.
B. cymosa ; Endlicher in Hueg. enum. 16 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,328	M.fr.II,101 ; IX,116.

ERIOSTEMON ; Smith in Transact. Linn. Soc. IX, 221 (1798). (Crowea, Phebalium, Asterolasia, Microcybe, Pleurandropsis, Geleznowia, Urocarpus, Sandfordia, Actinostigma.)

E. virgatus ; Cunningham in Hooker's Journ. of Bot. II, 417 (1840)	—	—	T.	—	—	—	—	B.fl.I,332	M.fr.IX,111.
E. nodiflorus ; Lindley, Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,336	M.fr.IX,110 ; X,53.
E. spicatus ; A. Richard, sert. Astrolab. 76, t. 27 (1833)	W.A.	—	—	—	—	—	—	B.fl.I,336	M.fr.IX,110.
E. pungens ; Lindley in Mitch. Three Exped. II, 156 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,338	M.fr.IX,110.

E. montanus; F. v. M., pl. of Vict. I, 129 (1860)	—	—	T.	—	—	—	—	B.fl.I,338	M.fr.IX,109.
E. phylicifolius; F. v. M., fragm. I, 105 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.I,105.
E. umbellatus; Turczaninow in Bull. Soc. Mosc. XXII, part III, 15 (1849) ...	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.I,104; IX,109.
E. Ralstoni; F. v. M., fragm. II, 101, t. 14 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.I,339	M.fr.III,165; IX,109.
E. Hillebrandi; F. v. M. in Transact. phil. Soc. Vict. I, 10 (1854)	—	S.A.	T.	V.	—	—	—	B.fl.I,340	M.fr.I,4; IX,109.
E. Coxii; F. v. M. in Melb. Chem., Dec. (1884)	—	—	—	—	N.S.W.	—	—	—	
E. lamprophyllus; F. v. M. in Journ. pharm. Soc. Vict. II, 43 (1858)	—	—	—	V.	N.S.W.	—	—	B.fl.I,340	M.fr.IX,109.
E. elatior; F. v. M., fragm. I, 181 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,340	M.fr.IX,109.
E. amplifolius; F. v. M. in Melb. Chem., Dec. (1884)	—	—	—	V.	N.S.W.	—	—	—	
E. ambiens; F. v. M., fragm. VI, 166 (1868)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,110.
E. Oldfieldii; F. v. M., fragm. I, 3 (1858)	—	—	T.	—	—	—	—	B.fl.I,340	M.fr.I,181; IX,109.
E. rotundifolius; Cunningham in Hueg. enum. 15 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,341	
E. microphyllus; F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854)	—	S.A.	—	—	—	—	—	B.fl.I,341	M.fr.IX.109.
E. phylicoides; F. v. M., fragm. I, 107 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,341	M.fr.IX.107.
E. ozothamnoides; F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,342	M.fr.IX,107.
E. Mortoni; F. v. M., fragm. IX, 108 (1875)	—	—	—	—	N.S.W.	—	—	B.fl.I,342	M.fr.IX,108.
E. sediflorus; F. v. M., fragm. I, 102 (1859)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,342	M.fr.I.102.
E. Nottii; F. v. M., fragm. VI, 22 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VI,22.
E. lepidotus; Sprengel, syst. II, 322 (1825)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,343	M.fr.I,104; IX,107.
E. stenophyllus; F. v. M., from Benth. Fl. Austr. I, 343 (1863)	—	S.A.	—	V.	N.S.W.	—	—	—	
E. alpinus; F. v. M., fragm. I, 103 (1859)	—	—	—	V.	N.S.W.	—	—	—	M.fr.I,244.
E. tuberculosus; F. v. M., pl. of Vict. I, 130 (1862)	W.A	—	—	—	—	—	—	B.fl.I,343	M.fr.IX,108.
E. Benthami; F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	B.fl.I,343	M.fr.IX,108.
E. Maxwelli; F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,103.
E. filifolius; F. v. M., fragm. IX, 108 (1875)	W.A.	—	—	—	—	—	—	B.fl.I,344	M.fr.IX,108.
E. squameus; Labillardière, Nov. Holl. pl. spec. I, 3, t. 141 (1804)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I.344	M.fr.I,104; IX,107.
E. anceps; Sprengel, syst. II, 322 (1825)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.I,103; IX,108.
E. ovatifolius; F. v. M., fragm. I, 103 (1859)...	—	—	—	V.	N.S.W.	—	—	B.fl.I,345	M.fr.I,103.
E. bilobus; F. v. M., fragm. I, 102 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,345	M.fr.IX,108.
E. Beckleri; F. v. M., fragm. IX, 109 (1875)...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,108.
E. capitatus; F. v. M., fragm. I, 106 (1859)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,346	M.fr.IX,107.
E. Geleznowii; F. v. M., fragm. I, 107 (1859)	W.A.	—	—	—	—	—	—	B.fl.I 347	M.fr.IX,107.
E. correifolius; F. v. M., fragm. I, 105 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,350	M.fr.IX,107.
E. Cunninghami; F. v. M., fragm. IX, 107 (1875)	—	—	—	—	N.S.W.	—	—	B.fl.I,351	M.fr.IX,107.
E. mollis; F. v. M., syst. cens. of Austr. pl. 11 (1882), from Bentham (1863) ...	—	—	—	—	N.S.W.	—	—	B.fl.I,349	
E. pleurandroides; F. v. M., fragm. I, 106 (1859)	—	—	—	V.	—	—	—	B.fl.I,349	M.fr.IX,107.
E. trymalioides; F. v. M., fragm. I, 106 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.I,351	M.fr.I,106.
E. Hookeri; F. v. M., fragm. I, 104 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,352	M.fr.VII,22; IX,107.
E. pallidus; F. v. M., syst. cens. of Austr. pl. 11 (1882), from Bentham (1863)	W.A.	—	—	—	—	—	—	B.fl.I,352	
E. Drummondii; F. v. M., fragm. I, 105 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,352	M.fr.VII,22.
E. grandiflorus; F. v. M., fragm. I, 105 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,353	M.fr.VI,21.
E. Crowei; F. v. M., pl. of Vict. I, 119 (1860)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,329	M.fr.IX,106
E. Turczaninowii; F. v. M., pl. of Vict. I, 120 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,330	M.fr.IX,106.
E. lanceolatus; K. F. Gaertner, de fruct. III, 154, t. 210 (1807)	—	—	—	V.	N.S.W.	—	—	B.fl.I,331	M.fr.IX,110.
E. Banksii; Cunningham in Hueg. enum. 15 (1837)	—	—	—	—	—	Q.	—	B.fl.I,332	
E. trachyphyllus; F. v. M. in Transact. phil. Soc. Vict. I, 99 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.I,332	M.fr.IX,110.
E. myoporoides; De Candolle, prodr. I, 720 (1824)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,333	M.fr.XI.110.
E. hispidulus; Sieber in Spreng. syst. cur. post. 164 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,333	M.fr.XI,110.

E. buxifolius; Smith in Rees' Cyclop. XIII, 2 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.I,333	M.fr.IX,111.
E. obovalis; Cunningham in Field's N.S. Wales, 331 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,334	M.fr.IX,110.
E. scaber; Paxton, Mag. of Bot. XIII, 127 (1846)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,334	M.fr.XI.111.
E. Brucei; F. v. M., fragm. VII, 38 (1869)	W.A.	—	—	—	—	—	—	—	M.fr.IX,110.
E. linearis; Cunningham in Hueg. enum. 16 (1837)	—	S.A.	—	—	N.S.W.	—	—	B.fl.I,334	
E. difformis; Cunningham in Hueg. enum. 15 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,335	M.fr.IX,110.
E. parvifolius; R. Brown in Benth. Fl. Austr. 1, 335 (1863)	—	—	—	—	—	Q.	—	B.fl.I,335	
E. ericifolius; Cunningham in Benth. Fl. Austr. I, 335 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,335	
PHILOTHECA; Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815) (Drummondita.)									
P. Australis; Rudge in Transact. Linn. Soc. XI, 298, t. 21 (1815)	—	—	--	—	N.S.W.	Q.	—	B.fl.I,348	M.fr.VII,20,141.
P. ericoides; F. v. M., fragm. I, 107 (1859) implied	W.A.	—	—	—	—	—	—	B.fl.I,349	M.fr.VII,21.
P. Hasselli; F. v. M. in Wing's S. Sc. Rec. III, 3 (1883)	W.A.	—	—	—	—	—	—	—	
P. calida; F. v. M., fragm. VII, 21 (1869)	—	—	—	—	—	Q.	—	—	M.fr.VII,38.
CORREA; Smith in Transact. Linn. Soc. IV, 219 (1798). (Didymeria, Mazeutoxeron, Automachia).									
C. aemula; F. v. M., fragm. I, 3 (1859)	—	S.A.	—	V.	—	—	—	B.fl.I,353	M.fr.IX,117.
C. alba; Andrews, Bot. Reposit. t. 18 (1797)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,354	M.fr.IX.117.
C. speciosa; Andrews, Bot. Reposit. t. 653 (1811)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,354	M.fr.IX,117.
C. Lawrenciana; Hooker, Journ. of Bot. I, 254 (1834)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,355	M.fr.VIII,142; IX,117.
C. decumbens; F. v. M. in Transact. phil. Soc. Vict. I, 30 (1854)	—	S.A.	—	—	—	—	—	B.fl.I,356	
C. Baeuerlenii; F. v. M. in proc. L. S. of N.S.W. IX, 960 (1884)	—	—	—	—	N.S.W.	—	—	—	
NEMATOLEPIS; Turczaninow in Bull. Soc Mosc. XXV, part III, 158 (1852). (Symphyopetalum.)									
N. phlebalioides; Turczaninow in Bull. Soc. Mosc. part III, 158 (1852)... ...	W.A.	—	—	—	—	—	—	B.fl.I,351	M.fr.II,102; IX,106.
N. Euphemiae; F. v. M., fragm. III, 149, t. 25 (1863)	W.A.	—	—	--	—	—	—	B.fl.I,345	M.fr.IX,166.
CHORILAENA; Endlicher in Hueg. enum. 17 (1837).									
C. quercifolia; Endlicher in Hueg. enum. 17 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,357	M.fr.IX,106.
C. hirsuta; Bentham, Fl. Austr. I, 357 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,357	M.fr.IX,106.
DIPLOLAENA; R. Brown in Append. Flind. voy. II, 546 (1814).									
D. Dampieri; Desfontaines in Mém. du Mus. 452, t. 20 (1817)	W.A.	—	—	—	—	—	—	B.fl.I,358	M.fr.IX.106.
BOSISTOA; F. v. M. in Benth. Fl. Austr. I, 359 (1863).									
B. sapindiformis; F. v. M. in Benth. Fl. Austr. I, 359 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,359	M.fr.IX,103; III,41,168; IV,175.
B. enodiformis; F. v. M., fragm. IX, 174 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,174.
ACRADENIA; Kippist in Transact. Linn. Soc. XXI, 207 (1855).									
A. Frankliniae; Kippist in Transact. Linn. Soc. XXI, 207 (1855)	—	—	T.	—	—	—	—	B.fl.I,323	M.fr.VI,167; IX,103.
EUODIA; R. et G. Forster, char. gen. 13, t. 7 (1776). (Melicope, Medicosma.)									
E. Cunninghami; F. v. M., fragm. III, 2 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,362	M.fr.I,27; II,178; IX,102.
E. Fareana; F. v. M., fragm. IX, 101 (1875)...	—	—	—	—	—	Q.	—	--	M.fr.IX,101.
E. erythrococca; F. v. M., fragm. I, 26 (1858)	-	—	—	—	N.S.W.	Q.	—	B.fl.I,360	M.fr.II,103.178; IX,102.
E. contermina; Moore & F. v. M., fragm. VII, 144 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII.144.
E. octandra; F. v. M., fragm. II, 102 (1861)	—	—	—	—	N.S.W.	—	—	B.fl.I,360	M.fr.IV,117; IX.102.
E. micrococca; F. v. M., fragm. I, 144 (1859)	-	—	—	—	N.S.W.	Q.	—	B.fl.I,361	M.fr.II,103; III,3,180; IX,102.
E. xanthoxyloides; F. v. M., fragm. IV, 155 (1864)...	—	—	—	—	—	Q.	—	—	M.fr.IV,155.
E. Bonwickii; F. v. M., fragm. V, 56 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,179.
E. alata; F. v. M., fragm. VII, 142 (1871)	—	—	—	—	—	Q.	—	—	M.fr.VII,142
E. accedens; Blume, Bijdr. 246 (1825)...	—	—	—	—	N.S.W.	Q.	—	--	M.fr.V,4,56,179; VII,22; IX,102. As.
E. litoralis; Endlicher, prodr. Fl. Norfolk. 86 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,102.

E. polybotrya; Moore & F. v. M., fragm. VII, 143 (1871) ...	...	...	...	—	—	—	—	N.S.W.	—	—		—	M.fr.IX,193.				
E. vitiflora; F. v. M., fragm. VII, 144 (1871)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VII,144.				
E. haplophylla; F. v. M., fragm. V, 179 (1866)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.V,179.				
BROMBYA; F. v. M., fragm. V, 4 (1865).																	
B. platynema; F. v. M., fragm. V, 4 (1865) ...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.V,56; IX,102.				
PAGETIA; F. v. M., fragm. V, 178 (1866).																	
P. medicinalis; F. v. M., fragm. V, 178 (1866)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.V,215; VI,167; IX,103.				
BOUCHARDATIA; Baillon in Adans. VII, 350 t. 10 (1867).																	
B. neurococca; Baillon in Adans. IX, 110 (1869)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,360	M.fr.I,28; II,103,178; IX,103.				
XANTHOXYLUM; Catesby Nat. Hist. of Carolina I, 26 (1731). (Zanthoxylum, Blackburnia.)																	
X. brachyacanthum; F. v. M., pl. of Vict. I, 108 (1862)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,363	M.fr.VII,141; IX,104.				
X. Blackburnia; Bentham, Fl. Austr. I, 363 (1863) ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I,363	M.fr.VII,141.				
X. parviflorum; Bentham, Fl. Austr. I, 363 (1863) ...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,363	M.fr; IX,104.				
X. torvum; F. v. M., fragm. VII, 140 (1871)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VII,140; IX,104.				
X. venenificum; Bailey, catal. of Ql. woods, 8 (1886)	...	...	...	...	—	—	—	—	—	Q.	—	—					
GEIJERA; Schott, Rutac. 7, t. 4 (1834). (Coatesia.)																	
G. Muelleri; Bentham, Fl. Austr. I, 364 (1863)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,364	M.fr.I,26; IX,104.				
G. salicifolia; Schott, Rutac. 7, t. 4 (1834)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,364	M.fr.VII,22; IX,105; XI,134.				
G. parviflora; Lindley in Mitch. Trop. Austr. 102 (1848)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,364	M.fr.VII,22; IX,105; X,53; XI,139.				
PLEIOCOCCA; F. v. M., fragm. IX, 117 (1875).																	
P. Wilcoxiana; F. v. M., fragm. IX, 117 (1875)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,117.				
ACRONYCHIA; R. & G. Forster, char. gen. 53, t. 27 (1776). (Cyminosma.)																	
A. Baueri; Schott, Rutac. 5, t. 3 (1834)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,366	M.fr.I,26; II,178; V,179; IX,103.				
A. laevis; R. & G. Forster, char. gen. 53, t. 27 (1776)	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,366	M.fr.I,27; V,179; IX,104.	P.			
A. imperforata; F. v. M., fragm. I, 26 (1858)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,367	M.fr.IX,104.				
A. melicopoides; F. v. M., fragm. V, 3 (1865)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	--	M.fr.IV,117; V,179; VII,145.				
A. acidula; F. v. M., fragm. IV, 154 (1864) ...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IV,154.				
A. vestita; F. v. M., fragm. IV, 155 (1864) ...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,104.				
A. Endlicheri; Schott, fragm. bot. III, t. 2 ...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,103.				
A. tetrandra; F. v. M., fragm. IX, 104 (1875)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.V,179.				
HALFORDIA; F. v. M., fragm. V, 43 (1865).																	
H. drupifera; F. v. M., fragm. V, 43 (1865) ...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,5,189; VII,142; IX,103.				
H. scleroxyla; F. v. M., fragm. VII, 142 (1871)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,103.				
GLYCOSMIS; Correa in Ann. du Mus. VI, 384 (1805).																	
G. pentaphylla; Correa in Annal. du Mus. VI, 384 (1805) ...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,367	M.fr.IX,105.	...	...	As., P.	
MICROMELUM; Blume, Bijdrag. 137 (1825).																	
M. pubescens; Blume, Bijdrag. 137 (1825)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,368	M fr.I,25; IX,105.	...	As., P.		
MURRAYA; Koenig in Linné, mant. alt. 563 (1771). (Chaslas, 1767.)																	
M. exotica; Koenig in Linné, mant. alt. 563 (1771) ...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,369	M.fr.IX,106.	...	...	As., P.	
M. crenulata; Oliver in Journ. Linn. Soc. V, suppl. II, 29 (1861)...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,369	...	...	...	...	As.	
CLAUSENA; Burmann, Fl. Indica index & t. 29 (1768).																	
C. brevistyla; Oliver in Journ. Linn. Soc. V, suppl. II, 31 (1861)	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,369	M.fr.IX,106.					
ATALANTIA; Correa in Ann. du Mus. VI, 383 (1805).																	
A. glauca; J. Hooker in B. & H. Gen. pl. I, 305 (1862)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,370	M.fr.XI,105.				
A. recurva; Bentham, Fl. Austr. I, 370 (1863)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,370	M.fr.VII,142.				

CITRUS ; Linné, gen. pl. 230 (1737), from Plinius.

C. Planchonii ; F. v. M., Report Intercol. Exhib. 23 (1867)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,371	M.fr.IX,105.				
C. Australasica ; F. v. M., fragm. I, 26 (1858)	...	...	..	..	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,371	M.fr.II,178 ; III,167 ; IX,105.		

PENTACERAS ; J. Hooker in B. & H. Gen. pl. I, 298 (1862).

P. Australis ; J. Hooker in B. & H. Gen. pl. I, 298 (1862)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,365	M.fr.I,25 ; IX,104.			

SIMARUBEAE.

De Candolle in Ann. du Mus. XVII, 323 (1811).—Genera 6, Species 7.

AILANTUS ; Desfontaines, in Act. Acad. Paris 265 (1786).

A. imberbiflora ; F. v. M., fragm. III, 42 (1862)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,373	M.fr.VI,166.		

BRUCEA ; J. S. Miller, icon. plantar. t. 25 (1780).

B. Sumatrana ; Roxburgh, Fl. Indic. I, 469 (1820)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,373	M.fr.VI,166	...	As.

HYPTIANDRA ; J. Hooker in B. & H. Gen. pl. 294 (1862).

H. Bidwilli ; J. Hooker in B. & H. Gen. pl. 294 (1862)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,374	M.fr.VI,165.			

CADELLIA ; F. v. M., fragm. II, 25 (1860).

C. pentastylis ; F. v. M., fragm. II, 26, t. 12 (1860)	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.I,374	M.fr.II,171.			
C. monostylis ; Bentham, Fl. Austr. I, 375 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,375	M.fr.VI,166 ; VIII,34.		

SURIANA ; Plumier, nov. plant. Amer. gen. 37, t. 40 (1703).

S. maritima ; Linné, sp. pl. 284 (1753)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,375	...	...	...	As., Afr., Am., P.

HARRISONIA ; R. Brown in Mém. du Mus. XII, 517 (1825). (Ebelingia.)

H. Brownii ; A. L. de Jussieu in Mém. du Mus. Par. XII, 530, t. 28 (1825)	...	—	—	—	—	—	—	Q.	N.A.	B.fl.I,376	...	..	...	As., P.

ZYGOPHYLLEAE.

R. Brown in App. Flind. voy. II, 545 (1814)—Genera 3, Species 22.

NITRARIA ; Linné, syst. nat. X, 1044 (1759).

N. Schoberi ; Linné, sp. pl. ed. sec. 639 (1762)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,291	M.fr.IX,75 ; XI,30	...	As., Afr.

ZYGOPHYLLUM ; Linné syst. nat. 8 (1735), Linné, gen. 126 (1737). (Sarcozygium, Roepera.)

Z. apiculatum ; F. v. M. in Linnaea XXV, 373 (1852)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,292	M.fr.XI,29.			
Z. glaucescens ; F. v. M., pl. of Vict. I, 228 (1862)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,293	M.fr.XI,29.			
Z. crenatum ; F. v. M. in Linnaea XXV, 374 (1852)	...	..	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	—				
Z. iodocarpum ; F. v. M. in Linnaea XXV, 372 (1852)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,293	M.fr.XI,29.			
Z. prismatothecum ; F. v. M. in Linnaea XXV, 375 (1852)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,293				
Z. ammophilum ; F. v. M. in Linnaea XXV, 376 (1852)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	—	M.fr.XI.28.			
Z. Billardieri ; De Candolle, prodr. I, 705 (1824)	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,293	M.fr.XI.29.			
Z. fruticulosum ; De Candolle, prodr. I, 705 (1824)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,594	M.fr.IX,75 ; XI,29.			
Z. Howittii ; F. v. M., fragm. III, 150 (1863)	...	...	...	...	—	S.A.	—	—	—	—	—	—	M.fr.XI,29	...	..	P.

TRIBULUS ; L'Obel, Icon. II, 84 (1581), from Theophrastos, Dioscorides & Plinius). (Tribulopsis.)

T. terrestris ; L'Obel, stirp. icones II, 84 (1581)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,288	M.fr.IX,75 ; XI,30 ; XII,4.	E., As., Afr, Am.,
T. ranunculiflorus ; F. v. M., fragm. I, 48 (1858)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,288	M.fr.I,48.		[P.
T. hystrix ; R. Brown in App. Sturt. Exped. 6 (1849)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	N.A.	B.fl.I,289	M.fr.XI,30.		
T. macrocarpus ; F. v. M. in Benth. Fl. Austr. I, 289 (1863)	...	...	..	W.A.	S.A.	—	—	—	—	—	N.A.	B.fl.I,289				
T. Forrestii ; F. v. M. in Wing's S. Sc. rec. Nov. (1885)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	—				

T. platypterus ; Bentham, Fl. Austr. I, 289 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,289					
T. hirsutus ; Bentham, Fl. Austr. I, 289 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,289					
T. minutus ; Leichhardt in Benth. Fl. Austr. I, 291 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,241					
T. Brownii ; F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,290	M.fr.I,48.				
T. bicolor ; F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	—	N.A.	B.fl.I,290	M.fr.I,47.				
T. Solandri ; F. v. M., pl. of Vict. I, 99 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,290	M.fr.I,47 ; XI,30.				
T. astrocarpus ; F. v. M., fragm. XII, 4 (1882)	W.A.	S.A.	—	—	—	—	—	—					

LINEAE.

De Candolle, Théor. élém. bot. 214 (1813).—Genera 3, Species 4.

HUGONIA ; Linné, gen. pl. 134 (1737).

H. Jenkinsii ; F. v. M., fragm. V, 7 (1865)	—	—	—	—	—	Q.	—	—	M.fr.IX,190.				

ERYTHROXYLUM ; P. Brown, Hist. pl. of Jamaica 278 (1756), from Theophrastos, Dioscorides & Plinius. (Erythroxylon.)

E. Australe ; F. v. M. in Transact. Vict. Inst. III, 22 (1855)	—	—	—	—	—	Q.	—	B.fl.I,284					
E. ellipticum ; R. Brown in Benth. Fl. Austr. I, 284 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,284					

LINUM ; Tournefort, inst. 339, t. 176 (1700), from Theophrastos, Dioscorides & Plinius.

L. marginale ; Cunningham in Hook. Lond. Jour. Bot. VII, 169 (1848)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,283	M.fr.XI,27.				

GERANIACEAE.

A. L. de Jussieu, Gen. 268 (1789), from B. de Jussieu (1759).—Genera 5, Species 8.

GERANIUM ; Tournefort, inst. 266, t. 142 (1700), from Dioscorides and Plinius.

G. pilosum ; Solander in G. Forst., Fl. Ins. Austr. prodr. 91 (1786)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,296	M.fr.XI,27	...	Am., N.Z.		
G. sessiliflorum ; Cavanilles, Diss. 198, t. 77 (1790)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,297	...	...	Am., N.Z.		

ERODIUM ; L. Héritier, Geraniologia, t. 7 (1787).

E. cygnorum ; Nees in Lehm. pl. Preiss. I, 162 (1844)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,297	M.fr.IX,75 ; XI,20.				

PELARGONIUM ; L. Héritier, Geraniologia, t. 1 (1787).

P. Australe ; Willdenow, sp. pl. III, 675 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,298	M.fr.IX,77	...	Afr., N.Z.		
P. Rodueyanum ; Mitchell, Three Exped. II, 144 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,299					

OXALIS ; Linné, gen. pl. 134 (1737), from Plinius.

O. Magellanica ; G. Forster in Comm. Goetting. IX, 33 (1789)	—	—	T.	V.	—	—	—	B.fl.I,300	...	...	Am., N.Z.		
O. corniculata ; Linné, sp. pl. 435 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,301	M.fr.IX,75,77	E., As., Afr., Am., P., N.Z.			

BIOPHYTUM ; De Candolle, prodr. I, 689 (1824).

B. apodoscias ; Turczaninow in Bullet. des Nat. de Mosc. XXXVI, 595 (1863)	—	—	—	—	—	Q.	—	—					

MALVACEAE.

Adanson in Mém. Acad. fr. 224 (1761).—Genera 15, Species 110.

LAVATERA ; Tournefort in Act. Acad. Par. 86, t. 3 (1706).

L. plebeja ; Sims in Bot. Mag. t. 2269 (1821)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,185	M.fr.IX,75,130 ; X,53.				

MALVASTRUM ; Asa Gray in Mem. Amer. Acad. IV, 21 (1849). (Malva partly.)

M. spicatum ; A. Gray in Mem. Amer. Acad. IV, 21 (1849)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,187	M.fr.VI,170 ; IX,130	As., Afr., Am., P.
M. tricuspidatum ; A. Gray in Mem. Amer. Acad. IV, 21 (1849)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,187	... ...	As., Afr., Am., P.

PLAGIANTHUS; R. & G. Forster, char. gen. 85, t. 43 (1776). (Asterotrichon, Blepharanthemum, Lawrencia, Halothamnus, Gynothrix.)

P. sidoides ; Hooker, Bot. Mag. t. 3396 (1835)	—	—	T.	—	—	—	—	B.fl.I,188	M.fr.VI,169.	
P. pulchellus ; A. Gray, Bot. Amer. Expl. Exped. I, 181 (1854)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,189	M.fr.IX,130.	
P. spicatus ; Bentham in proceed. Linn. Soc. VI, 103 (1862)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,189	M.fr.VI,170 ; IX,130.	
P. glomeratus ; Bentham in proceed. Linn. Soc. VI, 103 (1862)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,190	M.fr.IX,130.	
P. diffusus ; Bentham, Fl. Austr. I, 190 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,190		
P. microphyllus ; F. v. M., fragm. I, 29 (1858)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,190	M.fr.VI,170 ; IX,13.	
P. Berthae ; F. v. M., fragm. V, 103 (1866)	W.A.	S.A.	—	—	—	—	—	—	M.fr.V.114.	

SIDA ; Linné, gen. pl. 205 (1737) from Theophrastos.

S. corrugata ; Lindley in Mitch. Three Exped. II, 13 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,192	M.fr.IX,75.		
S. Spenceriana ; F. v. M. in Wing's S. Sc. Record, April 1885	—	—	—	—	N.S.W.	Q.	—				
S. intricata ; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1860)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,193			
S. macropoda ; F. v. M. in Benth. Fl. Austr. I, 193 (1863)...	—	—	—	—	—	—	N.A.	B.fl.I,193			
S. virgata ; Hooker in Mitch. Trop. Austr. 361 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,194			
S. echinocarpa ; F. v. M., fragm. XI, 62 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,62.		
S. cryphiopetala ; F. v. M., fragm. III, 4 (1862)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,194	M.fr.III,4.		
S. petrophila ; F. v. M. in Linnaea XXV, 381 (1852)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,194			
S. Kingii ; F. v. M. in Vict. Natur. III, 138 (1887)	W.A.	—	—	—	—	—	—				
S. calyxhymenia ; J. Gay in De Candolle, prodr. I, 462 (1824)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,194	M.fr.IX,75.		
S. physocalyx ; F. v. M., fragm. III, 3 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.I,195	M.fr.X,73.		
S. subspicata ; F. v. M. in Benth. Fl. Austr. I, 195 (1863)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,195			
S. pleiantha ; F. v. M. in Benth. Fl. Austr. I, 195 (1863)	—	—	—	—	—	Q.	—	B.fl.I,195	M.fr.VI,169.		
S. spinosa ; Linné, sp. pl. 683 (1753)	—	—	—	—	—	Q.	N.A.	B fl.I,196	... ...	As., Afr., Am., P.	
S. rhombifolia ; Linné, sp. pl. 684 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,196	... ...	As., Afr., Am., P.	
S. cordifolia ; Linné, sp. pl. 684 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,196	... ...	As., Afr., Am., P.	
S. platycalyx ; F. v. M. in Benth. Fl. Austr. I, 197 (1863)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,197	M.fr.X,73.		
S. cleisocalyx ; F. v. M., fragm. X, 73 (1876)...	—	—	—	—	—	—	N.A.	—	M.fr.X,73.		
S. inclusa ; Bentham, Fl. Austr. I, 197 (1863)	—	S.A.	—	—	—	—	N.A.	B.fl.I,197	M.fr.IX,181 ; XI,32.		
S. cardiophylla ; F. v. M., fragm. VIII, 242 (1874)	W.A.	S.A.	—	—	—	—	N.A.	—	M.fr.VIII,242.		
S. Hookeriana ; Miquel in Lehm. pl. Preiss. I, 242 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,197			
S. lepida ; F. v. M., fragm. VI, 168 (1868)	W.A.	S.A.	—	—	—	—	N.A.	—	M.fr XI,32.		

ABUTILON ; Tournefort. inst. 99, t. 25 (1700) from Camerarius (1586). (Fleischera, Abutilaea.)

A. tubulosum ; Hooker in Mitch. Trop. Austr. 300 (1848)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,200	M.fr.II,12 ; IX,131.		
A. amplum ; Bentham, Fl. Austr. I, 200 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,209			
A. leucopetalum ; F. v. M. in Benth. Fl. Austr. I, 200 (1863)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,200	M.fr.II,12.		
A. Mitchelli ; Bentham, Fl. Austr. I, 201 (1863)	—	S.A.	—	—	—	Q.	—	B.fl.I,201			
A. micropetalum ; F. v. M. in Benth. Fl. Austr. I, 201 (1863)	—	—	—	—	—	Q.	—	B.fl.I,201	M.fr.VI.169 ; IX,131 ; XI,32 ; M.fr.IX,131.		
A. cryptopetalum ; F. v. M. in Benth. Fl. Austr. I, 201 (1863)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.I,201	M.fr.II,11 ; IX,13.		
A. geranioides ; Bentham, Fl. Austr. I, 202 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,202	M.fr.VI,168.		
A. otocarpum ; F. v. M. in Transact. phil. Soc. Vic. I, 13 (1854)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,202	M.fr.XI,32.		
A. longilobum ; F. v. M., fragm. IX, 130 (1875)	W.A.	—	—	—	—	—	N.A.	—	M.fr.IX,130.		
A. subviscosum ; Bentham, Fl. Austr. I, 202 (1863)	—	—	—	—	—	Q.	—	B.fl.I,202			
A. Indicum ; J. Bauhin, hist. plant. univ. II, 959 (1651)	—	—	—	—	—	Q.	N.A.	B.fl.I,202	... .	As., Afr.	
A. auritum ; G. Don, gen. syst. I, 500 (1831)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,203	M.fr.XI,32 ...	As., Afr., P.	

A. Avicennae; Gerard, gen. hist. pl. 790 (1597)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,203						E., As., Am.
A. graveolens; Wight & Arnott, Prodr. Fl. Pen. Ind. or. I, 56 (1840)	—	—	—	—	—	Q.	—	B.fl.I,204						As., Afr.
A. oxycarpum; F. v. M. in Benth. Fl. Austr. I, 204 (1863)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,204	M.fr.II,12; IX,131; XI,32.					
A. muticum; G. Don, gen. syst. I, 502 (1831)	—	—	—	—	—	Q.	—	B.fl.I,204						As., Afr.
A. Cunninghamii; Bentham, Fl. Austr. I, 205 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,205						
A. exonemum; F. v. M., fragm. XI, 63 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,63.					
A. Fraseri; Hooker in Walpera, annal. II, 158 (1851)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,205	M.fr.II,11.					
A. halophilum; F. v. M. in Linnaea XXV, 381 (1852)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,206	M.fr.XI,27.					
A. macrum; F. v. M., fragm. IX, 59 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,59.					
A. crispum; G. Don, gen. syst. I, 502 (1831)	—	—	—	—	—	—	N.A.	B.fl.I,206	M.fr.VI,169					As., Afr., Am., P.
A. Julianae; Endl. prodr. fl. ins. Norf. 75 (1833)	—	—	—	—	N.S.W.	—	—	—						
URENA; Dillenius, Hort. Eltham. 430, t. 319 (1732).														
U. lobata; Linné, sp. pl. 692 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,206	M.fr.VII,69; IX,130					As., Afr., Am., P.
U. Armitiana; F. v. M., fragm. X, 78 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,78.					
PAVONIA; Cavanilles, monadelph. dissert. II, App. II (1786). (Greevesia.)														
P. hastata; Cavanilles, dissert. III, 138, t. 47 (1786)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,207	M.fr.IX,130; X,73					As., Afr., Am.
HOWITTIA; F. v. M. in Transact. Vict. Inst. I, 116 (1855).														
H. trilocularis; F. v. M. in Transact. Vict. Inst. I, 116 (1855)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,198	M.fr.VI,169.					
HIBISCUS; Linné, syst. nat. (1735); gen. pl. 207 (1737). (Fugosia partly, Abelmoschus, Paritium.)														
H. ficulneus; Linné, sp. pl. 605 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,209	M.fr.I,67.					As.
H. Abelmoschus; Linné, sp. pl. 696 (1753)	—	—	—	—	—	—	N.A.	—	M.fr.VI,169					As., Afr., P.
H. rhodopetalus; F. v. M. in Benth. Fl. Austr. I, 209 (1863)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,209	M.fr.II,113; V,210; IX,229; XI,31.					
H. Manihot; Linné, sp. pl. 696 (1753)	—	—	—	—	—	Q.	—	B.fl.I,210						As.
H. Notho-Manihot; F. v. M., fragm. V, 57 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,213; IX,130.					
H. Trionum; Linné, sp. pl. 697 (1753)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,210	M.fr.II,115; IX,129; XI,31.					E.,As.,Afr.,N.Z.
H. insularis; Endlicher, prodr. fl. ins. Norf. 74 (1833)	—	—	—	—	N.S.W.	—	—	—						
H. brachysiphonius; F. v. M., fragm. I, 67 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,210	M.fr.I,220,243; VI,169.					
H. Drummondii; Turczaninow in Bull. Soc. Mosc. XXXI, 195 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,211	M.fr.I,220; IX,129; XI,31.					
H. microlaenus; F. v. M., fragm. II, 116 (1861)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,211	M.fr.III,5; IX,129.					
H. Pinonianus; Gaudichaud in Freyc. voy. Bot. 476, t. 100 (1826)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,211	M.fr.II,116.					[As., Afr.
H. cannabinus; Linné, sp. pl. ed. sec., 979 (1763)	—	—	—	—	—	Q.	N.A.	B.fl.I,212	M.fr.II,117; III,166; VI,169; IX,129; XI, 31					
H. divaricatus; Graham in Edinb. phil. Journ. (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,212	M.fr.II,118; IX,129; XI,31.					
H. Fitzgeraldii; F. v. M., fragm. VIII, 242 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,242.					
H. Elaworthii; F. v. M., fragm. VIII, 241 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,241.					
H. Haynaldi; F. v. M. in Wing's S. Sc. Rec. March (1883)	W.A.	—	—	—	—	—	—	—						
H. heterophyllus; Ventenat, jard. de la Malm., t. 103 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,212	M.fr.VI,170; XI,31.					
H. diversifolius; N. Jacquin, icon. pl. rar. 551 (1789)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,213	M.fr.II,117; VI,169					As., Afr., P.
H. splendens; Fraser in Edinb. phil. journ. (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,213	M.fr.VI,170; XI,31&123.					
H. zonatus; F. v. M., fragm. I, 221 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,213	M.fr.VI,170.					
H. Coatesii; F. v. M., fragm. III, 5 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.III,5.					
H. leptocladus; Bentham, Fl. Austr. I, 214 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.VI,169; IX,129.					
H. setulosus; F. v. M., fragm. I, 221 (1859)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.IX,31.					
H. Goldsworthii; F. v. M., fragm. XI, 30 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.IX,30.					
H. pentaphyllus; F. v. M., fragm. II, 13 (1860)	—	—	—	—	—	—	N.A.	B.fl.I,214	M.fr.VI,169; XI,32.					As., Afr.
H. geranioides; Cunningham in Benth. Fl. Austr. I, 215 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,215						
H. vitifolius; Linné, sp. pl. 696 (1753)	—	—	—	—	—	Q.	—	B.fl.I,215	M.fr.I,220; II,114; VI,169.					As., Afr.
H. panduriformis; N. Burmann, Fl. Ind. 151, t. 47 (1768)	W.A.	—	—	—	—	Q.	N.A.	B.fl.I,215	M.fr.II,115; VI,169; IX,129.					As., Afr.

H. Normani ; F. v. M., fragm. III, 4 (1862) ...	—	—	—	—	—	Q.	—	B.fl.I,216	M.fr.V,44 ; VI,169 ; IX,128.					
H. Krichauffii ; F. v. M., Rep. Babb. Exped. 7 (1858)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,216	M.fr.VI,169 ; IX,75,129.					
H. Farragei ; F. v. M., fragm. VIII, 241 (1874)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	—	M.fr.VIII,284 ; IX,129&196 ; XI,32,134.					
H. Sturtii ; Hooker in Mitch. trop. Austr. 363 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,216	M.fr.II,13 ; IX,128.					
H. phyllochlaenus ; F. v. M., fragm. IX, 128 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,129.					
H. Huegelii ; Endlicher in Hueg. enum. 10 (1837)	W.A.	S.A.	—	—	—	—	—	B.fl.I,217	M.fr.IX,128.					
H. Wrayae ; Lindley, bot. Regist. t. 69 (1840)	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,128.					
H. tiliaceus ; Linné, sp. pl. 694 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,218	M.fr.VI,170 ; IX,129 ; XI,32.	As., Afr.,				
H. cuneiformis ; De Candolle. prodr. I, 454 (1824)	W.A.	—	—	—	—	—	N.A.	B.fl.I,219	M.fr.XI,32.	[Am., P.				
H. hakeaefolius ; Giordano, Mém. di nuov. sp. d'Ibisco (1833)	W.A.	S.A.	—	—	—	—	—	B.fl.I,220	M.fr.IX,128.					
LAGUNARIA ; G. Don, gen. syst. dichlam. pl. I, 485 (1831).														
L. Patersoni ; G. Don, gen. syst. I, 485 (1831)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,218	M.fr.IX,130 ; XI,32.					
THESPESIA ; Solander in Ann. du. Mus. IX, 290 (1807).														
T. populnea ; Solander in Ann. du Mus. IX, 290, t. 8 (1807)	—	—	—	—	—	Q.	N.A.	B.fl.I,221	M.fr.VI,170 ; IX,130 ; XI,32.	As., Afr., P.				
GOSSYPIUM ; Linné, gen. pl. 206 (1737) from Camerarius (1586). (Fugosia partly, Sturtia.)														
G. Sturtii ; F. v. M., fragm. III, 6 (1862)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.I,222	M.fr.IV,174 ; VI,169&251 ; IX,127 ; XI,63.					
G. Robinsoni ; F. v. M., fragm. IX, 126 (1875)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,63&135.					
G. Australe ; F. v. M., fragm. I, 46 (1858)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,220	M.fr.III,6 ; V,44 ; IX,127 ; XI,32&63.					
G. thespesioides ; F. v. M., fragm. IX, 127 (1875)	—	—	—	—	—	—	N.A.	B.fl.I,220	M.fr.V,44&177 ; VI,169 ; XI,32&63.					
G. Cunninghamii ; Todaro, Monogr. Gen. Gossyp. 7 (1878)	—	—	—	—	—	—	N.A.	B.fl.I,220	M.fr.VI,169 ; IX,129 ; XI,32&63.					
G. costulatum ; Todaro, Monogr. Gen. Gossyp. 7 (1878)	—	—	—	—	—	—	N.A.	B.fl.I,221	M.fr.IX,127 ; XI,32&63.					
G. flaviflorum ; F. v. M., fragm. IX, 127 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.V,44 ; XI,63.					
G. populifolium ; F. v. M., fragm. IX, 127 (1875)	—	—	—	—	—	—	N.A.	B.fl.I,221	M.fr.XI,32&63.					
ADANSONIA ; Linné, sp. pl. I, 1190 (1753).														
A. Gregorii ; F. v. M. in Hook. Kew misc. IX, 14 (1857)	—	—	—	—	—	—	N.A.	B.fl.I,223	M.fr.VII,40 ; XI,32.					
CAMPTOSTEMON ; Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876).														
C. Schultzii ; Masters in J. Hook. icon. pl. XII, 18, t. 1119 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.IX,43 ; XI,32.					
BOMBAX ; Linné, sp. pl. I, 511 (1753).														
B. Malabarica ; De Candolle, prodr. I, 479 (1824)	—	—	—	—	—	Q.	N.A.	B.fl.I,223	M.fr.XI,32	As.				

STERCULIACEAE.

VENTENAT ; Jard. de la Malm. II, 91 (1804).—Genera 20, Species 125.

STERCULIA ; Linné, Fl. Zeyl. 166 (1747).														
S. quadrifida ; R. Brown in Benn. pl. Jav. rar. 233 (1852)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,227	M.fr.VI,173.					
S. laurifolia ; F. v. M., fragm. VI, 172 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,172.					
BRACHYCHITON ; Schott & Endlicher, Melet. bot. 34 (1832). (Trichosiphon, Poecilodermis, Delabechea.)														
B. paradoxus ; Schott, Meletem. 34 (1832)	—	—	—	—	—	—	N.A.	B.fl.I,227						
B. Bidwilli ; Hooker, Bot. Mag. t. 5133 (1859)	—	—	—	—	—	Q.	—	B.fl.I,228						
B. discolor ; F. v. M., fragm. I, 1 (1858)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,228	M.fr.I,242 ; II,177.					
B. incanus ; R. Brown in Benn. pl. Jav. rar. 234 (1852)	—	—	—	—	—	—	N.A.	B.fl.I,228						
B. luridus ; C. Moore in F. v. M., fragm. I, 1 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,228	M.fr.II,177.					
B. platanoides ; R. Brown in Benn. pl. Jav. rar. 234 (1852)	W.A.	—	—	—	—	Q.	N.A.	B.fl.I,229	M.fr.VI,173.					
B. acerifolius ; F. v. M., fragm. I, 1 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,229	M.fr.II,177 ; V,209 ; VI,173.					

B. diversifolius; R. Brown in Benn. pl. Jav. rar. 234 (1852)	W.A.	—	—	—	—	Q.	N.A.	B.fl.I,230						
B. populneus; B. Brown in Benn. pl. Jav. rar. 234 (1852)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,229	M.fr.IV,137.					
B. populneo-acerifolius; F. v. M. in proceed. L.S. of N.S.W. IX, 380 (1884) ...	—	—	—	—	N.S.W.	—	—	—						
B. Delabechii; F. v. M., pl. of Vict. I, 157 (1860)	—	—	—	—	—	Q.	—	B.fl.I,230	M.fr.IV,137.					
B. Gregorii; F. v. M. in Hook. Kew misc. IX, 199 (1857)...	W.A.	S.A.	—	—	—	—	—	—	M.fr.IV,137; V,212; X,53.					
TARRIETIA; Blume, Bijdr. 227 (1825). (Argyrodendron.)														
T. Argyrodendron; Bentham, Fl. Austr. I, 230 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,230	M.fr.VI,173; IX,42.					
T. trifoliolata; F. v. M., fragm. IX, 43 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.I,2; II,177.					
HERITIERA; Dryander in Aiton, Hort. Kew III, 546 (1789).														
H. litoralis; Dryander in Aiton, Hort. Kew III, 546 (1789)	—	—	—	—	—	Q.	N.A.	B.fl.I,231	M.fr.VI,173 ...	As., Afr., P.				
UNGERIA; Schott & Endlicher, Melet. 27—31 (1832).														
U. floribunda; Schott & Endlicher, Melet. 27, t. 4 (1832)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.					
HELICTERES; Plukenet, Phytogr. 181, t. 245 (1692).														
H. Isora; Linné, sp. pl. 963 (1753)	—	—	—	—	—	—	N.A.	B.fl.I,232	M.fr.VI,173 ...	As.				
METHORIUM; Schott & Endlicher, Melet. 29—30 (1832).														
M. spicatum; F. v. M., syst. Cens. of Austr. pl. 15 (1882) from Colebr. (1831)	—	—	—	—	—	Q.	—	—	M.fr.VII,39 ...	As.				
M. canum; Schott, Meletem. 29, t. 5 (1832)	—	—	—	—	—	—	N.A.	B.fl.I,232						
M. dentatum; F. v. M. in Benth. Fl. Austr. I, 232 (1863)...	—	—	—	—	—	—	N.A.	B.fl.I,232						
M. semiglabrum; F. v. M., fragm. V, 43 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,43.					
MELHANIA; Forskael, Fl. Aeg. Arab. 64 (1775).														
M. incana; Heyne in Wall. numerit. list 1200 (1828)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,234	M.fr.I,69; VI,174	As., Afr., P.				
MELOCHIA; Dillenius, Hort. Eltham. 221, t. 176 (1732). (Riedleya.)														
M. pyramidata; Linné, sp. pl. 674 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,234	M.fr.VI,173. ...	As., Afr., Am., P.				
M. corchorifolia; Linné, sp. pl. 675 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,235	M.fr.VI,173. ...	As., Afr.				
HERMANNIA; Tournefort, inst. 656, t. 432 (1700). (Gilesia, Mahernia partly.)														
H. Gilesii; F. v. M., fragm. IX, 42 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,42.					
DICARPIDIUM; F. v. M. in Hook. Kew Misc. IX, 302 (1857).														
D. monoicum; F. v. M. in Hook. Kew misc. IX, 302 (1857)	—	—	—	—	—	—	N.A.	B.fl.I,235						
WALTHERIA; Linné, gen. pl. 203 (1737).														
W. Indica; Linné, sp. pl. 637 (1753)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,236	M.fr.VI,173. ...	As., Afr. Am., P.				
ABROMA; N. Jacquin, Hort. Vindob. III, t. 1 (1776).														
A. fastuosa; Salisbury, Parad. Lond. 102 (1807)	—	—	—	—	—	Q.	—	B.fl.I,236	M.fr.VI,173 ...	As.				
COMMERCONIA; R. & G. Forster, char. gen. 43, t. 22 (1776). (Ruelingia, Achilleopsis.)														
C. salvifolia; Bentham, Fl. Austr. I, 238 (1863)	—	—	—	—	—	Q.	—	B.fl.I,238						
C. dasyphylla; Andrews, Bot. Reposit. t. 603 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,238	M.fr.X,22.					
C. rugosa; F. v. M., syst. Cens. of Austr. pl. 16 (1882)	—	—	—	—	N.S.W.	—	—	B.fl.I,238						
C. Preissii; Steudel in Lehm. pl. Preiss. I, 237 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,339						
C. cinerea; Steudel in Lehm. pl. Preiss. I, 237 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,239						
C. cygnorum; Steudel in Lehm. pl. in Preiss. I, 238 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,239						
C. platycalyx; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Benth. (1863)	W.A.	—	—	—	—	—	—	B.fl,I,240						
C. parviflora; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Endl. (1837) ...	W.A.	—	—	—	—	—	—	B.fl.I,240						
C. hermanniaefolia; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Steetz (1847)	—	—	—	—	N.S.W.	—	—	B.fl.I,240						
C. Kempeana; F. v. M., fragm. IX, 113 (1881)	—	S.A.	—	—	—	—	—	—	M.fr.XI,113.					
C. loxophylla; F. v. M., fragm. X., 22 (1876)	—	S.A.	—	—	—	—	N.A.	B.fl.I,240	M.fr.I,68					

C. magniflora ; F. v. M., fragm. VIII, 223 (1874) from Turcz. (1852)	—	S.A.	—	—	—	—	—	—	—	M.fr.X,22 ; XI,114.	
C. cuneata ; F. v. M., syst. Cens. of Austr. pl. 16 (1882) from Turcz. (1852) ...	W.A.	—	—	—	-	—	—	B.fl.I,241			
C. rotundifolia ; F. v. M., syst. Cens. of Austr. pl. 16 (1882)	W.A.	—	—	—	—	—	—	B.fl.I,241			
C. densiflora ; F. v. M., fragm. X, 21 (1876)	W.A.	—	—	—	—	—	—	B.fl.I,241	M.fr.VI,175 ; X,22 ; XI,114.		
C. Fraseri ; J. Gay in Mém. du Mus. Par. X, 215, t. 15 (1823)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,242			
C. Leichhardtii ; Bentham, Fl. Austr. I, 242 (1863)...	—	—	—	—	—	Q.	—	B.fl.I,242	M.fr.VI,175 ; XI,114.		
C. echinata ; R. & G. Forster, char. gen. 43, t. 22 (1776)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,243	M.fr.VI,175 ; X,22 ...	As., P.	
C. Gaudichaudi ; J. Gay in Mém. du Mus. Par. X, 213, t. 14 (1823)	W.A.	—	—	—	—	—	—	B.fl.I,243			
C. crispa ; Turczaninow in Bull. Soc. Mosc. XIX, 501 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,243			
C. melanopetala ; F. v. M., fragm. X, 21 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,21.		
C. pulchella ; Turczaninow in Bull. Soc. Mosc. XIX, 502 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,244	M.fr.X,22.		
C. microphylla ; Bentham, Fl. Austr. I, 244 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,244			
C. craurophylla ; F. v. M., fragm. IX, 59 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,59.		
C. Tatei ; F. v. M. in Trans. Roy. Soc. of S. Austr. X (1888)	—	S.A.	—	—	—	—	—	—			
HANNAFORDIA ; F. v. M., fragm. II, 9 (1860).											
H. quadrivalvis ; F. v. M., fragm. II, 9 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,248	M.fr.X,96 ; XI,20&115		
H. Shanesii ; F. v. M., fragm. VI, 175 (1868)	—	—	—	—	—	Q.	—	—	M.fr.X,96.		
H. Bissillii ; F. v. M., fragm. X, 95 (1877)	W.A.	S.A.	—	—	N.S.W.	—	—	—	M.fr.XI,115.		
SERINGEA ; Sprengel, Anleit. II, 649 (1818). (Keraudrenia.)											
S. platyphylla ; J. Gay in Mém. du Mus. Par. VII, 443 (1821)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,244	M.fr.II,5 ; VI,175 ; X,97.		
S. lanceolata ; Steetz in Lehm. pl. Preiss. II, 349 (1847)	—	—	—	—	—	Q.	—	B.fl.I,245	M.fr.I,142 ; II,5 ; V,44 ; VI,174 ; X,97.		
S. corollata ; Steetz in Lehm. pl. Preiss. II, 330 (1847)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,246	M.fr.I,28,142 ; II,5 ; VI,174 ; X,96.		
S. Hillii ; F. v. M., syst. Cens. of Austr. pl. 16 (1882)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,246	M.fr.VI,174 ; X,97 ...	Afr.	
S. nephrosperma ; F. v. M. in Hook. Kew misc. IX, 15 (1857)	—	S.A.	—	—	—	—	N.A.	B.fl.I,246	M.fr.I,142 ; II,5 ; X,97.		
S. microphylla ; F. v. M., fragm. II, 5 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,247	M.fr.II,5.		
S. adenolasia ; F. v. M., fragm. X, 96 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,96.		
S. integrifolia ; F. v. M., fragm. II, 5 (1860)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.I,247	M.fr.VI,175 ; X,97.		
THOMASIA ; J. Gay in Mém. du Mus. VII, t. 6—7 (1821). (Leucothamnus, Rhynchostemon.)											
T. macrocarpa ; Huegel in Endlicher, nov. stirp. dec. 32 (1839)	W.A.	—	—	—	—	—	—	B.fl.I,250			
T. rugosa ; Turczaninow in Bull. Soc. Mosc. XIX, 501 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,250			
T. montana ; Steudel in Lehm. pl. Preiss. I, 230 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,250			
T. tenuivestita ; F. v. M., fragm. II, 7 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,251	M.fr.II,7.		
T. solanacea ; J. Gay in Mém. du Mus. VII, 456, t. 21 (1821)	W.A.	—	—	—	—	—	—	B.fl.I,251			
T. brachystachys ; Turczaninow in Bull. Soc. Mosc. XXV, 143 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,251			
T. discolor ; Steudel in Lehm. pl. Preiss. I, 233 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,251			
T. quercifolia ; J. Gay in Mém. du Mus. VII, 459, t. 21 (1821)	W.A.	—	—	—	—	—	—	B.fl.I,252			
T. foliosa ; J. Gay in Mém. du Mus. VII, 454, t. 22 (1821)...	W.A.	—	—	—	—	—	—	B.fl.I,252			
T. triloba ; Turczaninow in Bull. Soc. Mosc. XIX, 500 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,252			
T. triphylla ; J. Gay in Mém. du Mus. VII, 458 (1821)	W.A.	—	—	—	—	—	—	B.fl.I,252			
T. purpurea ; J. Gay in Mém. du Mus. VII, 452, t. 21 (1821)	W.A.	—	—	—	—	—	—	B.fl.I,253			
T. macrocalyx ; Steudel in Lehm. pl. Preiss. I, 230 (1844)...	W.A.	—	—	—	—	—	—	B.fl.I,253	M.fr.,VII,175.		
T. pauciflora ; Lindley Bot. Regist. XXV, App. XVII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,253			
T. rhynchocarpa ; Turczaninow in Bull. Soc. Mosc. XXV, 142 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,254	M.fr.II,8 ; VI,175.		
T. grandiflora ; Lindley, Bot. Regist. XXV, App. XVIII (1839)	W.A.	—	—	—	—	—	—	B.fl.I,254	M.fr.II,8.		
T. cognata ; Steudel in Lehm. pl. Preiss. I, 232 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,254			
T. rulingioides ; Steudel in Lehm. pl. Preiss. I, 232 (1844)...	W.A.	—	—	—	—	—	—	B.fl.I,255			

T. angustifolia; Steudel in Lehm. pl. Preiss. I, 232 (1844)...	W.A.	—	—	—	—	—	—	B.fl.I,255	
T. petalocalyx; F. v. M. in Transact. phil. Soc. Vict. I, 35 (1854)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,255	
T. Sarotes; Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,255	
GUICHENOTIA; J. Gay in Mém. du Mus. VII, 448, t. 5 (1821). (Sarotes, Ditomostrophe.)									
G. Iedifolia; J. Gay in Mém. du Mus. VII, 449, t. 5 (1821)	W.A.	—	—	—	—	—	—	B.fl.I,258	M.fr.VII,174; XI,114.
G. macrantha; Turczaninow in Bull. Soc. Mosc. XIX, 500 (1846)	W.A.	—	—	—	—	—	—	B.fl.I,258	
G. semihastata; Bentham, Fl. Austr. I, 258 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,258	M.fr.II,4,7.
G. Sarotes; Bentham, Fl. Austr. I, 258 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,258	
G. micrantha; Bentham, Fl. Austr. I, 258 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,258	M.fr.II,7.
LASIOPETALUM; Smith in Transact. Linn. Soc. IV, 216 (1789). (Corethrostylis, Asterochiton.)									
L. laxiflorum; F. v. M., fragm. XI, 112 (1881)	W.A.	—	—	—	—	—	—	B.fl.I,256	M.fr.XI,112.
L. pygmalum; Bentham, Fl. Austr. I, 256 (1863) implied...	W.A.	—	—	—	—	—	—	B.fl.I,257	M.fr.XI,113.
L. stelligerum; Bentham, Fl. Austr. I, 256 (1863) implied...	W.A.	—	—	—	—	—	—	B.fl.I,256	M.fr.XI,113.
L. glutinosum; F. v. M., fragm. XI, 113 (1881)	W.A.	—	—	—	—	—	—	B.fl.I,256	M.fr.XI,113.
L. discolor; Hooker, Comp. to Bot. Mag. I, 276 (1835)	W.A.	S.A.	T.	—	—	—	—	B.fl.I,260	M.fr.XI,110.
L. dasyphyllum; Sieber in Hook. journ. of bot. II, 414 (1840)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,261	M.fr.VI,174; XI,110.
L. indutum; Steudel in Lehm. pl. Preiss. I, 235 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,261	M.fr.XI,110.
L. Maxwelli; F. v. M., fragm. XI, 107 (1881)	W.A.	—	—	—	—	—	—	—	M.fr.XI,107.
L. Ogilvieanum; F. v. M., fragm. XI, 107 (1881)	W.A.	—	—	—	—	—	—	—	M.fr.XI,107.
L. Behrii; F. v. M. in Transact. phil. Soc. Vict. I, 36 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,261	
L. parviflorum; Rudge in Transact. Linn. Soc. X, 297, t. 19 (1811)	—	—	—	V.	N.S.W.	—	—	B.fl.I,262	M.fr.II,7; VI,174.
L. parvuliflorum; F. v. M., fragm. VI, 174 (1868)	W.A.	—	—	—	—	—	—	—	M.fr.VI,174.
L. Fitzgibbonii; F. v. M. in South. Sc. Record II, Apr. (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII.
L. oppositifolium; F. v. M., fragm. II, 5 (1860)	W.A.	—	—	—	—	—	—	B.fl.I,262	M.fr.XI,112.
L. micranthum; J. Hooker, Fl. Tasman. I, 51 (1860)	—	—	T.	—	—	—	—	B.fl.I,262	M.fr.XI,110.
L. macrophyllum; Graham in Bot. Mag. t. 3098 (1844)	—	—	—	—	N.S.W.	—	—	B.fl.I,262	
L. Baueri; Steetz in Lehm. pl. Preiss. II, 339 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,263	M.fr.VI,174; XI,110.
L. Tepperi; F. v. M., fragm. XI, 109 (1881)	—	S.A.	—	—	—	—	—	—	M.fr.IX,109.
L. rufum; R. Brown in Benth. Fl. Austr. I, 263 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,263	
L. ferrugineum; Smith in Bot. Reposit. t. 308 (1799)	—	—	—	V.	N.S.W.	—	—	B.fl.I,263	
L. acutiflorum; Turczaninow in Bull. Soc. Mosc. XXV, 145 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,264	M.fr.XI,109.
L. Oldfieldii; F. v. M., fragm. II, 6 (1860)	W.A.	—	—	—	—	—	—	—	M.fr.II,6.
L. quinquenervium; Turczaninow in Bull. Soc. Mosc. XXV, 146 (1852) ...	W.A.	—	—	—	—	—	—	—	M.fr.XI,109.
L. Drummondii; Bentham, Fl. Austr. I, 264 (1863)...	W.A.	—	—	—	—	—	—	B.fl.I,264	M.fr.VI,174; XI,112.
L. rosmarinifolium; Bentham, Fl. Austr. I, 264 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,264	M.fr.XI,112.
L. cordifolium; Endlicher in Hueg. enum. 10 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,265	M.fr.VI,174; XI,112.
L. Schulzenii; F. v. M., pl. of Vict. I, 145 (1861) implied...	—	S.A.	—	V.	—	—	—	B.fl.I,265	
L. florihundum; Bentham, Fl. Austr. I, 265 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,265	M.fr.XI,112.
L. molle; Bentham, Fl. Austr. I, 265 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,265	M.fr.XI,111.
L. membranaceum; Bentham, Fl. Austr. I, 266 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,266	M.fr.XI,112.
L. bracteatum; Bentham, Fl. Austr. I, 266 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,266	M.fr.XI,111.
LYSIOSEPALUM; F. v. M., fragm. I, 143 (1859). (Lasiopetali subgenus.)									
L. Barryanum; F. v. M., fragm. I, 143 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,267	M.fr.VI,174.
L. rugosum; Bentham, Fl. Austr. I, 267 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,267	

TILIACEAE.

A. L. DE JUSSIEU; Gen. 289 (1789).—Genera 7, Species 56.

ARISTOTELIA; L'Héritier, stirp. II, 21, t. 16 (1784). (Friesia.)												
A. peduncularis; J. Hooker, Fl. Nov. Zel. I, 33 (1853)	—	—	T.	—	—	—	—	B.fl.I,280	M.fr.II,80; VIII,2.			
A. Australasica; F. v. M., fragm. II, 79 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,280	M.fr.VIII,2.			
A. megalosperma; F. v. M., fragm. IX, 84 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,84.			
ELAEOCARPUS; Burmann, thesaur. Zeil. 39, t. 40 (1737). (Elaiocarpus, Dicera.)												
E. Bancroftii; F. v. M. & Bailey in proceed. Roy. Soc. of Q. II, pt. II (1885)	—	—	—	—	—	Q.	—	—				
E. holopetalus; F. v. M., fragm. II, 143 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.I,281	M.fr.IV,173; V,211; VI,172; VIII, 2.			
E. Arnhemicus; F. v. M., Decum. intercol. Exhib. 24 (1867)	—	—	—	—	—	—	N.A.	—	M.fr.X,4; VI,172.			
E. obovatus; G. Don, gen. syst. dichl. pl. I, 599 (1831)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,281	M.fr.II,80; VI,172; IX,85.			
E. reticulatus; Smith in Rees Cycl. XII (1809)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,281	M.fr.VI,172; VIII,2.			
E. grandis; F. v. M., fragm. II, 81 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,281	M.fr.II,175; VI,172; VIII,12.			
E. foveolatus; F. v. M., fragm. V, 157 (1866)	—	—	—	—	—	Q.	—	—	M.fr.VI,172; VIII,2.			
E. ruminatus; F. v. M., fragm. VIII, 1 (1872)	—	—	—	—	—	Q.	—	—	M.fr.X,4.			
E. Grahami; F. v. M., fragm. X, 3 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,3.			
E. sericopetalus; F. v. M., fragm. VI, 171 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,171.			
SLOANEA; Linné, hort. Cliffort. 210 (1737). (Echinocarpus.)												
S. Australis; F. v. M., fragm. IV, 91 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,279	M.fr.V,29; VI,172; VIII,2.			
S. Langii; F. v. M., fragm. V, 28 (1865)	—	—	—	—	—	Q.	—	—	M.fr.VIII,2.			
S. Macbrydei; F. v. M., fragm. VI, 170 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,170.			
S. Woollsii; F. v. M., fragm. VI, 171 (1868)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII,2.			
BERRYA; Roxburgh, Pl. Coromand. III, 60, t. 264 (1819).												
B. ammonilla; Roxburgh, pl. Coromand. III, 60, t. 264 (1819)	—	—	—	—	—	Q.	—	B.fl.I,268	...	...	...	As.
CORCHORUS; Tournefort, inst. 259, t. 135 (1700).												
C. Rothii; Triumfetta pilosa, Roth, Nov. pl. sp. 223 (1821)	—	—	—	—	—	Q.	—	—	M.fr.IV,28; VIII,5		As., Afr.	
C. Elderi; F. v. M. in transact., R. S. of S.A. IX, 58 (1886)	—	S.A.	—	—	—	—	N.A.	—				
C. echinatus; Bentham, Fl. Austr. I, 276 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,276	M.fr.III,8; VIII,5.			
C. hygrophilus; Cunningham in Fl. Austr. I, 276 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,276	M.fr.VIII,5.			
C. Cunninghamii; F. v. M., fragm. III, 8 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,276	M.fr.III,167; VI,172; VIII,3.			
C. olitorius; Linné, sp. pl. 529 (1753)	—	—	—	—	—	—	N.A.	B.fl.I,276	M.fr.VIII,5	...	As., Afr.	
C. tridens; Linné, mantiss. alter. 566 (1771)	—	—	—	—	—	—	N.A.	B.fl.I,276	...	...	...	As., Afr.
C. acutangulus; Lamarck, encycl. meth. II, 104 (1786)	—	—	—	—	—	Q.	N.A.	B.fl.I,277	M.fr.VIII,6	...	As., Afr.	
C. fascicularis; Lamarck, encycl. meth. II, 104 (1786)	—	—	—	—	—	—	N.A.	B.fl.I,277	...	...	...	As., Afr.
C. pumilio; Bentham, Fl. Austr. I, 277 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,277	...	...	...	As., Afr.
C. vermicularis; F. v. M., fragm. III, 10 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,277	M.fr.VIII,5.			
C. tomentellus; F. v. M., fragm. III, 10 (1862)	—	—	—	—	—	Q.	—	B.fl.I,278	M.fr.III,10.			
C. sidoides; F. v. M., fragm. III, 9 (1862)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.I,278	M.fr.VIII,6; XI,62.			
C. leptocarpus; Cunningham in Benth. Fl. Austr. I, 278 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,278				
C. elachocarpus; F. v. M., fragm. VIII, 6 (1872)	—	—	—	—	—	—	N.A.	—	M.fr.III,6.			
C. Walcotti; F. v. M., fragm. III, 9 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,278	M.fr.VI,172; VIII,6; XI,62.			
C. trilocularis; Linné, mantiss. 77 (1771)	—	—	—	—	—	Q.	—	—	M.fr.VIII,5.	...	As., Afr.	
TRIUMFETTA; Plumier, nov. gen. 40, t. 8 (1703).												
T. procumbens; G. Forster, prodr. fl. insul. Austr. 35 (1786)	—	—	—	—	—	Q.	—	B.fl.I,273	M.fr.III,8; VI,172.		As., P.	
T. appendiculata; F. v. M., fragm. III, 7 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,273	M.fr.VIII,5.			

T. chaetocarpa; F. v. M., fragm. XI, 61 (1879)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,61.					
T. glaucescens; R. Brown in Benth. Fl. Austr. I, 273 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,273						
T. denticulata; R. Brown in Benth. Fl. Austr. I, 273 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,274						
T. micracantha; F. v. M., fragm. III, 7 (1862)	—	—	—	—	—	—	N.A.	B.fl.I,274	M.fr.VIII,5; XI,62.					
T. leptacantha; F. v. M., fragm. XI, 62 (1879)	—	—	—	—	—	—	N.A.	—	M fr.XI,62.					
T. plumigera; F. v. M., fragm. I, 69 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,274	M.fr.III,8; VI,172.					
T. parviflora; Bentham, Fl. Austr. I, 274 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,274						
T. Winneckeana; F. v. M. in Winnecke's Report (1885)	—	S.A.	—	—	—	Q.	N.A.	—						
T. rhomboidea; N. Jacquin, select. stirp. Arn. hist. 147, t. 90 (1763)	—	—	—	—	—	Q.	—	—						As., Afr., Am., P.
T. Johnstonii; F. v. M. in Melb. Chem. Feb. (1885)	W.A.	—	—	—	—	—	—	—						
GREWIA; Linné, syst. nat. 9 (1735); gen. pl. 276 (1737). (Nettoa.)														
G. orientalis; Linné, sp. pl. 964 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,270						As.
G. laevigata; Vahl, symbol. Bot. I, 34 (1790)	—	—	—	—	—	Q.	N.A.	B.fl.I,270	M.fr.VI,172					As., Afr., P.
G. breviflora; Bentham, Fl. Austr. I, 270 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,270						
G. latifolia; F. v. M. in Benth. Fl. Austr. I, 271 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,271	M.fr.VI,172; VIII,4.					As.
G. polygama; Roxburgh, Fl. Ind. II, 588 (1832)	—	—	—	—	—	Q.	N.A.	B.fl.I,271						As.
G. xanthopetala; F. v. M. in Benth. Fl. Austr. I, 271 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,271						
G. scabrella; Bentham, Fl. Austr. I, 272 (1863)	—	—	—	—	—	Q.	—	B.fl.I,272						
G. orbifolia; F. v. M. in Benth. Fl. Austr. I, 272 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,272						
G. pleiostigma; F. v. M., fragm. VIII, 4 (1872)	—	—	—	—	—	Q.	—	—	M.fr.VIII,4.					

EUPHORBIACEAE.

A. L. de Jussieu, Gen. 384 (1789) from B. de Jussieu (1759).—Genera 39, Species 226.

CALYCOPEPLUS; Planchon in Bull. de la Soc. Bot. de Fr. VIII, 31 (1861).														
C. ephedroides; Planchon in Bull. Soc. Bot. Fr. VIII, 31 (1861)	W.A.	—	—	—	—	—	—	B.fl.VI,53						
C. marginatus; Bentham, Fl. Austr. VI, 53 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,53						
EUPHORBIA; Linné, gen. pl. 152 (1737) from Plinius.														
E. Atoto; G. Forster, florul. ins. Austr. prodr. 36 (1786)	—	—	—	—	—	Q.	N.A.	B.fl.VI,46						As., P.
E. obliqua; Bauer in Endl. fl. Norf. 85 (1833)	—	—	—	—	N.S.W.	—	—	—						
E. Sparmanni; Boissier, Cent. Euphorb. 5 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.VI,46	M.fr.IX,77					P.
E. Mitchelliana; Boissier in De Candolle, prodr. XV, 25 (1866)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VI,47	M.fr.VI,182.					
E. schizolepis; F. v. M. in De Candolle, prodr. XV, pt. II, 20 (1862)	—	—	—	—	—	Q.	N.A.	B.fl.VI,47						
E. Schultzii; Bentham in Fl. Austr. VI, 47 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,47						
E. Armstrongiana; Boissier in De Candolle, prodr. XV, 47 (1866)	—	—	—	—	—	—	N.A.	B.fl.VI,48						
E. erythrantha; F. v. M., fragm. II, 152 (1861)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,48	M.fr.XI,64.					
E. Muelleri; Boissier in De Candolle, prodr. XV, 27 (1866)	—	—	—	—	—	—	N.A.	B.fl.VI,48						
E. Drummondii; Boissier, Cent. Euphorb. 14 (1860)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,49	M.fr.VI,182; XI,64.					
E. alsiniflora; Baillon, Adans. VI, 288 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.VI,49						
E. Wheeleri; Baillon, Adans. VI, 286 (1866)	—	S.A.	—	—	—	—	N.A.	B.fl.VI,49						
E. myrtoides; Boissier in De Candolle, prodr. XV, 15 (1866)	W.A.	—	—	—	—	—	N.A.	B.fl.VI,50						
E. micradenia; Boissier in De Candolle, prodr. XV, 27 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.VI,50						
E. Macgillivrayi; Boissier in De Candolle, prodr. XV, 26 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,50						
E. serrulata; Reinwardt in Blum. Bijdrag. 635 (1826)	—	—	—	—	—	Q.	N.A.	B.fl.VI,51						As.
E. filipes; Bentham, Fl. Austr. VI, 51 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,51						
E. Careyi; F. v. M., fragm. XI, 64 (1879)	W.A.	—	—		—	—	N.A.	—	M.fr.XI,64					

E. pilulifera; Linné, dissert. de Euphorb. 19 (1752)	—	—	—	—	—	Q.	N.A.	B.fl.VI,51	M.fr.IV,35.	As., Afr., Am., P.
E. Tannensis; Sprengel, syst. veg. III, 791 (1826)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.	P.
E. eremophila; Cunningham in Mitch. Trop. Austr. 348 (1848)	W.A.	S.A.	V.	—	N.S.W.	Q.	N.A.	B.fl.VI,52		
E. corynoclada; F. v. M. in Wing's S. Sc. Rec., new ser., vol. II, Apr. (1886)	—	—	—	—	—	Q.	—	—		
MONOTAXIS; Brogniart in Duperr. Voy. Bot. 223, t. 49 (1829). (Hippocrepandra.)										
M. macrophylla; Bentham, Fl. Austr. VI, 79 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,79		
M. linifolia; Brongniart in Duperr. Voy. Bot. 223, t. 49 (1829)	—	—	—	—	N.S.W.	—	—	B.fl.VI,79		
M. occidentalis; Endlicher in Hueg. enum. 19 (1837)	W.A.	—	—	—	—	—	—	B.fl.VI,79		
M. lurida; Bentham, Fl. Austr. VI, 80 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,80		
M. megacarpa; F. v. M., fragm. IV, 143 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,80	M.fr.IV,143.	
M. gracilis; Baillon, Adans. VI, 293 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,80		
M. grandiflora; Endlicher in Hueg. enum. 19 (1837)	W.A.	—	—	—	—	—	—	B.fl.VI,81	M.fr.X,53.	
M. luteiflora; F. v. M., fragm. X, 51 (1877)	W.A.	—	—	—	—	—	—	—	M.fr.X,51.	
PORANTHERA; Rudge in Transact. Linn. Soc. X, 302 t. 22 (1811).										
P. ericifolia; Rudge in Transact. Linn. Soc. X, 302, t. 22 (1811)	—	—	—	—	N.S.W.	—	—	B.fl.VI,55		
P. ericoides; Klotzsch in Lehm. pl. Preiss. II, 239 (1847)	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,55		
P. Huegelii; Klotzsch in Lehm. pl. Preiss. II, 231 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,55		
P. corymbosa; Brongniart in Duperr. Voy. Coq. Bot. 219, t. 56 (1826)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,56	M.fr.VI,183.	
P. microphylla; Brongniart in Duperr. Voy. Coq. Bot. 218, t. 50 (1826)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,56	M.fr.XI,20; XII,11	
MICRANTHEUM; Desfontaines in Mém. du. Mus. hist. nat. IV, 253 (1818). (Caletia.)										
M. ericoides; Desfontaines in Mém. du Mus. IV, 253, t. 14 (1818)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,57	M.fr.I,32.	
M. hexandrum; J. Hooker in Lond. Journ. VI, 283 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,57	M.fr.XII,11.	
PSEUDANTHUS; Sieber in Sprengel, syst. veg. cur. post. 25 (1827). (Stachystemon, Chrysostemon, Choirizotheca.)										
P. pimeloides; Sieber in Sprengel, syst. veg. cur. post. 25 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,59	M.fr.I,107.	
P. ovalifolius; F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,59	M.fr.I,107.	
P. micranthus; Bentham, Fl. Austr. VI, 59 (1873)	—	S.A.	—	—	—	—	—	B.fl.VI,59	M.fr.XII,11.	
P. divaricatissimus; Bentham, Fl. Austr. VI, 60 (1873)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,60		
P. orientalis; F. v. M., fragm. II, 14 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,60	M.fr.II,14.	
P. occidentalis; F. v. M., fragm. I, 107 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,60	M.fr.I,107; IV,35.	
P. nematophorus; F. v. M., fragm. II, 14 (1860)	W.A.	—	—	—	—	—	—	B.fl.VI,61	M.II,14.	
P. polyandrus; F. v. M., fragm. II, 153 (1861)	W.A.	—	—	—	—	—	—	B.fl.VI,62	M.fr.II,153; IV,35,173.	
P. brachyphyllus; F. v. M. in Transact. R. Soc. N.S.W. 11 (1881)	W.A.	—	—	—	—	—	—	B.fl.VI,62		
P. vermicularis; F. v. M. in Transact. R. Soc. N.S.W. 11 (1881)	W.A.	—	—	—	—	—	—	B.fl.VI,62		
BEYERIA; Miquel in Ann. sc. nat. sér. trois. I, 350 (1844). (Calyptrostigma, Beyeriopsis.)										
B. viscosa; Miquel in Ann. des sc. nat. sér. trois. I, 350, t. 15 (1844)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,64	M.fr.I,230; II,181; XII,10.	
B. lasiocarpa; F. v. M. in Benth. Fl. Austr. VI, 65 (1873)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,65	M.fr.VI,182.	
B. opaca; F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,65	M.fr.VI,182; X,53.	
B. uncinata; F. v. M. in Benth. Fl. Austr. VI, 65 (1873)	—	S.A.	—	—	—	—	—	B.fl.VI,65		
B. latifolia; Baillon, Adans. VI, 304 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,66		
B. cygnorum; Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,66		
B. cinerea; Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,66		
B. cyanescens; Bentham, Fl. Austr. VI, 66 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,66		
B. lepidopetala; F. v. M., fragm. I, 230 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,67	M.fr.I,230; III,164.	
B. similis; Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,67		
B. brevifolia; Baillon, Adans. VI, 309 (1866)	W.A.	—	—	—	—	—	—	B.fl.VI,67		
B. Drummondii; J. Mueller in Linnaea XXXIV, 58 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,68		
B. tristigma; F. v. M., fragm. VI, 181 (1868)	—	—	—	—	—	Q.	—	B.fl.VI,68	M.fr.VI,181.	

RICINOCARPUS; Desfontaines in Mém. du Mus. III, 459, t. 22 (1817). (Roeperia.)

R. pinifolius; Desfontaines in Mém. du Mus. III, 459, t. 22 (1817)		—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,70	M.fr.VI,182.		
R. tuberculatus; J. Mueller in Linnaea XXXIV, 60 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.VI,70			
R. cyanescens; J. Mueller in Linnaea XXXIV, 60 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.VI,70			
R. psilocladus; Bentham, Fl. Austr. VI, 71 (1873)	W.A.	—	—	—	—	—	—	—	B.fl.VI,70			
R. glaucus; Eudlicher in Hueg. enum. 18 (1873)	W.A.	—	—	—	—	—	—	—	B.fl.VI,71	M.fr.VI,182.		
R. major; J. Mueller in Linnaea XXXIV, 59 (1865)		—	—	T.	—	—	—	—	B.fl.VI,72			
R. Bowmanii; F. v. M., fragm. I, 181 (1859)		—	—	—	—	N.S.W.	Q.	—	B.fl.VI,72	M.fr.III,164; IV,35.		
R. ledifolius; F. v. M., fragm. I, 76 (1858)		—	—	—	—	—	Q.	—	B.fl.VI,72	M.fr.IV,35.		
R. rosmarinifolius; Bentham, Fl. Austr. VI, 72 (1873)		—	—	—	—	—	—	N.A.	B.fl.VI,72			
R. marginatus; Bentham, Fl. Austr. VI, 73 (1873)		—	—	—	—	—	—	N.A.	B.fl.VI,73			
R. speciosus; J. Mueller in De Candolle, prodr. XV, 204 (1863)		—	—	—	—	N.S.W.	—	—	B.fl.VI,73			
R. trichoporus; J. Mueller in Linnaea XXXIV, 60 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.VI,73			
R. muricatus; J. Mueller in Linnaea XXXIV, 61 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.VI,73			
R. velutinus; F. v. M., fragm. IX, 2 (1875)	W.A.	—	—	—	—	—	—	—	—	M.fr.IX,2.		

BERTYA; Planchon in Hook. Lond. Journ. Bot. IV, 472, t. 16 (1845). (Lambertya).

B. gummifera; Planchon in Hook. Lond. Journ. IV, 473 (1845)		—	—	—	—	N.S.W.	—	—	B.fl.VI,75	M.fr.IV,35.		
B. pinifolia; Planchon in Hook. Lond. Journ. IV, 473 (1845)		—	—	—	—	N.S.W.	Q.	—	B.fl.VI,75			
B. quadrisepala; F. v. M., fragm. X, 52 (1876)	W.A.	—	—	—	—	—	—	—	—	M.fr.X,52.		
B. Cunninghamii; Planchon in Hook. Lond. Journ. IV, 473 (1845)		—	—	—	V.	N.S.W.	—	—	B.fl.VI,75	M.fr.IV,35; X,52.		
B. dimerostigma; F. v. M. in Wing's Science Journ. II (1882)	W.A.	—	—	—	—	—	—	—	—	M.fr.XII,9.		
B. rosmarinifolia; Planchon in Hook. Lond. Journ. IV, 473 (1845)		—	—	T.	—	N.S.W.	—	—	B.fl.VI,76	M.fr.IV,35; XII,10.		
B. Mitchelli; J. Mueller in Linnaea XXXIV, 63 (1865)		—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,76			
B. oppositifolia; F. v. M. & O'Shanesy in S. Sc. Rec. II, May (1882)		—	—	—	—	—	Q.	—	—	M.fr.XII,9.		
B. oleaefolia; Planchon in Hook. Lond. Journ. IV, 473 (1845)		—	—	—	—	N.S.W.	Q.	—	B.fl.VI,76			
B. rotundifolia; F. v. M., fragm. IV, 34 (1864)		—	S.A.	—	—	—	—	—	B.fl.VI,77	M.fr.XII,10.		
B. pedicellata; F. v. M., fragm. IV, 143 (1865)		—	—	—	—	—	Q.	—	B.fl.VI,77	M.fr.IV,143.		
B. pomaderrioides; F. v. M., fragm. IV, 34 (1864)		—	—	—	—	N.S.W.	—	—	B.fl.VI,77	M.fr.VIII,141.		
B. Findlayi; F. v. M., fragm. VIII, 141 (1874)		—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,141.		

AMPEREA; Adr. de Jussieu, Euphorb. gen. 35, t. 10 (1824).

A. protensa; Nees in Lehm. pl. Preiss. II, 229 (1847)	W.A.	—	—	—	—	—	—	—	B.fl.VI,82			
A. volubilis; F. v. M. in Benth. Fl. Austr. VI, 82 (1873)	W.A.	—	—	—	—	—	—	—	B.fl.VI,82	M.fr.VI,182.		
A. micrantha; Bentham, Fl. Austr. VI, 83 (1873)	W.A.	—	—	—	—	—	—	—	B.fl.VI,83			
A. conferta; Bentham, Fl. Austr. VI, 83 (1873)	W.A.	—	—	—	—	—	—	—	B.fl.VI,83			
A. ericoides; Adr. de Jussieu, Tent. Euphorb. 112, t. 10 (1824)	W.A.	—	—	—	—	—	—	—	B.fl.VI,83			
A. spartioides; Brongniart in Duperr. Voy. Coq. 221, t. 49 (1826)		—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,84			

ANTIDESMA; Linné, Fl. Zeyl. 169 (1747).

A. Ghaesembilla; Gaertner, de Fruct. I, 189, t. 39 (1788)		—	—	—	—	—	—	N.A.	B.fl.VI,85	M.fr.IV,86; VI,183.	...	As., P.
A. Dallachyanum; Baillon, Adans. VI, 337 (1866)		—	—	—	—	—	Q.	N.A.	B.fl.VI,85	M.fr.VI,183.		
A. Bunius; Sprengel, syst. veg. I, 826 (1825)		—	—	—	—	—	Q.	—	B.fl.VI,86		...	As., P.
A. parvifolium; Thwaites & F. v. M., fragm. IV, 86 (1864)		—	—	—	—	—	Q.	—	B.fl.VI,86	M.fr.VI,183.		
A. Schultzii; Bentham, Fl. Austr. VI, 86 (1873)		—	—	—	—	—	—	N.A.	B.fl.VI,86			
A. erostre; F. v. M. in Benth. Fl. Austr. VI, 87 (1873)		—	—	—	—	—	Q.	—	B.fl.VI,87			
A. sinuatum; Bentham, Fl. Aust. VI, 87 (1873)		—	—	—	—	—	Q.	—	B.fl.VI,87			

ANDRACHNE; Linné, syst. nat. 9 (1735); gen. pl. 287 (1737).

A. Decaisnei; Bentham, Fl. Austr. VI, 88 (1873)	W.A.	—	—	—	—	—	Q.	N.A.	B.fl.VI,88	M.fr.VI,183	...	As.

ACTEPHILA; Blume, Bijdr. XII, 581 (1825).

A. grandifolia; Baillon, Adans. VI, 330 & 360, t. 10 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,89				
A. Mooreana; Baillon, Adans. VI, 330 & 366 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,89				
A. latifolia; Bentham, Fl. Austr. VI, 89 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,89				
A. petiolaris; Bentham, Fl. Austr. VI, 89 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,89				
A. sessilifolia; Bentham, Fl. Austr. VI, 90 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,90				

DISSILIARIA; F. v. M. in Baillon, Adans. VII, 356, pl. 1 (1867). (Choriceras.)

D. baloghioides; F. v. M. in Baillon, Adans. VII, 356 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,90				
D. Muelleri; Baillon, Adans. VII, 359, t. 1 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,91				
D. tricornis; Bentham, Fl. Austr. VI, 91 (1873)	—	—	—	—	—	Q.	N.A.	B.fl.VI,91	M.fr.XII,11.			

PETALOSTIGMA; F. v. M. in Hook. Kew Misc. IX, 17 (1857). (Hylococcus.)

P. quadriloculare; F. v. M. in Hook. Kew Misc. IX, 17 (1857)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,92	M.fr.VI,182.			

PHYLLANTHUS; Linné, gen. pl. 282 (1737), from J. Commelin. (Glochidion, Bradleia, Synostemon.)

P. Ferdinandi; J. Mueller in Regensb. Flora, 379 (1865)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,96	M.fr.VI,183.			
P. lobocarpus; Bentham, Fl. Austr. VI, 97 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,97				
P. ditassoides; J. Mueller in Regensb. Flora, 487 (1864)	—	—	—	—	—	—	N.A.	B.fl.VI,97				
P. Adami; J. Mueller in De Candolle, prodr. XV, 327 (1866)	—	—	—	—	—	—	N.A.	B.fl.VI,97	M.fr.I,33.			
P. thesioides; Bentham, Fl. Austr. VI, 98 (1873)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,98				
P. hirtellus; J. Mueller in De Candolle, prodr. XV, 326 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,98	M.fr.III,89.			
P. rigens; J. Mueller in Regensb. Flora, 513 (1864)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VI,99	M.fr.II,153; XII,11.			
P. ochrophyllus; Bentham, Fl. Austr. VI, 99 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,99				
P. rigidulus; F. v. M. in Linnaea XXXIV, 72 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,99				
P. ramosissimus; J. Mueller in Linnaea XXXIV, 70 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,100	M.fr.I,33; III,163.			
P. Tatei; F. v. M. in Wing's South. Science Record II, 55 (1882)	—	S.A.	—	—	N.S.W.	—	—	—	M.fr.XII,10.			
P. rhytidospermus; F. v. M. in Linnaea XXXIV, 70 (1865)	—	S.A.	—	—	—	—	N.A.	B.fl.VI,100	M.fr,XII,11.			
P. albiflorus; F. v. M. in Linnaea XXXIV, 70 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,100				
P. crassifolius; J. Mueller in Regensb. Flora, 513 (1864)	W.A.	—	—	—	—	—	—	B.fl.VI,100				
P. elachophyllus; F. v. M. in Benth. Fl. Austr. VI, 101 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,101				
P. uberiflorus; F. v. M. in Adansonia, 343 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,101				
P. reticulatus; Poiret, encycl. meth. V, 298 (1804)	—	—	—	—	—	—	N.A.	B.fl.VI,101	...	...	As.	
P. baccatus; F. v. M. in Benth. Fl. Austr. VI, 102 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,102				
P. Urinaria; Linné, sp. pl. 982 (1753)	—	—	—	—	—	—	N.A.	B.fl.VI,102	...	...	As., Afr.	
P. trachygyne; Bentham, Fl. Austr. VI, 103 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,103				
P. Maderaspatanus; Linné, sp. pl. 982 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VI,103				
P. Mitchelli; Bentham, Fl. Austr. VI, 103 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,103	...	...	As., Afr., P.	
P. Gastroemii; J. Mueller in De Cand., prodr. XV, 358 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,104	M.fr.XII,11.			
P. Dallachyanus; Bentham, Fl. Austr. VI, 104 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,104				
P. subcrenulatus; F. v. M., fragm. I, 108 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,104	M.fr.I,108.			
P. calycinus; Labillardière, Nov. Holl. pl. spec. II, 75, t. 225 (1806)	W.A.	S.A.	—	—	—	—	—	B.fl.VI,105				
P. flagellaris; Bentham, Fl. Austr. VI, 106 (1873)	—	—	—	—	—	—	N.A.	B.fl.VI,106				
P. similis; J. Mueller in Linnaea XXXIV, 71 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,106				
P. microcladus; J. Mueller in Linnaea XXXIV, 71 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,106				
P. grandisepalus; F. v. M. in Linnaea XXXIV, 72 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,106				
P. Carpentariae; J. Mueller in Linnaea XXXIV, 72 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,107				
P. Fuernrohrii; F. v. M. in Transact. phil. Soc. Vict. I, 15 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,107				
P. hebecarpus; Bentham, Fl. Austr. VI, 108 (1873)	—	—	—	—	—	Q.	N.A.	B.fl.VI,108				

P. lacunarius ; F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854)... ... ... — S.A. — V. N.S.W. Q. — B.fl. VI,108
P. trachyspermus ; F. v. M. in Transact. phil. Soc. Vict. I, 14 (1854) ... ... — S.A. — V. N.S.W. Q. — B.fl. VI,108
P. Australis ; J. Hooker in Lond. Journ. VI, 284 (1847) ... ... — S.A. T. — N.S.W. — — B.fl. VI,108 M.fr. XII,11.
P. thymoides ; Sieber in Linnaea XXVIII, 566 (1856) ... ... — S.A. — V. N.S.W. Q. — B.fl. VI,109
P. scaber ; Klotzsch in Lehm. pl. Preiss. I, 179 (1844) ... ... W.A. — — — — — — B.fl. VI,109
P. indigoferoidea ; Bentham, Fl. Austr. VI, 110 (1873) ... ... — — — — — — N.A. B.fl. VI,110
P. aridus ; Bentham, Fl. Austr. VI, 110 (1873) ... ... — — — — — — N.A. B.fl. VI,110
P. Gunnii ; J. Hooker in Lond. Journ. VI, 284 (1847) ... ... — S.A. T. V. N.S.W. — — B.fl. VI,110
P. simplex ; Retzius, observ. V, 29 (1789) ... ... — — — — — Q. — B.fl. VI,111 M.fr. VI,183. ... As., P.
P. filicaulis ; Bentham, Fl. Austr. VI, 111 (1873) ... ... — — — — N.S.W. — — B.fl. VI,111
P. minutiflorus ; F. v. M. in Linnaea XXXIV, 75 (1865) ... . — — — — — Q. N.A. B.fl. VI,112
P. Armstrongii ; Bentham, Fl. Austr. VI, 112 (1873) ... ... — — — — — — N.A. B.fl. VI,112
P. buxifolius ; Reinwardt in Blume, catal. plant. Buitenz. 106 (1823) ... ... — — — — — Q. — — M.fr. X,121. ... As.

BREYNIA ; R. & G. Forster, char. gen. 145, t. 73 (1776). (Melanthesa, Melanthesiopsis.)

B. cernua ; J. Mueller in De Candolle, prodr. XV, 439 (1866) ... ... — — — — — Q. N.A. B.fl. VI,113
B. oblongifolia ; J. Mueller in De Candolle, prodr. XV, 110 (1866) .. ... — — — — N.S.W. Q. — B.fl. VI,114
B. stipitata ; J. Mueller in De Candolle, prodr. XV, 442 (1866) .. .. — — — — — Q. N.A. B.fl. VI,114
B. rhynchocarpa ; Bentham, Fl. Austr. VI, 114 (1873) ... ... — — — — — — N.A. B.fl. VI,114

SECURINEGA ; A. L. de Jussieu, Gen. 388 (1789). (Flueggea.)

S. Abyssinica ; A. Richard, tentam. Fl. Abyss. II, 256 (1851) ... ... W.A. — — — — Q. N.A. B.fl. VI,116 ... ... As., Afr., P.
S. Leucopyrus ; J. Mueller in De Candolle, prodr. XV, 451 (1866) ... ... — S.A. — — — Q. N.A. B.fl. VI,116 ... ... As.

NEOROEPERA ; F. & J. Mueller in De Candolle, prodr. XV, 489 (1866).

N. buxifolia ; F. & J. Mueller in De Candolle, prodr. XV, 489 (1866) ... ... — — — — — Q. — B.fl. VI,116
N. Banksii ; Bentham, Fl. Austr. VI, 117 (1873) ... ... — — — — — Q. — B.fl. VI,117

BISCHOFFIA ; Blume, Bijdr. 1168 (1825).

B. Javanica ; Blume, Bijdr. 1168 (1825) ... ... — — — — — Q. N.A. — M.fr. VIII,141 ... As., P.

HEMICYCLIA ; Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833).

H. sepiaria ; Wight & Arnott in Edinb. phil. Journ. XIV, 297 (1833) ... .. — — — — — — N.A. B.fl. VI,117 M.fr. IV,119 ; VI,182 ; IX,77. As.
H. Australasica ; J. Mueller in De Candolle, prodr. XV, 487 (1866) ... ... — — — — N.S.W. Q. — B.fl. VI,118
H. lasiogyna ; F. v. M., fragm. IV, 119 (1864) ... ... — — — — — — N.A. B.fl. VI,118 M.fr. IV,119.

APOROSA ; Blume, Bijdrag. 514 (1824).

A. Australiana ; F. v. M., syst. cons. of Austr. pl. 20 (1882) ... ... — — — — — Q. — — M.fr. VIII,141.

BRIDELIA ; Willdenow, Spec. pl. IV, 978 (1805).

B. exaltata ; F. v. M., fragm. III, 32 (1862) ... ... — — — — N.S.W. Q. — B.fl. VI,119 M.fr. III,32.
B. ovata ; Decaisne in nouv. ann. du Mus. V, 484 (1835) ... ... — — — — — — N.A. B.fl. VI,120 ... ... As.
B. tomentosa ; Blume, Bijdrag. 597 (1825) . ... — — — — — Q. N.A. B.fl. VI,120 M.fr. III,32 ; VI,182 As., P.
B. faginea ; F. v. M. in Benth. Fl. Austr. VI, 120 (1873) ... ... — — — — — Q. — B.fl. VI,120

CLEISTANTHUS ; J. Hooker, icon. pl. t. 779 (1847). (Lebediera, Amanoa.)

C. Cunninghamii ; J. Mueller in De Candolle, prodr. XV, 506 (1866) ... ... — — — — N.S.W. Q. — B.fl. VI,122
C. apodus ; Bentham, Fl. Austr. VI, 122 (1873) ... ... — — — — — Q. — B.fl. VI,122
C. Dallachyanus ; Baillon in Benth. Fl. Austr. VI, 122 (1873) ... ... — — — — — Q. — B.fl. VI,122 M.fr. VI,182.
C. semiopacus ; F. v. M. in Benth. Fl. Austr. VI, 123 (1873) ... ... — — — — — Q. — B.fl. VI,123

CROTON ; Linné, gen. pl. 288 (1737).

C. Schultzii ; Bentham, Fl. Austr. VI, 124 (1873) ... ... — — — — — — N.A. B.fl.VI,124
C. insularis ; Baillon, Adans. II, 217 (1862) ... ... — — — — N.S.W. Q. — B.fl.VI,124
C. phebalioides ; F. v. M. in Benth. Fl. Austr. VI, 125 (1873) ... ... — — — — N.S.W. Q. — B.fl.VI,125 M.fr.IV,140 ... P.
C. opponens ; F. v. M. in Benth. Fl. Austr. VI, 125 (1873) ... ... — — — — — Q. — B.fl.VI,125
C. tomentellus ; F. v. M., fragm. IV, 141 (1864) ... ... — — — — — — N.A. B.fl.VI,126 M.fr.IV,141.
C. Verreauxii ; Baillon, Etud. Euph. 357 (1858) ... ... — — — — N.S.W. Q. N.A. B.fl.VI,126 M.fr.IV,141 ; VI,186.
C. acronychioides ; F. v. M., fragm. IV, 142 (1864) ... ... — — — — — Q. — B.fl.VI,127 M.fr.VI,186.
C. triacros ; F. v. M., fragm. VI, 185 (1867) ... ... — — — — — Q. — B.fl.VI,127 M.fr.VI,185.
C. Arnhemicus ; F. v. M. in Linnaea XXXIV, 112 (1865) ... ... — — — — — Q. N.A. B.fl.VI,127 M.fr.VI,186.

ALEURITES ; R. & G. Forster, char. gen. 111, t. 56 (1776).

A. triloba ; R. & G. Forster, char. gen. 111, t. 56 (1776) ... ... — — — — — Q. — B.fl.VI,128 M.fr.V,30 ; VI,182. As., P.

CLAOXYLON ; Adr. de Jussieu, Euphorb. gen. 43, t. 14 (1824).

C. angustifolium ; J. Mueller in Linnaea XXXIV, 165 (1865) ... ... — — — — — Q. — B.fl.VI,129
C. tenerifolium ; F. v. M. in Baillon, Adans. VI, 323 (1866) ... ... — — — — — Q. — B.fl.VI,130 M.fr.VI,183.
C. Australe ; Baillon, Etud. Euphorb. 493 (1858) ... ... — — — V. N.S.W. Q. — B.fl.VI,130 M.fr.IV,142.
C. Hillii ; Bentham, Fl. Austr. VI, 131 (1873) ... ... — — — — — Q. — B.fl.VI,131

ACALYPHA ; Royen in Linné, hort. Cliffort. 495 (1737).

A. nemorum ; F. v. M. in Linnaea XXXIV, 38 (1865) ... ... — — — — N.S.W. Q. — B.fl.VI,132
A. eremorum ; F. v. M. in Regensb. Flora 440 (1864) ... ... — — — — — Q. — B.fl.VI,132
A. capillipes ; F. v. M. in Linnaea XXXIV, 40 (1865) ... . — — — — N.S.W. Q. — B.fl.VI,133

ADRIANA ; Gaudichaud in Ann. sc. nat. V. 223 (1825). (Trachycaryon.)

A. tomentosa ; Gaudichaud in Ann. sc. nat. V, 223 (1825) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VI,134 M.fr.IX,2 ; XII,11.
A. quadripartita ; Gaudichaud in Freyc. voy. Bot. 489 (1826) ... ... W.A. S.A. — V. — — — B.fl.VI,135 M.fr.XII,11.

ALCHORNEA ; Solander in Swartz, prodr. 698 (1788). (Coelebogyne, Cladodes.)

A. ilicifolia ; J. Mueller in Linnaea XXXIV, 170 (1865) ... ... — — — — N.S.W. Q. — B.fl.VI,136
A. Thozetiana ; F. v. M. in Benth. Fl. Austr. VI, 137 (1873) ... ... — — — — — Q. — B.fl.VI,137

TRAGIA ; Plumier, gen. 14, t. 12 (1703).

T. Novae Hollandiae ; J. Mueller in Linnaea XXXIV, 180 (1865) ... ... — — — — N.S.W. Q. — B.fl.VI,137 M.fr.VI,183 ; XII,11.

MALLOTUS ; Loureiro, Fl. Cochinch. II, 635 (1790). (Echinus, Rottlera, Echinocroton.)

M. ricinoides ; J. Mueller in Linnaea XXXIV, 187 (1865) ... ... — — — — — Q. — B.fl.VI,140 M.fr.IV,138 ; VI,184 As., P.
M. Chinensis ; Loureiro, Fl. Cochinchin. II, 635 (1770) ... ... — — — — — Q. — B.fl.VI,140 M.fr.VI,184 ... As., P.
M. claoxyloides ; J. Mueller in Linnaea XXXIV, 192 (1865) ... . — — — — N.S.W. Q. — B.fl.VI,140 M.fr.I,32 ; VI,184.
M. Philippinensis ; J. Mueller in Linnaea XXXIV, 196 (1865) ... ... — — — — N.S.W. Q. — B.fl.VI,141 M.fr.VI,183 ... As., P.
M. angustifolius ; Bentham in Fl. Austr. VI, 141 (1873) ... ... — — — — — Q. — B.fl.VI,142
M. polyadenos ; F. v. M., fragm. VI, 184 (1867) ... ... — — — — — Q. — B.fl.VI,142 M.fr.VI,184.
M. repandus ; J. Mueller in Linnaea XXXIV, 197 (1865) ... ... — — — — — Q. — B.fl.VI,142 M.fr.VI,185.
M. nesophilus ; F. v. M. in Linnaea XXXIV, 196 (1865) ... ... — — — — — Q. N.A. B.fl.VI,143 M.fr.VI,184.
M. discolor ; F. v. M. in Benth. Fl. Austr. VI, 143 (1873) ... ... — — — — N.S.W. Q. — B.fl.VI,143 M.fr.IV,139.

MACARANGA ; Petit-Thouars, Gen. Madag. n. 88 (1809). (Mappa.)

M. Dallachyi ; F. v. M. in Benth. Fl. Austr. VI, 144 (1873) ... ... — — — — — Q. — B.fl.VI,144 M.fr.VI,185.
M. subdentata ; Bentham, Fl. Austr. VI, 145 (1873) ... ... — — — — — Q. — B.fl.VI,145
M. inamoena ; F. v. M. in Benth. Fl. Austr. VI, 145 (1873) ... . — — — — — Q. — B.fl.VI,145
M. involucrata ; Baillon, Etud. Euphorb. 432 (1858) ... ... — — — — — Q. — B.fl.VI,146 M.fr.IV,139,177 ; VI,183 As.
M. Tanaria ; J. Mueller in De Candolle, prodr. XV, 997 (1866) ... ... — — — — N.S.W. Q. N.A. B.fl.VI,146 M.fr.IV,140 ; VI,183 ... As., P.

CODIAEUM ; Rumphius, herbar. Amboin. IV, 65, t. 25 (1743).

C. variegatum ; Blume, Bijdrag. 605 (1825) ... ... — — — — — Q. — B.fl.VI,147 M.fr.VI,182 ... ... As., P.

BALOGHIA ; Endlicher, prodr. fl. Norfolk, 84 (1833).

B. lucida ; Endlicher, prodr. fl. Norfolk, 84 (1833) ... ... — — — — N.S.W. Q. — B.fl.VI,148 M.fr.VI,182 ; IX,77 .. P.

FONTAINEA ; Heckel, thése inaugurale, Montpell. (1870).

F. Pancheri ; Heckel, thése inaugurale (1870) ... ... — — — — — Q. — B.fl.VI,149 M.fr.XII,11 ... ... P.

OMALANTHUS ; Adr. de Jussieu, Euphorb. gen. 50, t. 16 (1824). (Carumbium, Wartmannia, Homalanthus.)

O. Leschenaultianus ; A. de Jussieu, Euph. gen. 50, t. 16, f. 53 (1824) ... ... — — — V. N.S.W. Q. N.A. B.fl.VI,150 M.fr.I,32 ; VI,183 ; IX,77 ; XII,11 As., P.
O. stillingiifolius ; F. v. M., fragm. I, 32 (1858) ... ... — — — — N.S.W. Q. — B.fl.VI,150 M.fr.I,33 ; III,163

SEBASTIANIA ; Sprengel, Neue Entd. II, 118 (1821). (Microstachys, Elachocroton.)

S. chamelaea ; J. Mueller in De Candolle, prodr. XV, 1175 (1866) ... ... — — — — — Q. N.A. B.fl.VI,151 ... ... As., Afr., P.

EXCAECARIA ; Linné, syst. veg. ed. decim. 1288 (1759).

E. Agallocha ; Linné, syst. ed. decim. 1288 (1759) ... ... — — — — N.S.W. Q. N.A. B.fl.VI,152 M.fr.XI,64 ... As., P.
E. Dallachyana ; Baillon, Adans. VI, 324 (1866) ... ... — — — — — Q. — B.fl.VI,153
E. parvifolia ; J. Mueller in Flora, 433 (1864) ... ... — — — — — Q. N.A. B.fl.VI,153

URTICACEAE.

VENTENAT ; Tabl. du règn. vég. III, 524 (1799).—Genera 19, Species 65.

CELTIS ; Tournefort, inst. 612, t. 383 (1700) from Camerarius (1586). (Solenostigma.)

C. Philippinensis ; Blanco, Fl. de Filipinas 197 (1837) ... ... — — — — — Q. N.A. B.fl.VI,156 M.fr.IV,88 ... As., P.
C. paniculata ; Planchon in Ann. sc. nat. sér. trois. X, 305 (1848) ... ... — — — — N.S.W. Q. N.A. B.fl.VI,156 M.fr.IV,88 ... As.
C. amblyphylla ; F. v. M., fragm. IX, 76 (1875) ... ... — — — — N.S.W. — — — M.fr.IX,76.

TREMA ; Loureiro, Fl. Cochinch. II, 565 (1790). (Sponia.)

T. cannabina ; Loureiro, Fl. Cochinch. II, 689 (1790) ... ... — — — V. N.S.W. Q. N.A. B.fl.VI,158 M.fr.VI,194 ; VIII,247 As., P.

APHANANTHE ; Planchon in Ann. sc. nat. X, 265 (1848).

A. Philippinensis ; Planchon in Ann. sc. nat. X, 337 (1848) ... ... — — — — N.S.W. Q. — B.fl.VI,160 M.fr.VI,192 ... As.

FICUS ; Tournefort, inst. 662 (1700) from Plinius. (Urostigma, Covellia.)

F. colossea ; F. v. M. in Benth. Fl. Austr. VI, 163 (1873) ... ... — — — — — Q. — B.fl.VI,163
F. pilosa ; Reinwardt in Blume, Bijdr. 446 (1825) ... ... — — — — — Q. — B.fl.VI,164 M.fr.VI,195 ... As.
F. glabella ; Blume, Bijdr. 452 (1825) ... ... — — — — — Q. N.A. B.fl.VI,164 M.fr.VI,195 ... As.
F. Cunninghamii ; Miquel in Ann. Mus. Lugd. Bat. III. 286 (1867) ... ... — — — — — Q. — B.fl.VI,165 M.fr.VI,195 ; VIII,246.
F. Henneana ; Miquel in Ann. Mus. Lugd. Bat. III, 216 (1867) ... ... — — — — — Q. N.A. B.fl.VI,165
F. validinervis ; F. v. M. in Benth. Fl. Austr. VI, 166 (1873) ... ... — — — — — Q. — B.fl.VI,166
F. retusa ; Linné, mantissa plant. 129 (1767) ... ... — — — — — Q. N.A. B.fl.VI,166 M.fr.VI,195 ... As., P.
F. eugenioides ; F. v. M., Docum. Interc. Exhib. 26 (1866) ... ... — — — — N.S.W. Q. N.A. B.fl.VI,166 M.fr.VI,195
F. Benjaminea ; Linné, mantissa plant. 129 (1767) ... ... — — — — — Q. — B.fl.VI,167 M.fr.VI,195. ... As.
F. Muelleri ; Miquel in Ann. Mus. Lugd. Bat. III, 287 (1867) ... ... — — — — N.S.W. — — B.fl.VI,167 M.fr.VI,195 ; VIII,246.
F. leucotricha ; Miquel in Ann. Mus. Lugd. Bat. III, 285 (1867) .. .. — — — — — Q. N.A. B.fl.VI,167 M.fr.VI,195.
F. rubiginosa ; Desfontaines, catal. pl. hort. Paris. 209 (1804) ... ... — — — — N.S.W. Q. — B.fl.VI,168 M.fr.IV,194 ; VIII,247.
F. columnaris ; F. v. M. & Moore in proc. Accl. Soc. Vict. III, 71 (1874) ... — — — — N.S.W. — — — M.fr.VIII,247.
F. macrophylla ; Desfontaines, catal. pl. hort. Paris. 209 (1804) ... ... — — — — N.S.W. Q. — B.fl.VI,170 M.fr.VII,194 ; VIII,247.
F. pubcrula ; Cunningham in Hook. Lond. Journ. VI, 562 (1847) ... ... — — — — — — N.A. B.fl.VI,169

F. platypoda; Cunningham in Hook. Lond. Journ. VI, 561 (1847) ... ... W.A. S.A. — — — Q. N.A. B.fl.VI,169 M.fr.VI,194; VIII,247.
F. dictyophleba; F. v. M. in Benth. Fl. Austr. VI, 170 (1873) ... ... — — — — — Q. — B.fl.VI,170
F. magnifolia; F. v. M., fragm. IV, 50 (1864) ... ... — — — — — Q. — B.fl.VI,171 M.fr.IV,177; VI,196.
F. ehretioides; F. v. M. in Benth. Fl. Austr. VI, 171 (1873) ... ... — — — — — Q. — B.fl.VI,171
F. pumila; Linné, spec. plant. 1060 (1753) ... ... — — — — N.S.W. Q. — B.fl.VI,171 M.fr.VI,196 ... As.
F. coronulata; F. v. M. in Journ. Bot. Neerland. 241 (1861) ... ... — — — — — — N.A. B.fl.VI,172 M.fr.IV,49; VI,196; VIII,246.
F. leptoclada; Bentham, Fl. Austr. VI, 172 (1873) ... ... — — — — — Q. — B.fl.VI,172
F. depressa; Bentham, Fl. Austr. VI, 172 (1873) ... ... — — — — — Q. — B.fl.VI,172
F. subulata; Blume Bijdr., 461 (1825) ... ... — — — — Q. — — ... ... As.
F. infectoria; Roxburgh. flor. Indic. III, 550 (1832) ... ... — — — — — Q. — — ... ... As.
F. Philippinensis; Miquel in Hooker, Lond. Jonrn. VII, 435 (1848) ... ... — — — — — Q. — B.fl.VI,173 ... ... As., P.
F. mollior; F. v. M. in Benth. Fl. Austr. VI, 173 (1873) ... ... — — — — — Q. — B.fl.VI,173
F. stenocarpa; F. v. M. in Benth. Fl. Austr. VI, 174 (1873) ... ... — — — — N.S.W. Q. — B.fl.VI,174 M.fr.IX,152.
F. scabra; G. Forster, florul. ins. Austr. prodr. 76 (1786) ... ... — — — V. N.S.W. Q. N.A. B.fl.VI,174 M.fr.IV,85; VI,196; IX,152; X,114. As., P.
F. subglabra; F. v. M., fragm. IX, 152 (1875) ... ... — — — — — Q. — —
F. Pinkiana; F. v. M. in Wing's S. Sec. Rec. II, 273 (1882) ... ... — — — — — Q. — —
F. orbicularis; Cunningham in Hook. Lond. Journ. VII, 426 (1848) ... ... — S.A. — — — — N.A. B.fl.VI,175 M.fr.VI,196.
F. aculeata; Cunningham in Hook. Lond. Journ. VII, 426 (1848) ... ... — — — — — — N.A. B.fl.VI,175
F. opposita; Miquel in Hook. Lond. Journ. VII, 426 (1848) ... ... — — — — N.S.W. Q. — B.fl.VI,176 M.fr.VI,196; VIII,246.
F. scobina; Bentham, Fl. Austr. VI, 176 (1873) ... ... — — — — — — N.A. B.fl.VI,176
F. hispida; Linné, fil. suppl. 442 (1781) ... ... — — — — — Q. N.A. B.fl.VI,176 ... ... As.
F. fasciculata; F. v. M. in Benth. Fl. Austr. VI, 177 (1873) ... ... — — — — — Q. — B.fl.VI,177
F. casearia; F. v. M. in Benth. Fl. Austr. VI, 177 (1873) ... ... — — — — — Q. — B.fl.VI,177 ... ... As.
F. glomerata; Willdenow, spec. pl. IV, 1148 (1806) ... ... — — — — — Q. N.A. B.fl.VI,178 M.fr.VI,195 ... As.
F. pleurocarpa; F. v. M., fragm. VIII, 246 (1874) ... ... — — — — — Q. — — M.fr.VIII,246.

CUDRANIA; Trécul in Ann. des sc. nat. VIII, 122 (1847).
C. Javanensis; Trécul in Ann. des sc. nat. VIII, 123 (1847) ... ... — — — — N.S.W. Q. — B.fl.VI,179 M.fr.IV,88 ... As., Afr., P.

ANTIARIS; Leschenault in Ann. du Mus. XVI, 476 (1810).
A. macrophylla; R. Brown in Flind. voy. II, 602, t. 5 (1814) ... ... — — — — — — N.A. B.fl.VI,179 M.fr.VI,194.

MALAISIA; Blanco, Fl. Filip. 789 (1837). (Caturus.) [As., P.
M. tortuosa; Blanco, Fl. Filip. 789 (1837) ... ... — — — — N.S.W. Q. N.A. B.fl.VI,180 M.fr.IV,88; VI,193&255; VIII,247; IX,77.

PSEUDOMORUS; Bureau in Ann. des sc. nat. sér. cinq. XI, 372 (1872).
P. Brunoniana; Bureau in Ann. des sc. nat. sér. cinq. XI, 372 (1872) ... ... — — — — N.S.W. Q. — B.fl.VI,181 M.fr.IV,88; VI,192. P.

FATOUA; Gaudichaud in Freyc. voy. Bot. 509 (1826).
F. pilosa; Gaudichaud in Freyc. voy. Bot. 509 (1826) ... ... — — — — — — N.A. B.fl.VI,182 ... ... As., P.

ELATOSTEMMA; R. & G. Forster, char. gen. 105, t. 53 (1776).
E. reticulatum; Weddell in Ann. des sc. nat. sér. 4, I, 188 (1854) ... ... — — — — N.S.W. Q. — B.fl.VI,183 M.fr.IX,77.
E. stipitatum; Weddell in Ann. des sc. nat. sér. 4, I, 190 (1854) ... ... — — — — N.S.W. Q. — B.fl.VI,184

PROCRIS; Commerçon in A. L. de Jussieu, gen. 403 (1789).
P. montana; Steudel, nomencl. bot. II, 398 (1841) ... ... — — — — N.S.W. — — — M.fr.IX,169.

BOEHMERIA; N. J. Jacquin, stirp. Amer. hist. 246, t. 157 (1763).
B. calophleba; Moore & Mueller, fragm. VIII, 11 (1870) ... ... — — — — N.S.W. — — B.fl.VI,184 M.fr.VIII,11.
B. Australis; Endlicher, prodr. Fl. Norfolk. 38 (1833) ... ... — — — — N.S.W. — — — M.fr.VIII,12; IX,169.

PIPTURUS ; Weddell in Ann. des sc. nat. sér. quatr. I, 196 (1854).

P. propinquus ; Weddell in Archiv. du Mus. 447, t. 15 (1856)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 185	M.fr. IV, 87 ; VI, 197 ; VIII, 12.						As., P.

POUZOLZIA ; Gaudichaud in Freyc. voy. Bot. 503 (1826). (Memorialis, Hyrtanandra.)

P. Indica ; Gaudichaud in Freyc. voy. Bot. 503 (1826)	—	—	—	—	—	Q.	N.A.	B.fl. VI, 186	M.fr. IV, 87					...	As.
P. quinquenervis ; Bennett, Pl. Jav. rar. 66 (1852)	—	—	—	—	—	Q.	N.A.	B.fl. VI, 187	M.fr. V, 194					...	As.

PARIETARIA ; Tournefort, inst. 509, t. 289 (1700) from C. Bauhin (1623). Freirea, from Brunfels (1532). [N.Z.

P. debilis ; G. Forster, florul. ins. Austr. prodr. 73 (1786)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl. VI, 188	M.fr. IV, 88 ; IX, 77 ; XI, 20.						As., Afr., Am., P.,

AUSTRALINA ; Gaudichaud in Freyc. voy. Bot. 497 (1826).

A. pusilla ; Gaudichaud in Freyc. voy. Bot. 505 (1826)	—	—	T.	V.	N.S.W.	Q.	—	B.fl. VI, 189	M.fr. IV, 88 ; VIII, 247					...	N.Z.

FLEURYA ; Gaudichaud in Freyc. voy. Bot. 497 (1826).

F. interrupta ; Gaudichaud in Freyc. voy. Bot. 497 (1826)...	—	—	—	—	—	Q.	—	—	M.fr. X, 114					...	As., Afr., P.

URTICA ; Tournefort, inst. 534, t. 308 (1700) from Plinius.

U. incisa ; Poiret, Encycl. meth. suppl. IV, 223 (1817)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 190	M.fr. IV, 87					...	N.Z.

LAPORTEA ; Gaudichaud in Freyc. voy. Bot. 498 (1826).

L. gigas ; Weddell in Arch. du Mus. IX, 129, t. 3 & 4 (1856)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 192	M.fr. IV, 87.						
L. photiniphylla ; Weddell in Arch. du Mus. IX, 138 (1856)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 192	M.fr. IV, 88, 194.						
L. moroides ; Weddell in Arch. du Mus. IX, 142 (1856)	—	—	—	—	—	Q.	—	B.fl. VI, 192	M.fr. VIII, 248.						

CUPULIFERAE.

L. Cl. Richard ; Anal. de fr. 32 & 92 (1808).—Genera 2, Species 4.

FAGUS ; Tournefort, inst. 584, t. 351 (1700) from Camerarius (1586).

F. Gunnii ; J. Hooker, icon. plant., t. 881 (1852)	—	—	T.	—	—	—	—	B.fl. VI, 210	M.fr. V, 110.						
F. Cunninghamii ; Hooker, Journ. of Bot. II, 152, t. 7 (1840)	—	—	T.	V.	—	—	—	B.fl. VI, 210	M.fr. V, 110.						
F. Moorei ; F. v. M., fragm. V, 109 (1866)	—	—	—	—	N.S.W.	—	—	B.fl. VI, 211	M.fr. V, 109.						

BALANOPS ; Baillon, Hist. des pl. VI, 237 & 258 (1876).

B. Australiana ; F. v. M., fragm. X, 114 (1877)	—	—	—	—	—	Q.	—	—	M.fr. X, 114.						

CASUARINEAE.

Mirbel in Ann. du Mus. XVI, 451 (1810).—Genus 1, Species 24.

CASUARINA ; Rumphius, herb. Amboin. III, 87, t. 58 (1743).

C. quadrivalvis ; Labillardière, nov. Holl. pl. spec. II, 67, t. 218 (1806) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VI, 195	M.fr. VI, 18&19 ; X, 62.						
C. trichodon ; Miquel in Lehm. pl. Preiss. I, 641 (1845)	W.A.	—	—	—	—	—	—	B.fl. VI, 196	M.fr. VI, 17.						
C. lepidophloia ; F. v. M., fragm. X, 115 (1877)	—	S.A.	—	—	N.S.W.	—	—	—	M.fr. X, 115.						
C. glauca ; Sieber in Sprengel, syst. III, 803 (1826)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VI, 196	M.fr. VI, 18.						
C. Huegeliana ; Miquel in Lehm. pl. Preiss. I, 640 (1845)	W.A.	—	—	—	—	—	—	B.fl. VI, 196	M.fr. VI, 19 ; X, 62.						
C. equisetifolia ; R. & G. Forster, char. gen. 103, t. 52 (1776)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VI, 197	M.fr. VI, 17					...	As., Afr., P.
C. suberosa ; Otto & Dietrich, allgem. Gartenzeit. 155 (1841)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VI, 197	M.fr. VI, 20&254.						
C. Cunninghamiana ; Miquel, Revis. Casuarin. 56, t. 6 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI, 198							
C. paludosa ; Sieber in Spreng. syst. veg. III, 803 (1826)	—	—	—	V.	N.S.W.	—	—	—							

C. distyla; Ventenat, pl. dans le jard. de Cels. t. 62 (1800)	W.A. S.A. T.	V.	N.S.W.	—	—	B.fl. VI,198	M.fr. VI,19.
C. inophloia; F. v. M. & Bailey in Melb. Chemist, April (1882)	— — —	—	N.S.W.	Q.	—	--	M.fr. XII.
C. Fraseriana; Miquel, Revis. Casuarin. 50, t. 6 (1848)	W.A. — —	—	—	—	—	B.fl. VI,199	M.fr. VI,19.
C. nana; Sieber in Sprengel, syst. III, 804 (1826)	— — —	V.	N.S.W.	—	—	B.fl. VI,199	
C. humilis; Otto & Dietrich, allgem. Gartenzeit. 163 (1841)	W.A. S.A. —	—	—	—	—	B.fl. VI,200	M.fr. VI,18.
C. torulosa; Aiton, hort. Kew. III, 320 (1789)	— — —	—	N.S.W.	Q.	—	B.fl. VI,200	M.fr. VI,17.
C. decussata; Bentham, Fl. Austr. VI, 200 (1873)	W.A. — —	—	—	—	—	B.fl. VI,200	
C. Decaisneana; F. v. M., fragm. I, 61 (1858)	W.A. S.A. —	—	—	Q.	N.A.	B.fl. VI,201	M.fr. I,61; X,62.
C. Drummondiana; Miquel, Revis. Casuar. 26, t. 1 (1848)	W.A. — —	—	—	—	—	B.fl. VI,201	
C. microstachya; Miquel in Lehm. pl. Preiss. I, 642 (1845)	W.A. — —	—	—	—	—	B.fl. VI,201	M.fr. VI,17; X,62.
C. acutivalvis; F. v. M., fragm. X, 61 (1876)	W.A. — —	—	—	—	—	—	M.fr. X,61.
C. bicuspidata; Bentham, Fl. Austr. VI, 202 (1873)	W.A. S.A. T.	—	—	—	—	B.fl. VI,202	
C. thuyoides; Miquel in Lehm. pl. Preiss. I, 641 (1845)	W.A. — —	—	—	—	—	B.fl. VI,202	M.fr. VI,17.
C. corniculata; F. v. M., fragm. X, 62 (1876)	W.A. — —	—	—	—	—	—	M.fr. X,62.
C. acuaria; F. v. M., fragm. VI, 16 (1868)	W.A. — —	—	—	—	—	B.fl. VI,202	M.fr. VI,250.

CELASTRINAE.

R. Brown in App. Flind. voy. 554 (1814).—Genera 10, Species 17.

EUONYMUS; Tournefort, inst. 617, t. 388 (1700) from Theophrastos & Plinius. (Evonymus.)									
E. Australianus; F. v. M., fragm. IV, 118 (1864)	— — —	—	—	Q.	—	—		M.fr. VI,202.	
CELASTRUS; Linné, gen. pl. 59 (1737).									
C. Australis; Harvey & F. v. M. in Trans. phil. Soc. Vict. I, 41 (1854)	— — —	V.	N.S.W.	Q.	—	B.fl. I,398		M.fr. III,93.	
C. Muelleri; Bentham, Fl. Austr. I, 399 (1863)	— — —	—	—	—	N.A.	B.fl. I,399			
C. dispermus; F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859)	— — —	—	—	Q.	—	B.fl. I,399		M.fr. VI,167,203.	
C. bilocularis; F. v. M. in Transact. phil. Inst. Vict. III, 31 (1859)	— — —	—	N.S.W.	Q.	—	B.fl. I,399		M.fr. VI,204.	
C. Cunninghamii; F. v. M. in Transact. phil. Inst. Vict. III, 30 (1859)	— — —	—	N.S.W.	Q.	N.A.	B.fl. I,399		M.fr. VI,203.	
GYMNOSPORIA; Wight & Arnott, prodr. Fl. penins. Ind. I, 159 (1834).									
G. montana; Wight & Arnott, prodr. I, 159 (1834)	— — —	—	—	Q.	—	B.fl. I,400		...	As.
HEDRAIANTHERA; F. v. M., fragm. V, 58 (1865).									
H. porphyropetala; F. v. M., fragm. V, 59 (1865)	— — —	—	N.S.W.	Q.	—	—		M.fr. V,214.	
HYPSOPHILA; F. v. M. in Vict. Nat. III, 169 (1887).									
H. Halleyana; F. v. M. in Vict. Nat. III, 169 (1887)	— — —	—	—	Q.	—	—			
LEUCOCARPUM; A. Richard, sert. Astrolabe 46 (1834). (Denhamia, Leucocarpon.)									
L. Oleaster; F. v. M. in Transact. Roy. Soc. of N.S.W. 15 (1881)	— — —	—	—	Q.	—	B.fl. I,401		M.fr. VI,203.	
L. obscurum; A. Richard, sert. Astrol. 46, t. 18 (1834)	— — —	—	—	Q.	N.A.	B.fl. I,401		M.fr. VI,203.	
L. pittosporoides; F. v. M. in Transact. Roy. Soc. of N.S.W. 15 (1881)	— — —	—	N.S.W.	Q.	—	B.fl. I,402		M.fr. VI,203.	
ELAEODENDRON; J. F. Jacquin in Nov. Act. Helv. I, 36 (1787). (Portenschlagia.)									
E. Australe; Ventenat, Jard. Malm. t. 117 (1804)	— — —	—	N.S.W.	Q.	N.A.	B.fl. I,402		M.fr. III,61,168; IV,175; VI,204; IX,77.	
E. curtipendulum; Endlicher, prodr. fl. ins. Norf. 81 (1833)	— — —	—	N.S.W.	—	—	—			
CARYOSPERMUM; Blume, Mus. Bot. Lugd. I, 175 (1850). (Perrottetiae subgenus.)									
C. arborescens; F. v. M., fragm. VI, 202 (1868)	— — —	—	—	Q.	—	—		M.fr. VIII,279.	

SIPHONODON ; Griffith in Calc. Journ. of Nat. Hist. IV, 150 (1843).

S. Australe ; Bentham, Fl. Austr. I, 403 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,403	M.fr.VI,204.				

HIPPOCRATEA ; Linné, gen. pl. 363 (1737).

H. obtusifolia ; Roxburgh, flor. Ind. edit. Carey, I, 170 (1820)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,404	...	...	...	...	As.		

SALACIA ; Linné, mantiss. II, 159 (1771).

S. prinoides ; De Candolle, prodr. I, 571 (1824)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VI,202.				

SAPINDACEAE.

A. L. DE JUSSIEU, Gen. plant. 246 (1789).—Genera 14, Species 100.

CARDIOSPERMUM ; Linné, syst. nat. (1735) ; gen. pl. 117 (1737).

C. Halicacabum ; Linné, sp. pl. 366 (1753)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,453	M.fr.IX,89	...	As., Afr., Am., P.	

ATALAYA ; Blume, Rumphia III, 186 (1847). (Pseudatalaya, Sapindus partly.)

A. coriacea ; Radlkofer in Sitzungsber. der Akad. zu Münch. 326 (1878)	...	—	—	—	—	N.S.W.	—	—	—									
A. salicifolia ; Blume, Rumphia III, 186 (1847)	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,463	...	...	As.		
A. Australis ; Radlkofer in Sitzungsber. der Akad. zu Münch. 327 (1878)	...	—	—	—	—	—	Q.	—	B.fl.I,464									
A. multiflora ; Bentham, Fl. Austr. I, 463 (1863)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,463	M.fr.IX,90.				
A. hemiglauca ; F. v. M. in Benth. Fl. Austr. I, 463 (1863)	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,463	M.fr.IX,90.						
A. variifolia ; F. v. M. in Benth. Fl. Austr. I, 463 (1863)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,463	M.fr.I,45 ; IX,90.					

DIPLOGLOTTIS ; J. Hooker in H. & B. gen. pl. 395 (1862). (Stadmannia partly.)

D. Cunninghamii ; J. Hooker in H. & B. gen. pl. 395 (1862)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,454	M.fr.V,145 ; IX,90.						

ERIOGLOSSUM ; Blume, Bijdr. 229 (1829). (Pancovia partly.)

E. edule ; Blume, Bijdr. 229 (1829)	...	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,454	M.fr.IX,100	...	As.

CASTANOSPORA ; F. v. M., fragm. IX, 92 (1875).

C. Alphandi ; F. v. M., fragm. IX, 92 (1875)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IV,158 ; IX,197.			

ALLOPHYLUS ; Linné, amoen. acad. 124 (1747). (Schmidelia.)

A. ternatus ; Loureiro, Fl. Cochinch. ed. Willd. 286 (1790)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,455	M.fr.IX,90	...	As., P.				

CUPANIA ; Plumier, Gen. 45, t. 19 (1703). (Guioa, Ratonia, Aryteria, Elattostachys, Mischocarpus, Euphoria partly.)

C. Wadsworthii ; F. v. M., syst. Cens. of Austr. pl. 24 (1882)	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IV,1 ; V,147 ; IX,89,197.						
C. anacardicidea ; A. Richard, sert. Astrol. 33, t. 13 (1833)	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,458	M.fr.IX,91.						
C. Robertsonii ; F. v. M., fragm. V, 146 (1866)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,94.				
C. serrata ; F. v. M., fragm. III, 43 (1862)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,458	M.fr.IX,94,198.			
C. foveolata ; F. v. M., fragm. IX, 95 (1875)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,95.			
C. tomentella ; F. v. M. in Benth. Fl. Austr. I, 458 (1863)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,458	M.fr.IX,94.					
C. Pseudorhus ; A. Richard, sert. Astrol. 34, t. 14 (1833)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,459	M.fr.V,146 ; IX,92.					
C. pleurophylla ; F. v. M. in Vict. Natural. III, 170 (1887)	...	..	...	—	—	—	—	—	Q.	—	—							
C. xylocarpa ; Cunningham in Benth. Fl. Austr. I, 459 (1863)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,459	M.fr.V,7 ; IX,95.						
C. erythrocarpa ; F. v. M., fragm. V, 7 (1865)	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.IX,91.					
C. Mortoniana ; F. v. M., fragm. V, 177 (1866)	...	..	..	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,94.				
C. lachnocarpa ; F. v. M., fragm. IV, 157 (1864)	..	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.V,6 ; IX,91.				
C. grandissima ; F. v. M., fragm. IV, 156 (1864)	...	...	..	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,91.				
C. pyriformis ; F. v. M., App. Rep. Intercol. Exhib. 25 (1867)	...	..	...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,461	M.fr.I,2 ; II,76 ; IX,90.						
C. anodonta ; F. v. M., fragm. II, 76 (1860)	...	...	..	...	..	..	—	—	—	—	—	Q.	—	B.fl.I,461	M.fr.I,2 ; IX,90.			
C. stipitata ; F. v. M., fragm. II, 75 (1860)	...	.	..	.	..	..	—	—	—	—	N.S.W.	—	—	B.fl.I,461	M.fr.III,12 ; IV,157 ; V,6,147,210 ; IX,91.			
C. punctulata ; F. v. M., fragm. III, 12 (1862)	..	..	...	...	..	—	—	—	—	—	Q.	—	B.fl.I,458	M.fr.IX,91.				

C. exangulata ; F. v. M., fragm. IV, 156 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IX,91.		
C. O'Shanesiana ; F. v. M., fragm. IX, 96 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,96.		
C. Daemeliana ; F. v. M., fragm. IX, 96 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,96.		
C. Martyana ; F. v. M., fragm. V, 6 (1865)	—	—	—	—	—	Q.	—	—	M.fr.IX,94.		
C. Cordieri ; F. v. M., fragm. IX, 93 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,93.		
C. tenax ; Cunningham in Benth. Fl. Austr. I, 461 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,461	M.fr.IX,94.		
NEPHELIUM ; Linné, mantiss. 18 (1767). (Euphoria partly, Alectryon, Spanoghea.)											
N. semiglaucum ; F. v. M. in Transact. phil. Soc. Vict. III, 25 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,457	M.fr.IV,158 ; V,147 ; IX,97.		
N. connatum ; F. v. M. in Benth. Fl. Austr. I, 465 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,465	M.fr.IV,158 ; IX,98.		
N. semicinereum ; F. v. M., fragm. IV, 158 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,158.		
N. subdentatum ; F. v. M. in Benth. Fl. Austr. I, 465 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,465	M.fr.IX,99.		
N. tomentosum ; F. v. M. in Transact. Vict. Inst. II, 64 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,466	M.fr.IX,99.		
N. coriaceum ; Bentham, Fl. Austr. I, 466 (1863)	—	—	—	—	—	Q.	—	B.fl.I,466			
N. foveolatum ; F. v. M. in Benth. Fl. Austr. I, 466 (1863)	—	—	—	—	—	Q.	—	B.fl.I,466	M.fr.IX,99.		
N. leiocarpum ; F. v. M. in Transact. phil. Inst. Vict. III, 25 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,467	M.fr.IX,98.		
N. Beckleri ; Bentham, Fl. Austr. I, 467 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,467			
N. divaricatum ; F. v. M. in Benth. Fl. Austr. 467 (1863)	—	—	—	—	—	Q.	—	B.fl.I,467	M.fr.IX,98.		
N. microphyllum ; Bentham, Fl. Austr. I, 468 (1863)	—	—	—	—	—	Q.	—	B.fl.I,468			
N. Leichhardtii ; F. v. M., App. Rep. Intercol. Exhib. 25 (1867)	—	—	—	—	—	Q.	—	B.fl.I,468	M.fr.IX,99.		
N. distyle ; F. v. M., fragm. IX, 99 (1875)	—	—	—	—	—	Q.	—	B.fl.I,462	M.fr.IX,99.		
HETERODENDRON ; Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818).											
H. oleaefolium ; Desfontaines in Mém. du Mus. IV, 8, t. 3 (1818)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,469	M.fr.IX,75,90 ; X,82.		
H. diversifolium ; F. v. M., fragm. I, 46 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,469	M.fr.IX,90 ; X,82.		
HARPULLIA ; Roxburgh, Fl. Ind. ed. Carey II, 44 (1826).											
H. alata ; F. v. M., fragm. II, 103 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,470	M.fr.IX,89,198.		
H. Hillii ; F. v. M. in Transact. Vict. Inst. III, 26 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,470	M.fr.II,104 ; IX,100,197.		
H. Leichhardtii ; F. v. M. in Benth. Fl. Austr. I, 470 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,470			
H. pendula ; Planchon in Transact. Vict. Inst. III, 26 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,471	M.fr.II,104 ; IX,89,100&187.		
AKANIA ; J. Hooker in B. & H. Gen. pl. 409 (1862).											
A. Hillii ; J. Hooker in B. & H. Gen. pl. 409 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,471	M.fr.III,44 ; IX,89&197.		
DIPLOPELTIS ; Endlicher in Huegel, enum. 13 (1837).											
D. Hnegelii ; Endlicher in Hueg. enum. 13 (1837)	W.A.	—	—	—	—	—	N.A.	B.fl.I,456	M.fr.III,12; IX,89.		
D. Stuartii; F. v. M., fragm. III, 12 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.I,456	M.fr.III,12&167; IX,89.		
DODONAEA ; Linné, gen. pl. 341 (1737). (Empleurosma.)											
D. triquetra ; Wendland, Bot. Beobacht. p. 44 (1798)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,474	M.fr.I,73; IX,88.		
D. lanceolata ; F. v. M., fragm. I, 73 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.I,73.		
D. petiolaris ; F. v. M., fragm. III, 13 (1862)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.IX,89.		
D. viscosa ; Linné, mantissa altera 149 & 228 (1771)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,475	M.fr.IX,81	...	As., Afr., Am., P., N.Z.
D. peduncularis ; Lindley in Mitch. Trop. Austr. 361 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,478	M.fr.IX,87.		
D. procumbens ; F. v. M. in Transact. Vict. Inst. I, 3 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,478			
D. ericifolia ; G. Don, gen. syst. I, 674 (1831)	—	—	T.	—	—	—	—	B.fl.I,478	M.fr.IX,87.		
D. filifolia ; Hooker in Mitch. Trop. Austr. 241 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,478	M.fr.I,71.		
D. lobulata ; F. v. M. in Linnaea XXV, 372 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,479	M.fr.IX,87,197.		
D. ptarmicifolia ; Turczaninow in Bull. Soc. Mosc. XXV, part III, 165 (1852)	W.A.	—	—	—	—	—	—	B.fl.I,479	M.fr.I,97; IX,87.		
D. calycina ; Cunningham in A. Gray, Bot. of Wilk. Expl. Exped. I, 262 (1854)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,479	M.fr.II,143; IX,89.		
D. platyptera ; F. v. M., fragm. I, 73 (1858)	W.A.	—	—	—	—	—	N.A.	B.fl.I,480	M.fr.IX,87.		
D. pachyneura ; F. v. M. in Transact. R. S. of Vict. XXIII, 51 (1886)	W.A.	—	—	—	—	—	N.A.	—			
D. stenophylla ; F. v. M., fragm. I, 72 (1858)	—	—	—	—	—	Q.	—	B.fl.I,480	M.fr.IX,87.		

D. pinifolia; Miquel in Lehm. pl. Preiss. I, 227 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,480	M.fr.IX,88.		
D. ceratocarpa; Endlicher in Hueg. enum. 13 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,481	M.fr.X,88.		
D. divaricata; Bentham, Fl. Austr. I, 481 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,481	M.fr.IX,87.		
D. triangularis; Lindley in Mitch. Trop. Austr. 219 (1848)...	—	—	—	—	N.S.W.	Q.	—	B.fl.I,481	M.fr.IX,87.		
D. aptera; Miquel in Lehm. pl. Preiss. I, 225 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,481	M.fr.IX,88.		
D. bursarifolia; Behr & F. v. M. in Transact. Vict. Inst. I, 8 (1855)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,482	M.fr.IX,87.		
D. Baueri; Endlicher in Hueg. enum. 13 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,482	M.fr.IX,87.		
D. trifida; F. v. M., fragm. IX, 88 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,88.		
D. humifusa; Miquel in Lehm. pl. Preiss. I, 266 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,482	M.fr.IX,88.		
D. hexandra; F. v. M. in Transact. Vict. Inst. I, 117 (1855)	—	S.A.	—	—	—	—	—	B.fl.I,483			
D. ericoides; Miquel in Lehm. pl. Preiss. I, 227 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,483	M.fr.IX,88.		
D. polyzyga; F. v. M., fragm. I, 74 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,483	M.fr.I,219.		
D. megazyga; F. v. M. in Benth. Fl. Austr. I, 483 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.I,483	M.fr.IX,86.		
D. macrozyga; F. v. M., fragm. IV, 135 (1864)	—	S.A.	—	—	—	Q.	—	—	M.fr.IX,86.		
D. physocarpa; F. v. M., fragm. I, 74 (1858)...	—	—	—	—	—	Q.	N.A.	B.fl.I,484	M.fr.IX,86.		
D. vestita; Hooker in Mitch. Trop. Austr. 265 (1848)	—	—	—	—	—	Q.	—	B.fl.I,484	M.fr.IX,86.		
D. pinnata; Smith in Rees' Cycl. XII (1809)	—	—	—	—	N.S.W.	—	—	B.fl.I,484			
D. oxyptera; F. v. M., fragm. I, 74 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,484	M.fr.I,74.		
D. humilis; Endlicher, nov. stirp. dec. 26 (1839)	—	S.A.	—	—	—	—	N.A.	B.fl.I,485	M.fr.IX,86.		
D. Macrossani; F. v. M. & Scortechini in Melb. Chemist Jan. 1882	—	—	—	—	—	Q.	—	—	M.fr.XII.		
D. microzyga; F. v. M., annual Rep. Bot. Gard. 12 (1863)	—	S.A.	—	—	—	—	—	—	M.fr.IX,86.		
D. boronifolia; G. Don, gen. syst. I, 674 (1831)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,485	M.fr.IX,86.		
D. multijuga; G. Don, gen. syst. I, 674 (1831)	—	—	—	—	N.S.W.	—	—	B.fl.I,485	M.fr.IX,86.		
D. larraeoides; Turczaninow in Bull. Soc. Mosc. XXXI, 408 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,486	M.fr.I,210; II,182.		
D. inacquifolia; Turczaninow in Bull. Soc. Mosc. XXXI, 408 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,486	M.fr.I,219; IV,135; IX,89.		
D. adenophora; Miquel in Linnaea XVIII, 95 (1845)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.I,486	M.fr.I,98.		
D. tenuifolia; Lindley in Mitch. Trop. Austr. 248 (1848)	—	S.A.	—	—	N.S.W.	Q.	N.A.	—	M.fr.IX,85.		
D. stenozyga; F. v. M., fragm. I, 98 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,486	M.fr.IX,87.		
D. concinna; Bentham, Fl. Austr. I, 487 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,487	M.fr.IX,87.		
DISTICHOSTEMON; F. v. M. in Hook. Kew Misc. IX, 306 (1857). (Pleistichostemon.)											
D. phyllopterus; F. v. M. in Hook. Kew Misc. IX, 306 (1857)	—	—	—	—	—	Q.	N.A.	B.fl.I,487	M.fr.IX,89.		

MALPIGHIACEAE.

VENTENAT; Tabl. III, 131 (1799).—Genera 2, Species 2.

RYSSOPTERYS; Blume in Delessert, ic. sel. III, 21, t. 35 (1837).											
R. Timorensis; Blume in Deless. ic. sel. III, 21, t. 35 (1837)	—	—	—	—	—	Q.	—	B.fl.I,285	...	...	As.
TRISTELLATEIA; Petit-Thouars in Roemer, Collect. bot. 206 (1809).											
T. Australasica; A. Richard, Sert. Astrol. 38, t. 15 (1833)...	—	—	—	—	—	Q.	—	B.fl.I,286	...	...	As., P.

BURSERACEAE.

KUNTH in Ann. des sc. nat. II, 346 (1824).—Genera 3, Species 3.

GARUGA; Roxburgh Pl. of Coromandel III, 5, t. 208 (1819).											
G. floribunda; Descaisne, in nouv. annal. du Mus. III, 477 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.I,377	...	...	As.

CANARIUM; Rumphius, herb. Amboin. II, 145, t. 47, 49, 54 (1741). (Sonzaya.)

C. Australianum; F. v. M., fragm. III, 15 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,378	M.fr.V,211; VII,25.							

GANOPHYLLUM; Blume, mus. bot. Lugd. L, 230 (1850).

G. falcatum; Blume, mus. bot. Lugd. I, 230 (1850)	—	—	—	—	—	Q.	—	—	M.fr.VII,24; IX,193; XI,16.	As.						

ANACARDIACEAE.

R. Brown in Tuckey's Expedit. Congo 431 (1818).—Genera 6, Species 9.

EUROSCHINUS; J. Hooker in B. & H. Gen. pl. I, 422 (1862).

E. falcatus; J. Hooker in B. & H. Gen. pl. I, 422 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,490	M.fr.VII,24.							

RHUS; Tournefort, Inst. 611, t. 381 (1700), from Hippocrates, Theophrastos, Dioscorides & Plinius. (Rhodosphaera.)

R. rhodanthema; F. v. M. in Journ. pharm. soc. Vict. pl. I, 43 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,489	M.fr.VII,23.							
R. rufa; Teysman & Binnendyk in Naturk. Tijd. XXVII, 52 (1863)	—	—	—	—	—	Q.	—	—	M.fr.VII,22.							

BLEPHAROCARYA; F. v. M., fragm. XI, 15 (1878).

B. involucrigera; F. v. M., fr. XI, 16 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,137.							

BUCHANANIA; Sprengel in Schrader's Journ. II, 234 (1800). (Coniogeton.)

B. mangoides; F. v. M., fragm. VII, 23 (1869)	—	—	—	—	—	Q.	—	—	M.fr.VII,23.							
B. obovata; Engler in A. & C. de Candolle, monogr. phaner. IV, 187 (1883) ...	—	—	—	—	—	—	N.A.	—								
B. Muelleri; Engler in A. & C. de Candolle, monogr. phaner. IV, 190 (1883) ...	—	—	—	—	—	Q.	N.A.	B.fl.I,490	M.fr.VII,23.							

SEMECARPUS; Linné fil., suppl. 25 (1781).

S. Anacardium; Linné, fil., suppl. 182 (1781)...	—	—	—	—	—	Q.	N.A.	B.fl.I,491	M.fr.VII,23 ...	As., P.						

SPONDIAS; Linné, gen. pl. 365 (1737). (Pleiogynium.)

S. Solandri; Bentham, Fl. Austr. I, 492 (1863)	—	—	—	—	—	Q.	—	B.fl.I,492	M.fr.III,15; V,177.							

STACKHOUSIEAE.

R. Brown in App. to Flind. Voy. 555 (1814).—Genera 2, Species 13.

STACKHOUSIA; Smith in Transact. Linn. Soc. IV, 218 (1798). (Tripterococcus, Plokiostigma.)

S. pulvinaris; F. v. M. in Transact. phil. Soc. Vict. I, 101 (1855)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,405	M.fr.III,88.							
S. linarifolia; Cunningham in Field's N. S. Wales 365 (1825)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,406	M.fr.III,87; XI,27.							
S. pubescens; A. Richard, sert. Astrol. 89, t. 33 (1834)	W.A.	—	—	—	—	—	—	B.fl.I,407								
S. Huegelii; Endlicher in Hueg. enum. 17 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,407								
S. flava; Hooker, icon. pl. t. 269 (1840)	W.A.	S.A.	T.	V.	—	—	—	B.fl.I,407								
S. muricata; Lindley, Bot. Reg. XXII, 1917 (1837)...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,408	M.fr.VIII,36.							
S. viminea; Smith in Rees' Cycl. XXXIII (1816)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,408	M.fr.III,86; VIII,36.							
S. elata; F. v. M., fragm. III, 86 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,408	M.fr.III,86.							
S. scoparia; Bentham, Fl. Austr. I, 409 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,409								
S. spathulata; Sieber in Spreng. syst. cur. post. 124 (1827)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,406	M.fr.III,86.							
S. megaloptera; F. v. M., fragm. VIII, 35 (1873)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,119.							
S. Brunonis; Bentham, Fl. Austr. I, 409 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.I,409	M.fr.VIII,36.							

MACGREGORIA; F. v. M. in Caruel, Giorn. Bot. Italian. 128 (1873).

M. racemigera; F. v. M. in Caruel, Giorn. Bot. Ital. 128 (1873)	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.VIII,161; XI,134.							

FRANKENIACEAE.

A. de St. Hilaire in Bull. de la Soc. Philom. 22 (1815).—Genus 1, Species 7.

FRANKENIA; Linné, gen. pl. 92 (1737).

F. laevis; Linné, spec. plant. 331 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.I,151	M.fr.IX,75	...	E., As., Afr.
F. bracteata; Turczaninow in Bull. Soc. Mosc. XXVII, 367 (1854)	W.A.	—	—	—	—	—	—	B.fl.I,150			
F. glomerata; Turczaninow in Bull. Soc. Mosc. XXVII, 368 (1854)	W.A.	—	—	—	—	—	—	B.fl.I,151			
F. parvula; Turczaninow in Bull. Soc. Mosc. XXVII, 368 (1854)	W.A.	—	—	—	—	—	—	B.fl.I,152			
F. Drummondii; Bentham, Fl. Austr. I, 152 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,152			
F. tetrapetala; Labillardière, nov. Holl. pl. spec. I, 88, t. 114 (1804)	W.A.	—	—	—	—	—	—	B.fl.I,152			
F. punctata; Turczaninow in Bull. Soc. Mosc. XXVII, 367 (1854)	W.A.	—	—	—	—	—	—	B.fl.I,153			

PLUMBAGINEAE.

A. L. de Jussieu, gen. plant. 92 (1789).—Genera 3, Species 4.

PLUMBAGO; Tournefort, inst. 140, t. 58 (1700).

P. Zeilanica; Linné, sp. pl. 151 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,267	M.fr.XI,7	...	As., Afr., P.

STATICE; Linné, gen. pl. 88 (1737). (Taxanthema.)

S. Taxanthema; Roemer & Schultes, syst. veg. VI, 798 (1820)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,267	M.fr.XI,7	...	As., P.
S. salicorniacea; F. v. M., fragm. XI, 7 (1878)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,7.		

AEGIALITIS; R. Brown, prodr. 426 (1810). (Aegialinites.)

A. annulata; R. Brown, prodr. 426 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,266	M.fr.XI,7	...	P.

PORTULACEAE.

A. L. de Jussieu, gen. plant. 312 (1789).—Genera 3, Species 32.

PORTULACA; Tournefort, inst. 236, t. 118 (1700) from W. Turner (1538).

P. oleracea; Linné, spec. plant. 445 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.I,169	M.fr.X,71	...	E., As., Afr., Am., P.
P. napiformis; F. v. M. in Benth. Fl. Austr. I, 169 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.I,169	M.fr.X,72	...	As.
P. Australis; Endlicher, Atakta 7, t. 6 (1833)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,169	M.fr.X,72.		
P. filifolia; F. v. M., fragm. I, 169 (1859)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,169	M.fr.III,164; X,72.		
P. digyna; F. v. M., fragm I, 170 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.I,170.		
P. oligosperma; F. v. M., fragm. I, 170 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.III,164; X,72.	...	As.
P. bicolor; F. v. M., fragm. I, 171 (1859)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,170	M.fr.III,164; X,72.		
P. Armitii; F. v. M., fragm. X, 97 (1877)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,97.		

CLAYTONIA; Gronovius in Linné, gen. pl. 339 (1737). (Calandrinia, Talinum, Montiae subgenus.)

C. spergularina; F. v. M., Cens. 27 (1882)	—	—	—	—	—	Q.	N.A.	B.fl.I,176	M.fr.I,175; IV,137; X,71.		
C. ptychosperma; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	—	—	Q.	N.A.	—	M.fr.IV,137; X,70.		
C. strophiolata; F. v. M., fragm. XI, 82 (1880)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,82.		
C. Lehmanni; F. v. M., Cens. 27 (1882) from Endl. (1847)	W.A.	—	—	—	—	—	—	B.fl.I,172	M.fr.X,71.		
C. uniflora; F. v. M., Cens. 27 (1882)	—	—	—	—	—	Q.	N.A.	B.fl.I,172	M.fr.X,71.		
C. Balonnensis; F. v. M., Cens. 27 (1882) from Lindl. (1848)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,172	M.fr.I,177; X,71.		
C. polyandra; F. v. M., Cens. 27 (1882) from Hook. (1855)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,172	M.fr.I,177; X,11.		
C. pleiopetala; F. v. M., Cens. 27 (1882)	—	S.A.	—	—	—	Q.	—	—	M.fr.X,70.		
C. quadrivalvis; F. v. M., Cens. 27 (1882)	—	—	—	—	—	—	N.A.	B.fl.I,173	M.fr.I,176.		

C. liniflora ; F. v. M., Cens. 27 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,173	M.fr.X,71.		
C. gracilis ; F. v. M., Cens. 27 (1882) from Benth. (1863)	—	—	—	—	—	—	N.A.	B.fl.I,173			
C. polypetala ; F. v. M., Cens. 27 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,174			
C. Pickeringi ; F. v. M., Cens. 27 (1882) from A. Gray (1854)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,69 ; XI,136.		
C. volubilis ; F. v. M., Cens. 27 (1882) from Benth. (1863)...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,174	M.fr.X,68 ; XI,27.		
C. calyptrata ; F. v. M., fragm. III, 89 (1862)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,174	M.fr.I,175,177 ; IV,175 ; X,68.		
C. pumila ; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	—	M.fr.X,68.		
C. composita ; F. v. M., Cens. 27 (1882) from Nees (1844)...	W.A.	—	—	—	—	—	—	B.fl.I,175	M.fr.,X,69.		
C. corrigioloides ; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,175	M.fr.X,69 ; XI,83.		
C. brevipedata ; F. v. M., Cens. 27 (1882)	W.A.	S.A.	—	V.	—	—	—	—	M.fr.X,69 ; IX,136.		
C. granulifera ; F. v. M., Cens. 27 (1882) from Benth. (1863)	W.A.	—	—	—	—	—	—	B.fl.I,176			
C. pogonophora ; F. v. M., Cens. 27 (1882)	—	—	—	—	—	Q.	—	—	M.fr.X,69.		
C. pygmaea ; F. v. M., fragm. III, 89 (1862)...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,176	M.fr.I,175 ; X,71 ; XI,83.		
C. Australasica ; J. Hooker, icon. plant. t. 293 (1840)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,177	M.fr.III,89 ; X,71 ...	N.Z.	
MONTIA ; Micheli, nov. pl. gen. 17, t. 13 (1729).											
M. fontana ; Linné, spec. plant. 82 (1753)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,177	... ...	E., As., Am., N.Z.	

CARYOPHYLLEAE.

LINNE, Philosoph. Bot. 31 (1751).—Genera 10, Species 26.

STELLARIA ; Linné, syst. nat. ed. VI, (1748).											
S. pungens ; Brongniart, Voy. sur la Coquille, t. 78 (1826)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,157			
S. palustris ; Retzius, Fl. Scandinav., prodr. sec. ed. n. 548 (1779)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,158	... ...	E., As., Am.	
S. flaccida ; Hooker, Comp. Bot. Mag. I, 275 (1835)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,158			
S. multiflora ; Hooker, Comp. Bot. Mag. I, 275 (1835)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,159	M.fr.IX,27.		
SAGINA ; Linné, syst. nat. 8 (1735) ; gen. pl. 118 (1737).											
S. procumbens ; Linné, spec. plant. 128 (1753)	—	—	—	V.	N.S.W.	—	—	B.fl.I,160	.. ..	E., As., Afr. Am.	
S. apetala ; Arduino, animadv. Bot. spec. alt. 22, t. 8 (1764)	W.A.	S.A.	T.	V.	—	—	—	B.fl.I,160	.. ..	E., As., Afr.	
COLOBANTHUS ; Bartling, Reliq. Haenk. II, 13, t. 49 (1830).											
C. Benthamianus ; Fenzl in Ann. des Wien. Mus. I, 49 (1838)	—	—	—	V.	N.S.W.	—	—	B.fl.I,160	... .	Am., N.Z.	
C. Billardieri ; Fenzl in Ann. des Wien. Mus. I, 48 (1838)...	—	—	T.	V.	—	—	—	B.fl.I,161			
SCLERANTHUS ; Linné, gen. plant. 130 (1737). (Mniarum, Ditoca.)											
S. pungens ; R. Brown, prodr. 412 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,260			
S. diander ; R. Brown, prodr. 412 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,260			
S. biflorus ; J. Hooker, Fl. N. Zeal. I, 74 (1853)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,259	... ...	N.Z.	
S. mniaroides ; F. v. M., pl. of Vict. I, 215 (1862)	—	—	—	V.	N.S.W.	—	—	B.fl.V,259			
SAPONARIA ; Linné, gen. plant. 130 (1737) from Bock (1529). (Gypsophila, Dichoglottis.)											
S. tubulosa ; F. v. M., native pl. of Vict. I, 136 (1879)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.I,155			
SPERGULARIA ; Persoon, synops. pl. I, 504 (1805). (Tissa (1763), Lepigonum, Arenaria partly.)											
S. rubra ; Cambessèdes in St. Hilaire, Fl. Brazil. II, 179 (1829)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,162	M.fr.XI,27...	..	E., As., Am.
DRYMARIA ; Wildenow in Roemer & Schultes, syst. V, 406 (1819).											
D. filiformis ; Bentham, Fl. Austr. I, 162 (1863)	W.A.	S.A.	—	—	—	—	—	B.fl.I,162			
D. diandra ; Blume, Bijdr. 63 (1823)	—	—	—	—	—	Q.	—	—	... ...	As.	
HERNIARIA ; Tournefort, instit. rei. herb. 507, t. 288 (1700).											
H. incana ; Lamarck, encyl. meth. III, 124 (1789)	—	S.A.	—	—	—	—	—	—	... ...	E., As., Afr.	

POLYCARPON; Loefling, iter Hispanic. 7 (1758).

P. tetraphyllum; Loefling in Linné, syst. nat. ed. X, 881 (1759)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,163	...	...	...	E., As., Afr.

POLYCARPAEA; Lamarck in Journ. d'hist. nat. II, 8, t. 25 (1792). (Aylmeyera.)

P. longiflora; F. v. M., Rep. of Babb. Exped. 8 (1858)	W.A.	—	—	—	—	—	N.A.	B.fl.I,164				
P. spirostylis; F. v. M., Rep. of Babb. Exped. 8 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,165				
P. synandra; F. v. M., Rep. of Babb. Exped. 8 (1858)	—	S.A.	—	—	—	Q.	N.A.	B.fl.I,165				
P. violacea; Bentham, Fl. Austr. I, 165 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,165				
P. Indica; Lamarck in Journ. d'hist. nat. II, 8 (1792)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,166	...	...	...	As., Afr., Am.
P. breviflora; F. v. M., Rep. of Babb. Exped. 9 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.I,166				
P. spicata; Arnott in Ann. of nat. hist. III, 91 (1839)	—	—	—	—	—	—	N.A.	B.fl.I,167				
P. involucrata; F. v. M., Rep. of Babb. Exped. 9 (1858)	—	—	—	—	—	—	N.A.	B.fl.I,167				

AMARANTACEAE.

A. L. de Jussieu, gen. plant. 87 (1789).—Genera 12, Species 100.

GOMPHRENA; Linné, gen. plant. 69 (1737). (Philoxerus.)

G. canescens; R. Brown, prodr. 416 (1810)	W.A.	—	—	—	—	Q.	N.A.	B.fl.V,253				
G. flaccida; R. Brown, prodr. 416 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,254	M.fr.III,123.			
G. affinis; F. v. M. in Benth. Fl. Austr. V, 254 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,254				
G. humilis; R. Brown, prodr. 416 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,254				
G. platandra; F. v. M. in Trans. R. S. Vict. XXIII, 52 (1886)	W.A.	—	—	—	—	—	N.A.	—				
G. lanata; R. Brown, prodr. 416 (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.V,255				
G. brachystylis; F. v. M., fragm. III, 124 (1862)	—	—	—	—	—	—	N.A.	B.fl.V,255	M.fr.III,124.			
G. leptoclada; Bentham, Fl. Austr. V, 255 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,255				
G. Maitlandi; F. v. M., fragm. III, 124, t. 23 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.V,256	M.fr.VI,234.			
G. pusilla; Bentham, Fl. Austr. V, 256 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,256				
G. tenella; Bentham, Fl. Austr. V, 256 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,256				
G. conica; Sprengel, syst. I, 824 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.V,256	M.fr.III,125.			
G. conferta; Bentham, Fl. Austr. V, 257 (1870)	W.A.	—	—	—	—	Q.	N.A.	B.fl.V,257				
G. diffusa; Sprengel, syst. I, 824 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.V,257				
G. parviflora; Bentham, Fl. Austr. V, 257 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,257				

ALTERNANTHERA; Forskael, flor. Aegypt. Arab. 28 (1775). (Telanthera.)

A. triandra; Lamarck, encycl. meth. I, 95 (1783)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,249	...	...	As., Afr.	
A. nana; R. Brown, prodr. 417 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,250				
A. angustifolia; R. Brown, prodr. 417 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.V,250				
A. decipiens; Bentham, Fl. Austr. V, 251 (1870)	—	—	—	—	—	Q.	—	B.fl.V,251				
A. polycephala; Bentham, Fl. Austr. V, 251 (1870)...	—	—	—	—	—	—	N.A.	B.fl.V,251				
A. leptophylla; Bentham, Fl. Austr. V, 251 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,251				
A. longipes; Bentham, Fl. Austr. V, 252 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,252				

ACHYRANTHES; Linné, gen. plant. 34 (1737).

A. aspera; Linné, spec. plant. 204 (1753)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,246	M.fr.IX,77.			
A. arborescens; R. Brown, prodr. 417 (1810)...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.			

NYSSANTHES; R. Brown, prodr. 418 (1810).

N. erecta; R. Brown, prodr. 418 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,247				
N. diffusa; R. Brown, prodr. 418 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,247				

DIPTERANTHEMUM; F. v. M. in Wing's S. Sc. Rec. III, 281 (1884).

D. Crosslandi; F. v. M. in Wing's S. Sc. Rec. III, 281 (1884)	W.A.	—	—	—	—	—	—	—					

PTILOTUS; R. Brown, prodr. 415 (1810). (Trichinium, Psilotrichum, Goniotriche, Hemisteirus, Arthrotrichum.)

P. conicus; R. Brown, prodr. 415 (1810)	—	—	—	—	—	—	N.A.	B.fl.V,242					
P. Macleayi; F. v. M. in proceed. L. S. of N.S.W. sec. ser. III, 162 (1888) ...	—	—	—	—	—	—	N.A.	—					
P. corymbosus; R. Brown, prodr. 415 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,243	M.fr.III,125.				
P. grandiflorus; F. v. M., fragm. I, 237 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,243	M.fr.I,237.				
P. spicatus; F. v. M. in Benth. Fl. Austr. V, 243 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,243					
P. Murrayi; F. v. M., fragm. III, 145 (1863)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,243	M.fr.III,145.				
P. gomphrenoides; F. v. M., fragm. VI, 233 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.V,244					
P. latifolius; R. Brown in App. Sturt's Exped. 25 (1849)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.V,244	M.fr.III,145; VI,232.				
P. macrotrichus; F. v. M., fragm. IV, 90 (1864)	W.A.	—	—	—	—	—	N.A.	B.fl.V,244	M.fr.VI,232.				
P. villosiflorus; F. v. M., fragm. III, 125 (1863)	—	—	—	—	—	—	N.A.	B.fl.V,245	M.fr.III,125.				
P. humilis; F. v. M., fragm. VI, 229 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,245	M.fr.III,161; VI,229.				
P. obovatus; F. v. M., fragm. VI, 228 (1868)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,220	M.fr.XI,27.				
P. incanus; Poiret, Encycl. méth. suppl. IV, 620 (1817)	—	S.A.	—	—	—	—	N.A.	B.fl.V,221	M.fr.VI,228.				
P. parviflorus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Lindley (1838)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,222					
P. astrolasius; F. v. M., fragm. VI, 233 (1868)	—	—	—	—	—	—	N.A.	B.fl.V,222	M.fr.VI,233.				
P. rotundifolius; F. v. M., fragm. VI, 230 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,223	M.fr.III,122.				
P. dissitiflorus; F. v. M., syst. Cens. of Austr. pl. 28 (1882)	—	—	—	—	—	—	N.A.	B.fl.V,223	M.fr.IV,89.				
P. distans; Poiret, Encycl. suppl. IV, 620 (1817)	—	—	—	—	—	Q.	N.A.	B.fl.V,223	M.fr.VI,228.				
P. alopecuroideus; F. v. M., fragm. VI, 227 (1868)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,224	M.fr.VI,227.				
P. nobilis; F. v. M., fragm. VI, 227 (1868)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,224	M.fr.VI,227.				
P. macrocephalus; Poiret, Encycl. suppl. IV, 620 (1817)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,225	M.fr.VI,228; XI,27.				
P. hemisteirus; F. v. M., fragm. IV, 90 (1864)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.V,226	M.fr.VI,231.				
P. parvifolius; F. v. M., fragm. VI, 229 (1868)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,241	M.fr.VI,229.				
P. Polakii; F. v. M. in S. Sc. Rec. II, 12 (1882)	W.A.	—	—	—	—	—	—	—					
P. exaltatus; Nees in Lehm. pl. Preiss. I, 630 (1845)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,227					
P. Manglesii; F. v. M., fragm. VI, 230 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.V,228	M.fr.VI,230.				
P. Beckeri; F. v. M., fragm. VI, 233 (1868)	—	S.A.	—	—	—	—	—	B.fl.V,228					
P. gomphrenoides; Moquin in De Cand., prodr. XIII, 2, 287 (1849)	W.A.	S.A.	—	—	—	—	—	B.fl.V,229					
P. esquamatus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Bentham (1838)	W.A.	—	—	—	—	—	—	B.fl.V,229					
P. declinatus; Nees in Lehm. pl. Preiss. I, 631 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,230					
P. erubescens; Schlechtendal in Linnaea XX, 575 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,230	M.fr.VI,229.				
P. divaricatus; F. v. M., fragm. VI, 229 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.V,231	M.fr.VI,229.				
P. helipteroides; F. v. M., fragm. VI, 231 (1868)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,231	M.fr.III,122&161.				
P. Stirlingi; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Lindley (1839) ...	W.A.	—	—	—	—	—	N.A.	B.fl.V,232					
P. laxus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Bentham (1870) ...	W.A.	—	—	—	—	—	—	B.fl.V,232					
P. axillaris; F. v. M., syst. Cens. of Austr. pl. 28 (1882)	W.A.	—	—	—	—	—	N.A.	B.fl.V,232					
P. striatus; F. v. M., syst. Cens. of Austr. pl. 28 (1882) from Moquin (1870) ...	W.A.	—	—	—	—	—	N.A.	B.fl.V,233					
P. auriculifolius; F. v. M., Cens. of Austr. pl. 28 (1882) from Cunningham (1849)	—	—	—	—	—	—	N.A.	B.fl.V,233					
P. sericostachyus; F. v. M., fragm. VI, 231 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,234	M.fr.VI,230.				
P. roseus; F. v. M., syst. Cens. of Austr., pl. 28 (1882) from Moquin (1849) ...	W.A.	—	—	—	—	—	—	B.fl.V,234					
P. fusiformis; Poiret, Encycl. suppl. IV, 619 (1817)	—	—	—	—	—	Q.	N.A.	B.fl.V,234					
P. gracilis; Poiret, Encycl. suppl. IV, 620 (1817)	W.A.	—	—	—	—	Q.	N.A.	B.fl.V,235	M.fr.VI,228.				
P. Drummondii; F. v. M., fragm. VI, 229 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,235	M.fr.VI,229.				
P. calostachyus; F. v. M., fragm. VI, 231 (1868)	—	—	—	—	—	Q.	N.A.	B.fl.V,236	M.fr.VI,231.				
P. Fraseri; F. v. M., Cens. 29 (1882) from Cunningham (1849)	—	—	—	—	N.S.W.	—	—	B.fl.V,236					
P. Carlsoni; F. v. M. in Vict. Nat. V, 74 (1888)	W.A.	—	—	—	—	—	—						
P. Sewelliae; F. v. M. in Vict. Nat. Jan. (1885)	W.A.	—	—	—	—	—	—						

P. Polakii; F. v. M. in Wing's S. Sc. Rec. II, 274 (1882)	W.A.	—	—	—	—	—	—							
P. spathulatus; Poiret, Encycl. suppl. IV, 620 (1817)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,236	M.fr.VI,228; XI,27.					
P. pyramidatus; F. v. M., fragm. VI, 230 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,237	M.fr.VI,230.					
P. holosericeus; F. v. M., fragm. VI, 229 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,237	M.fr.VI,229.					
P. lanatus; Cunningham in De Candolle, prodr. XIII, p. II, 281 (1849)... ...	—	—	—	—	—	—	N.A.	B.fl.V,238						
P. leucocoma; F. v. M., Cens. 29 (1882) from Moquin (1849)	—	S.A.	—	—	—	—	—	B.fl.V,238						
P. brachyanthus; F. v. M., Cens. 29 (1882)	—	—	—	—	—	—	N.A.	B.fl.V,239						
P. arthrolasius; F. v. M., fragm. VI, 232 (1868)	—	—	—	—	—	—	N.A.	B.fl.V,239	M.fr.VI,232.					
P. Forrestii; F. v. M., pl. of N. W. Austr. 7 (1881)	—	—	—	—	—	—	N.A.	—						
P. aeruoides; F. v. M., fragm. VI, 231 (1868)	—	—	—	—	—	Q.	N.A.	B.fl.V,240	M.fr.III,123.					
P. Roei; F. v. M., Cens. 29 (1882)	W.A.	—	—	—	--	—	—	B.fl.V,240						
P. Hoodii; F. v. M., fragm. VIII, 232 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,238.					
P. caespitulosus; F. v. M., fragm. VI, 232 (1868)	W.A.	—	—	—	—	—	--	B.fl.V,240	M.fr.VI,232.					
P. helichrysoides; F. v. M., fragm. VI, 231 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,241	M.fr.I,237.					
P. psilotrichoides; F. v. M., fragm. VI, 231 (1869)...	W.A.	—	—	—	--	—	N.A.	B.fl.V,209	M.fr.I,238; XI,97.					
CYATHULA; Loureiro, flora Cochinchin. I, 101 (1790).														
C. geniculata; Loureiro, fl. Coch. I, 101 (1790)	—	—	—	—	—	—	N.A.	...	...	...	...	As., Afr., Am., P.		
EUXOLUS; Rafinesque, Fl. Tell. 42 (1836).														
E. Mitchelli; F. v. M. in Giles, geogr. trav. Centr. Austr. 214 (1875)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,214						
E. interruptus; Moquin in De Candolle, prodr. XIII, pars II, 275 (1849) ...	—	—	—	—	—	Q.	N.A.	B.fl.V,215						
E. viridis; Moquin in De Candolle, prodr. XIII, pars II, 273 (1849)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.V,215	...	...	...	E., As., Afr., Am., P.		
E. macrocarpus; F. v. M., Cens. 29 (1882) from Benth. (1870)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,216						
AMARANTUS; Dodoens, stirp. hist. pempt. 185 (1583).														
A. leptostachyus; Bentham, Fl. Austr. V, 214 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,213						
A. pallidiflorus; F. v. M., fragm. I, 140 (1859)	W.A.	—	—	—	—	Q.	N.A.	B.fl.V,214	M.fr.I,140.					
POLYCNEMON; Linné, gen. pl. ed. sec. 21 (1742). (Hemichroa.)														
P. pentandrum; F. v. M. in papers R.S. of Tasm. 115 (1877)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,211	M.fr.VI,234.					
P. diandrum; F. v. M., native pl. of Vict. I, 163 (1879)	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,211	M.fr.VI,234.					
P. mesembrianthemum; F. v. M., fragm. VIII, 38 (1873)...	—	S.A.	—	—	—	—	—	—	M.fr.VIII,38.					
CELOSIA; Linné, gen. plant. 34 (1737).														
C. cristata; Linné, sp. pl. 205 (1753)	—	—	—	—	—	Q.	—	—	...	...	...	As., P.		
DEERINGIA; R. Brown, prodr. 413 (1810). (Lestibudesia partly, Celosia partly, Lagrezia.)														
D. celosioides; R. Brown, prodr. 413 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,210	M.fr.II,92	...	As., P.			
D. altissima; F. v. M., fragm. II, 92 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,210	M.fr.VI,251					

SALSOLACEAE.

LINNE, Class. pl. 507 (1738).—Genera 13, Species 111.

ATRIPLEX; Tournefort, inst. 506, t. 288 (1700) from l'Obel (1581) from Turner (1538). (Obione, Theleophyton.)

A. fissivalve; F. v. M., fragm. IX, 123 (1875)	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.XII,17.					
A. crystallinum; J. Hooker in Lond. Journ. of Bot. VI, 279 (1847)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,180	M.fr.IV,49; VII,9; XII,15.					
A. leptocarpum; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,178	M.fr.VII,9; XI,27; XII,17.					
A. limbatum; Bentham, Fl. Austr. V, 178 (1870)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,178						
A. velutinellum; F. v. M., Rep. Babb. Exp. 20 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,174	M.fr.VIII,9.					
A. elachophyllum; F. v. M., fragm. VII, 8 (1870)	—	—	—	—	—	—	N.A.	B.fl.V,176	M.fr.VII,8.					

A. Muelleri; Bentham, Fl. Austr. V, 175 (1870)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,175	M.fr.VII,9.
A. semibaccatum; R. Brown, prodr. 406 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,175	M.fr.VI,19.
A. humile; F. v. M., fragm. IV, 48 (1866)	—	—	—	—	—	—	N.A.	B.fl.V,174	M.fr.IV,48.
A. prostratum; R. Brown, prodr. 406 (1810)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.V,176	M.fr.VII,9; XII,17.
A. angulatum; Bentham, Fl. Austr. V, 174 (1870)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,174	
A. Quinii; F. v. M. in Vict. Nat. V, 96 (1888)	—	S.A.	—	—	N.S.W.	Q.	—		
A. stipitatum; Bentham, Fl. Austr. V, 168 (1870)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,168	M.fr.VII,9.
A. paludosum; R. Brown, prodr. 406 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.V,169	M.fr.VII,9; XII,16.
A. Moquinianum; Webb in De Candolle, prodr. XIII, pars II, 97 (1849)	W.A.	—	—	—	—	—	N.A.	B.fl.V,169	
A. Drummondi; Moquin in De Candolle, prodr. XIII, pars II, 102 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,170	
A. cinereum; Poiret, Encycl. suppl. I, 471 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,171	M.fr.VII,9; IX,77.
A. rhagodioides; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,172	
A. incrassatum; F. v. M., Rep. Babb. Exp. 20 (1858)	—	[S.A.	—	—	—	—	—	B.fl.V,172	
A. nummularium; Lindley in Mitch. Trop. Austr. 64 (1848)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,170	M.fr.VII,10; XI,27; XII,17.
A. hymenothecum; Moquin in De Candolle, prodr. XIII, pars II, 101 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,173	M.fr.VII,10.
A. vesicarium; Heward in Benth. Fl. Austr. V, 172 (1870)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,172	
A. balimoides; Lindley in Mitch. Three Exped. I, 282 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,179	M.fr.VII,10; XI,27; XII,17.
A. spongiosum; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,179	M.fr.XII,17.
RHAGODIA; R. Brown, prodr. 408 (1810).									
R. Billardieri; R. Brown, prodr. 408 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,152	M.fr.VII,10; IX,77; XII,16.
R. parabolica; R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,153	M.fr.VII,10; XII,15.
R. dioica; Nees in Lehm. pl. Preiss. I, 636 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,154	
R. Gaudichaudiana; Moquin, Chenop. monogr. enum. 11 (1840)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,154	M.fr.VII,10; XII,15.
R. crassifolia; R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,154	M.fr.XII,15.
R. Pressii; Moquin in De Candolle, prodr. XIII, part II, 49 (1849)	W.A.	S.A.	—	—	—	—	—	B.fl.V,155	M.fr.VII,10; XII,15.
R. obovata; Moquin, Chenop. monogr. enum. 10 (1840)	W.A.	—	—	—	—	—	—	B.fl.V,155	
R. spinescens; R. Brown, prodr. 408 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,155	M.fr.VII,10; XI,20,75; XII,15.
R. hastata; R. Brown, prodr. 408 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,156	M.fr.VII,10.
R. nutans; R. Brown, prodr. 408 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,156	M.fr.VII,10; XII,15.
R. linifolia; R. Brown, prodr. 408 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,157	M.fr.VII,10; XII,15.
CHENOPODIUM; Tournefort, inst. 506, t. 288 (1700). (Blitum, Ambrina.)									
C. nitrariaceum; F. v. M. in Benth. Fl. Austr. V, 158 (1870)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,158	M.fr.XI,20&27; XII,16.
C. auricomum; Lindley in Mitch. Trop. Austr. 94 (1848)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,159	M.fr.VII,11.
C. triangulare; R. Brown, prodr. 407 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,160	M.fr.VII,10.
C. microphyllum; F. v. M. in Transact. phil. Inst. Vict. II, 74 (1857)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,161	M.fr.VII,11; XII,16.
C. atriplicinum; F. v. M., fragm. VII, 11 (1869)	—	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,163	M.fr.I,140; XI,27.
C. cristatum; F. v. M., fragm. VII, 11 (1869)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,163	M.fr.XI,27; IX,75; XII,17.
C. carinatum; R. Brown, prodr. 407 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,162	M.fr.VII,11 ... N.Z., P.
C. rhadinostachyum; F. v. M. in Wing's S. Sc. Rec. May (1880)	—	S.A.	—	—	—	—	—	—	M.fr.XII.
C. simulans, F. v. M. & Tate; Dysphania, F. v. M. & T. in proc. Roy. Soc. S.A. VIII, 71 (1885)	—	S.A.	—	—	—	—	—	—	
DYSPHANIA; R. Brown, prodr. VI, 411 (1810).									
D. plantaginella; F. v. M., fragm. I, 61 (1858)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,164	M.fr.IX,75
D. litoralis; R. Brown, prodr. 411 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,164	M.fr.VII,11; XII,17.
KOCHIA; Roth in Schrader's Journ. Bot. I, 303 (1799). (Maireana, Sclerochlamys.)									
K. fimbriolata; F. v. M., fragm. IX, 75 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,75.
K. lobiflora; F. v. M. in Benth. Fl. Austr. V, 184 (1870)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,184	M.fr.IX,75.
K. lanosa; Lindley in Mitch. Trop. Austr. 88 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,184	M.fr.VII,13; XII,15
K. prosthecochaeta; F. v. M., fragm. XII, 14 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII,14.
K. melanocoma; F. v. M., fragm. XII, 14 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII,14.

K. triptera; Bentham, Fl. Austr. V, 185 (1870)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,185	
K. oppositifolia; F. v. M. in Transact. Vict. Inst. 134 (1855)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,185	M.fr.VII,12; XII,15.
K. brevifolia; R. Brown, prodr. 409 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,185	M.fr.VII,13.
K. pyramidata; Bentham, Fl. Austr. V, 186 (1870)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,186	M.fr.XII,15.
K. spongiocarpa; F. v. M. in Vict. Nat. III, 92, (1886)	W.A.	S.A.	—	—	N.S.W.	Q.	—	—	
K. eriantha; F. v. M., Rep. Babb. Exped. 20 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,186	M.fr.VII,13.
K. villosa; Lindley in Mitch. Trop. Austr. 91 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,186	M.fr.VII,12; IX,75; XII,15.
K. planifolia; F. v. M., fragm. I, 213 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,187	M.fr.VII,12.
K. sedifolia; F. v. M. in Transact. Vict. Inst. I, 134 (1855)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,187	M.fr.VII,12; VIII,37; XII,15.
K. humillima; F. v. M., fragm. IX, 168 (1875)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.XII,15.
K. microphylla; F. v. M., fragm. VIII, 148 (1874)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,181	M.fr.VIII,37; XII,15.
K. dichoptera; F. v. M., fragm. VIII, 37 (1873)	—	—	—	—	—	Q.	—	—	M.fr.VIII,163.
K. decaptera; F. v. M., fragm. IX, 75 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,75.
K. ciliata; F. v. M., Rep. Babb. Exped. 20 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,188	M.fr.VII,12; XII,15.
K. brachyptera; F. v. M., second gen. Rep. 15 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,189	M.fr.VII,13; XII,15.
K. stelligera; F. v. M., fragm. VII, 13 (1869)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,189	M.fr.I,139; XII,15.
DIDYMANTHUS; Endlicher, nov. stirp. decad. 7 (1839).									
D. Roei; Endlicher, nov. stirp. decad. 8 (1839)	W.A.	—	—	—	—	—	N.A.	B.fl.V,193	M.fr.VII,11.
BABBAGIA; F. v. M., Rep. on pl. of Babb. Exped. 21 (1858).									
B. dipterocarpa; F. v. M., Rep. of Babb. Exped. 21 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,192	M.fr.VII,11.
B. scleroptera; F. v. M. in Wing's S. Sc. Rec. Nov. (1885)	—	—	—	—	N.S.W.	Q.	—	—	
B. acroptera; F. v. M. & Tate in Transact. Roy. Soc. of S. Austr. VI, 108 (1883)	—	S.A.	—	—	—	Q.	N.A.	—	
B. pentaptera; Tate in Transact. Roy. Soc. of S. Austr. VI, 108 (1883)	—	S.A.	—	—	—	—	—	—	
BASSIA; Allioni in Misc. Taurin. III, 117, t. 4 (1766). (Chenolea, Sclerolaena, Anisacantha, Echinopsilon, Kentropsis, Dissocarpus, Eriochiton, Osteocarpum, Coilocarpus.)									
B. carnosa; F. v. M., Cens. of Austr. pl. 30 (1882) from Bentham (1870)	W.A.	—	—	—	—	—	—	B.fl.V,190	M.fr.VIII,13; XII,13.
B. Dallachyana; F. v. M., Cens. of Austr. pl. 30 (1882) from Bentham (1870)	—	—	—	V.	N.S.W.	—	—	B.fl.V,191	
B. tricornis; F. v. M., Cens. of Austr. pl. 30 (1882) from Bentham (1870)	W.A.	S.A.	—	—	—	—	—	B.fl.V,191	
B. tridens; F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII,12.
B. eurotioides; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A.	—	—	—	—	—	—	B.fl.V,191	
B. Muelleri; F. v. M., Cens. of Austr. pl. 30 (1882) from Bentham (1870)	—	—	—	—	—	—	N.A.	B.fl.V,191	M.fr.XII,13.
B. astrocarpa; F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII,12.
B. sclerolaenoides; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,192	M.fr.VII,13; XII,13.
B. Cornishiana; F. v. M. in Austr. Chemist (1885)	—	S.A.	—	—	—	—	—	—	
B. uniflora; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.V,194	M.fr.VIII,38.
B. diacantha; F. v. M., Cens. of Austr. pl. 30 (1882) from Nees (1844)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,194	M.fr.VII,14; IX,75; X,91.
B. lanicuspis; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,195	M.fr.II,140,170; VII,14; IX.27; XII,13.
B. bicornis; F. v. M., Cens. of Austr. pl. 30 (1882) from Lindley (1838)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,195	M.fr.I,139; VII,14; XII,13.
B. biflora; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1838)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,196	M.fr.VII,11; VIII,38; IX.27; X,91; XII,13.
B. paradoxa; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,196	M.fr.VII,11; VIII,38,149; X,91; XII,13.
B. quinquecuspis; F. v. M., Cens. of Austr. pl. 30 (1882)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,199	M.fr.VII,14; X,91.
B. Forrestiana; F. v. M., fragm. XII, 12 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.XII,12.
B. Drummondii; F. v. M., Cens. of Austr. pl. 30 (1882) from Bentham (1870)	W.A.	—	—	—	—	—	—	B.fl.V,199	
B. Birchii; F. v. M., Cens. of Austr. pl. 30 (1882)	—	—	—	—	—	Q.	—	—	M.fr.VIII,163.
B. divaricata; F. v. M., Cens. of Austr. pl. 30 (1882) from R. Brown (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,200	M.fr.VII,14; X,92; XII,13.
B. bicuspis; F. v. M., Cens. of Austr. pl. 30 (1882)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,200	M.fr.VII,13.
B. glabra; F. v. M., Cens. of Austr. pl. 30 (1882)	—	—	—	—	—	—	N.A.	B.fl.V,200	M.fr.I,139.
B. echinopsila; F. v. M., Cens. of Austr. pl. 30 (1882)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,201	M.fr.I,139; VII,14; X,92; XII,13.
B. enchylaenoides; F. v. M., Cens. of Austr. pl. 30 (1882)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,182	M.fr.X,92; XII,13.

B. micrantha; F. v. M., Cens. of Austr. pl. 30 (1882) from Bentham (1870) ... W.A. — — — — — — B.fl.V,181
B. brevicuspis; F. v. M., Cens. of Austr. pl. 30 (1882) ... ... — — — — N.S.W. Q. — B.fl.V,198 M.fr.IV,150; VII,14; VIII,38.
B. salsuginosa; F. v. M., Cens. of Austr. pl. 30 (1882) ... ... — S.A. — V. N.S.W. — — B.fl.V,197 M.fr.VII,12; X,92; XI,27; XII,13.

THRELKELDIA; R. Brown, prodr. 409 (1810).

T. proceriflora; F. v. M., fragm. VIII, 38 (1873) ... ... — — — — N.S.W. Q. — — M.fr.VIII,38.
T. diffusa; R. Brown, prodr. 410 (1810) ... ... W.A. S.A. T. V. N.S.W — — B.fl.V,197 M.fr.VII,12.

ENCHYLAENA; R. Brown, prodr. 408 (1810).

E. tomentosa; R. Brown, prodr. 408 (1810) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,181 M.fr.VII,12; IX,75; XI,27; XII,15.

SALICORNIA; Tournefort, coroll. 51 (1703). (Halocnemum, Arthrocnemum, Tecticornia, Pachycornia) from Gerard (1597).

S. robusta; F. v. M., fragm. VI, 251 (1868) ... ... W.A. S.A. — V. N.S.W. — N.A. B.fl.V,202 M.fr.I,139; VII,14; XII,18.
S. arbuscula; R. Brown, prodr. 411 (1810) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,203 M.fr.VII,15; XII,18.
S. cinerea; F. v. M., fragm. VI, 251 (1869) ... ... — — — — — Q. N.A. B.fl.V,203 M.fr.I,140; VII,15; XII,18.
S. leiostachya; Bentham, Fl. Austr. V, 203 (1870) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.V,203
S. bidens; Bentham, Fl. Austr. V, 204 (1870) ... ... W.A. — — — — — — B.fl.V,204
S. tenuis; Bentham, Fl. Austr. V, 204 (1870) ... ... — S.A. — — N.S.W. Q. — B.fl.V,204
S. Australis; Solander in G. Forster, prodr. 88 (1786) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,205 M.fr.VII,15; IX,75; XI,20,27; XII,18. N.Z.

SUAEDA; Forskael, Fl. Aegypt. Arab. 69 (1779). (Chenopodium parlty, Schoberia, Chenopodina.)

S. maritima; Dumortier, Fl. Belg. 22 (1827) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.V,206 M.fr.XI,20. ... E., As., Afr., Am., P., N.Z.

SALSOLA; Linné, gen. pl. 67 (1737), from Cisalpini. [Am., P.

S. Kali; Linné, spec. plant. 222 (1753) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.V,207 M.fr.IX,75; XI,20,27; XII,18. E., As., Afr.,

FICOIDEAE.

A. L. de Jussieu, Gen. plant. 315 (1789).—Genera 9, Species 28.

MESEMBRIANTHEMUM; Breyne, prodr. fasc. rar. pl. II, 67 (1689).

M. aequilaterale; Haworth, misc. nat. 77 (1803) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.V,158; XI,20,27. ... Am.
M. Australe; Solander in G. Forster, prodr. 90 (1786) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,324 M.fr.V,158; X,83; XI,20,27. N.Z.

TETRAGONIA; Linné, syst. nat. 9 (1735); gen. pl. 144 (1737). (Tetragonella.)

T. expansa; Murray in Comm. Goetting. VI, 13, t. 5 (1783) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.III,325 M.fr.VII,129; X,84; XI,27. As.,Am.,P.,N.Z.
T. diptera; F. v. M., fragm. XI, 8 (1878) ... ... W.A. — — — — — N.A. — M.fr.XI,8.
T. implexicoma; J. Hooker, Fl. Tasman. I, 148 (1859) ... ... W.A. S.A. T. V. N.S.W. — N.A. B.fl.III,326 M.fr.X,84; XI,8 ... ... N.Z.

AIZOON; Linné, gen. pl. 161 (1737). (Aizoum, Gunniopsis.)

A. quadrifidum; F. v. M., fragm. II, 148 (1861) ... ... W.A. S.A. — — N.S.W. Q. — B.fl.III,327 M.fr.VII,129; X,83.
A. zygophylloides; F. v. M., fragm. VII, 129 (1871) ... ... W.A. S.A. — — N.S.W. Q. — — M.fr.X,83.

GUNNIA; F. v. M., Rep. on pl. of Babb. Exped. 9 (1858).

G. septifraga; F. v. M., Rep. of Babb. Exped. 9 (1858) ... ... — S.A. — — N.S.W. Q. — B.fl.III,327
G. Drummondii; Bentham, Fl. Austr. III, 327 (1866) ... ... W.A. — — — — — — B.fl.III,327

SESUVIUM; Linné, syst. ed. decim. 1058 (1759). [Afr., Am

S. portulacastrum; Linné, syst. ed. dec. 1058 (1759) ... ... — — — — N.S.W. Q. N.A. B.fl.III,328 M.fr.III,164; VII,129; IX,77; X,83. As.,

ZALEYA; N. L. Burmann, Fl. Indic. 110, t. 31 (1768).

Z. decandra; N. L. Burmann, Fl. Ind. 110, t. 31 (1768) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.III,329 M.fr.I,172; III,164; X,83. As.

TRIANTHEMA; Sauvage, Méth. fol. 127 (1751). (Ancistrostigma.)

T. turgidifolia; F. v. M., fragm. X, 83 (1876) ... ... — S.A. — — — Q. N.A. — M.fr.X,83.
T. crystallina; Vahl, symbol. I, 32 (1790) .. .. W.A. S.A. — — N.S.W. Q. N.A. B.fl.III,330 M.fr.I,171-2. As., Afr.
T. glaucifolia; F. v. M., fragm. I, 172 (1859).. ... ... W.A. — — — — Q. — — M.fr.I,172.

T. pilosa; F. v. M., fragm. I, 174 (1859)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.III,330	M.fr.X,83.				
T. oxycalyptra; F. v. M., fragm. I, 173 (1859)	—	—	—	—	—	—	N.A.	B.fl.III,330	M.fr.I,173.				
T. rhynchocalyptra; F. v. M., fragm. I, 174 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.III,330	M.fr.I,174.				
T. glossostigma; F. v. M. in Wing's S. Sc. Rec. III, 282 (1884)	W.A.	—	—	—	—	—	—	—					
T. cypseloides; Bentham, Fl. Austr. III, 331 (1866)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,331	M.fr.X,83.				
MACARTHURIA; Huegel, enum. pl. Nov. Holl. Austr. occ. 11 (1837).													
M. apetala; Harvey in Hook. Kew. Misc. VII, 55 (1855)	W.A.	—	—	—	—	—	N.A.	B.fl.III,332	M.fr.XI,20.				
M. Australis; Huegel, enum. pl. 11 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,332	M.fr.II,11; V,28.				
M. Neo-Cambrica; F. v. M., fragm. V, 28 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,332	M.fr.V,28.				
MOLLUGO; Linné, gen. plant. 336 (1737). (Glinus, Trigastrotheca.)													
M. hirta; Thunberg, prodr. pl. cap. I, 24 (1794)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,333	...	...	E., As., Afr., Am.		
M. orygioides; F. v. M. in Benth. Fl. Austr. III, 333 (1866)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,333					
M. Spergula; Linné, spec. pl. ed. alt. 131 (1762)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,334	M.fr.II,148	...	As., Afr.		
M. stricta; Linné, spec. pl. ed. alt. 31 (1762)	—	—	—	—	—	Q.	—	—	...	...	As., P.		
M. trigastrotheca; F. v. M., pl. of Vict. I, 201 (1860)	W.A.	—	—	—	—	—	N.A.	B.fl.III,334					
M. Cerviana; Séringe in De Candolle, prodr. I, 392 (1824)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,334	M.fr.II,148	...	E., As., Afr.		

POLYGONACEAE.

A. L. de Jussieu, Gen. pl. 82 (1789), from B. de Jussieu (1759).—Genera 4, Species 25.

EMEX; Necker, elem. bot. II, 214 (1790). (Perhaps immigrated.)													
E. Australis; Steinheil in Ann. des sc. nat. ser. 2, IX, 195, t. 7 (1839)	W.A.	S.A.	—	—	—	—	—	B.fl.V,262	...	...	Afr.		
RUMEX; Linné, gen. pl. 105 (1737), from Plinius.													
R. Brownii; Campdera, Monogr. Rum. 81 (1819)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,263					
R. flexuosus; Solander in G. Forst., fl. ins. Austr. prodr. 90 (1786)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,264	...	...	N.Z.		
R. crystallinus; Lange, ind. sem. hort. Hafn. 28 (1861)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,265	M.fr.IV,48.				
R. bidens; R. Brown, prodr. 421 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,265					
POLYGONUM; l'Obel, stirp. hist. 228 (1576), from Dioscorides and Plinius.													
P. plebejum; R. Brown, prodr. 420 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,267	...	...	E., As., Afr.		
P. strigosum; R. Brown, prodr. 420 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,268	...	...	As., N.Z.		
P. prostratum; R. Brown, prodr. 419 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,268					
P. hydropiper; Linné, spec. plant. 361 (1753)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,269	...	...	E., As., Afr., Am.		
P. minus; Hudson, Fl. Anglica I, 148 (1762)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,269	...	...	E., As., Afr.		
P. subsessile; R. Brown, prodr. 419 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,269	...	...	P.		
P. barbatum; Linné, spec. plant. 362 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,270	...	...	As., Afr.		
P. articulatum; R. Brown, prodr. 420 (1810)	—	—	—	—	—	Q.	—	B.fl.V,270					
P. lapathifolium; Linné, spec. plant. 360 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,270	...	...	E., As., Afr., Am.		
P. orientale; Linné, spec. plant. 362 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,271	...	...	As., P.		
P. attenuatum; R. Brown, prodr. 420 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,272	...	...	Am.		
MUEHLENBECKIA; Meissner, gen. pl. vasc. 316 (1840). (Sarcogonum, Coccoloba partly.)													
M. adpressa; Meissner, gen. pl. comm. 227 (1843)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,273	M.fr.XI,20.	...	N.Z., Am.		
M. Australis; Meissner, gen. pl. comm. 227 (1843)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,169.				
M. gracillima; Meissner in De Candolle, prodr. XIV, 145 (1856)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,274					
M. rhyticarya; F. v. M., fragm. V, 92 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,274					
M. axillaris; J. Hooker in Lond. Journ. VI, 278 (1847)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,275	M.fr.IX,77	...	N.Z.		
M. polybotrya; Meissner in Lehm. pl. Preiss. I, 623 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,475	M.fr.IV,130.				

M. polygonoides; F. v. M., fragm. V, 73 (1865)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,275	M.fr.V,73.			
M. stenophylla; F. v. M., fragm. I, 138 (1859)	—	—	—	V.	N.S.W.	—	—	B.fl.V,275	M.fr.I,138.			
M. Cunninghamii; F. v. M., fragm. V, 91 (1865)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,275	M.fr.IV,131.			

PHYTOLACCEAE.

R. Brown in Tuck. Exped. Cong. App. V, 454 (1818).—Genera 6, Species 11.

MONOCOCCUS; F. v. M., fragm. I, 47 (1858).

M. echinophorus; F. v. M., fragm. I, 47 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,144	M.fr.I,47	...	P.	

DIDYMOTHECA; J. Hooker in Lond. Journ. Bot. VI, 278 (1847). (Cyclotheca).

D. thesioides; J. Hooker in Lond. Journ. V, 279 (1847)	W.A.	S.A.	T.	—	—	—	—	B.fl.V,145				
D. pleiococca; F. v. M., fragm. I, 202 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.V,146	M.fr.I,202.			

GYROSTEMON; Desfontaines in Mém. du Mus. VI, 16, t. 6 (1820).

G. brachystigma; F. v. M. in Benth. Fl. Austr. V, 146 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,146				
G. ramulosus; Desfontaines in Mém. du Mus. V, 17, t. 6 (1820)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,147				

CODONOCARPUS; Cunningham in Hook. Bot. Misc. I, 244 (1830). (Hymenotheca.)

C. pyramidalis; F. v. M., plants of Vict. I, 201 (1862)	—	S.A.	—	—	—	—	—	B.fl.V,148	M.fr,I,202.			
C. Australis; Cunningham in Hook. bot. misc. I, 244 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,148				
C. cotinifolius; F. v. M., pl. of Vict. I, 200 (1862)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,149	M.fr.II,181; X,53.			

TERSONIA; Moquin in De Candolle, prodr. XIII, pars sec. 40 (1849).

T. brevipes; Moquin in De Candolle, prodr. XIII, pars II, 40 (1849)	W.A.	—	—	—	—	—	—	B.fl.V,149				
T. subvolubilis; Bentham, Fl. Austr. V, 150 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,150				

CYPSELOCARPUS; F. v. M., fragm. VIII, 36 (1873).

C. haloragoides; F. v. M., fragm. VIII, 36 (1873)	W.A.	—	—	—	—	—	—	B.fl.V,198	M.fr.VIII,36.			

NYCTAGINEAE.

A. L. de Jussieu, Gen. plant. 90 (1789).—Genera 2, Species 6.

BOERHAAVIA; Vaillant, serm. de struct. fl. 50 (1718).

B. diffusa; Linné, fl. Zeylan. 4 (1747)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,277	M.fr.IX,75,77			As., Afr., P.
B. elegans; Choisy in De Candolle, prodr. XIII, pars II, 453 (1849)	—	—	—	—	—	—	N.A.	—		...	...	As., Afr.
B. plantaginea; Cavanilles, icon. et descr. pl. II, 7, t. 112 (1793)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.V,278		...	...	E., As., Afr.

PISONIA; Plumier, nov. pl. Amer. gen. 7, t. 11 (1703).

P. aculeata; Linné, sp. pl. 1026 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,279	M.fr.VI,197	...	As., Afr., Am., P.	
P. inermis; G. Forster, prodr. 75 (1786)	—	—	—	—	—	Q.	N.A.	B.fl.V,280	M.fr.VI,197	...	P.	
P. Brunoniana; Endlicher, prodr. fl. Norf. 43 (1833)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,280	M.fr.I,20; IV,169; VI,198; IX,77.		N.Z.	

CHORIPETALEAE PERIGYNAE.

F. v. M. in Woolls, pl. of the neighb. of Sydney 18 (1880).

THYMELEAE.

A. L. de Jussieu, Gen. plant. 76 (1789).—Genera 4, Species 75.

PIMELEA; Banks & Solander in Gaert. de fruct. I, 186 (1788). (Thecanthes, Gymnococca, Heterolaena, Calyptrostegia, Macrostegia;—Banksia, Forst., 1776.)

P. punicea; R. Brown, prodr. 359 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,6	M.fr.V,74; VII,3.				
P. concreta; F. v. M., fragm. V, 73 (1865)	—	—	—	—	—	—	N.A.	B.fl.VI,6	M.fr.V,73.				
P. cornucopiae; Vahl, enum. plant. I, 305 (1804)	—	—	—	—	—	Q.	—	B.fl.VI,6	M.fr.VII,3.				
P. sanguinea; F. v. M., fragm. I, 84 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.VI,7	M.fr.VII,3; XI,47.				
P. alpina; F. v. M., second general report, 14 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,7	M.fr.VII,4.				
P. longifolia; Banks & Solander in Rees, Cycl. XXVI (1814)	—	—	—	—	N.S.W.	—	—	B.fl.VI,7	M.fr.VIII,9.	...	N.Z.		
P. cinerea; R. Brown, prodr. 361 (1810)	—	—	T.	—	—	—	—	B.fl.VI,8					
P. Milligani; Meissner in De Candolle, prodr. XIV, 509 (1856)	—	—	T.	—	—	—	—	B.fl.VI,8					
P. spectabilis; Lindley, Bot. Regist. t. 33 (1841)	W.A.	—	—	—	—	—	—	B.fl.VI,9	M.fr.VII,4.				
P. Lehmanniana; Meissner in Lehm. pl. Preiss. I, 603 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,9	M.fr.VII,4.				
P. hispida; R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,10					
P. rosea; R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,10					
P. ferruginea; Labillardière, Nov. Holl. pl. spec. I, 10, t. 5 (1804)	W.A.	—	—	—	—	—	—	B.fl.VI,10	M.fr.VII,4.				
P. brachyphylla; Bentham, Fl. Austr. VI, 11 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,11					
P. sylvestris; R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,11	M.fr.VII,2&5.				
P. brevifolia; R. Brown, prodr. 359 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,12	M.fr.VII,4.				
P. Maxwelli; F. v. M. in Benth. Fl. Austr. VI, 12 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,12					
P. angustifolia; R. Brown, prodr. 360 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,13	M.fr.VII,3.				
P. nervosa; Meissner in Lehm. pl. Preiss. II, 269 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,13	M.fr.VII,3.				
P. sulphurea; Meissner in Mohl & Schlecht Bot. Zeitung, 396 (1848)	W.A.	—	—	—	—	—	—	B.fl.VI,14	M.fr.VII,5.				
P. floribunda; Meissner in De Candolle, prodr. XIV, 505 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,14	M.fr.VII,5.				
P. suaveolens; Meissner in Lehm. pl. Preiss. I, 603 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,14	M.fr.VI,2,4,5.				
P. physodes; Hooker, icon. plant. t. 865 (1852)	W.A.	—	—	—	—	—	—	B.fl.VI,15	M.fr.VII,4.				
P. glauca; R. Brown, prodr. 360 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,15	M.fr.VII,4; XI,123.				
P. colorans; Cunningham in De Candolle, prodr. XIV, 499 (1856)	—	—	—	—	N.S.W.	—	—	B.fl.VI,16					
P. collina; R. Brown, prodr. 359 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,16					
P. spathulata; Labillardière, Nov. Holl. pl. spec. I, 9, t. 4 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,17					
P. linifolia; Smith, Specim. of Bot. of N. Holl. 31, t. 11 (1793)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,17	M.fr.VII,3.				
P. ligustrina; Labillardière, Nov. Holl. pl. spec. I, 9, t. 3 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,18	M.fr.I,87; IV,163; VII,5.				
P. humilis; R. Brown, prodr. 361 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,19	M.fr.VII,4; XI,47.				
P. sericea; R. Brown, prodr. 361 (1810)	—	—	T.	—	—	—	—	B.fl.VI,19	M.fr.VII,5.				
P. nivea; Labillardière, Nov. Holl. pl. spec. I, 10, t. 6 (1804)	—	—	T.	—	—	—	—	B.fl.VI,20	M.fr.VII,5.				
P. imbricata; R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,20	M.fr.VII,5.				
P. villifera; Meissner in Lehm. pl. Preiss. II, 271 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,21					
P. drupacea; Labillardière, Nov. Holl. pl. spec. I, 10, t. 7 (1804)	—	—	T.	V.	—	—	—	B.fl.VI,21	M.fr.VII,8.				
P. haematostachya; F. v. M., fragm. I, 84 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,21	M.fr.I.84; VII,8; XI,47.				
P. spicata; R. Brown, prodr. 362 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,22	M.fr.VII,7; VIII,9.				
P. spiculigera; F. v. M., in Bentham Fl. Austr. VI, 23 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,23	M.fr.XI,46.				
P. Forrestiana; F. v. M., fragm. XI, 46 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,46.				
P filiformis; J. Hooker in Lond. Journ. VI, 280 (1847)	—	—	T.	—	—	—	—	B.fl.VI,23	M.fr.VII,7.				
P. latifolia; R. Brown, prodr. 362 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,23	M.fr.IV,49; VII,7.				
P. simplex; F. v. M. in Linnaea XXV, 443 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,23					

P. sericostachya; F. v. M., fragm. IV, 162 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,24	M.fr.XI,47.						
P. trichostachya; Lindley in Mitch. Trop. Austr. 355 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,24	M.fr.VII,7.						
P. leptostachya; Bentham, Fl. Austr. VI, 24 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,24							
P. argentea; R. Brown, prodr. 362 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,25	M.fr.VII,7.						
P. clavata; Labillardière, Nov. Holl. pl. spec. I, 11 (1804)...	W.A.	—	—	—	—	—	—	B.fl.VI,25	M.fr.VI,160; VII,5.						
P. axiflora; F. v. M., First general Report 17 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,26	M.fr.VII,7.						
P. leptospermoides; F. v. M., fragm. VII, 2 (1870)...	—	—	—	—	—	Q.	—	B.fl.VI,27	M.fr.VII,2.						
P. microcephala; R. Brown, prodr. 361 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,27	M.fr.XI,27,47.						
P. pauciflora; R. Brown, prodr. 360 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,27	M.fr.VII,8; XI,47.						
P. elachantha; F. v. M., First general Report 17 (1853)	—	S.A.	—	V.	—	—	—	B.fl.VI,28	M.fr.VII,6.						
P. pygmaea; F. v. M., in Schlecht. Linnaea XXVI, 346 (1853)	—	—	T.	—	—	—	—	B.fl.VI,28							
P. serpyllifolia; R. Brown, prodr. 360 (1810)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,29	M.fr.VII,6.						
P. flava; R. Brown, prodr. 361 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,29	M.fr.VII,6.						
P. petrophila; F. v. M., in Schlecht. Linnaea XXV, 442 (1852)	—	S.A.	—	—	N.S.W.	—	—	B.fl.VI,30							
P. Bowmanni; F. v. M. in Benth. Fl. Austr. VI, 30 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,30	M.fr.VII,6.						
P. ammocharis; F. v. M. in Hook. Kew Misc. IX, 24 (1857)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.VI,30	M.fr.VII,5; XI,47.						
P. penicillaris; F. v. M. in Melb. Chemist Oct. (1883)	—	—	—	—	N.S.W.	Q.	—	—							
P. curviflora; R. Brown, prodr. 362 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,31	M.fr.VII,6; VIII,9.						
P. hirsuta; Meissner in De Candolle, prodr. XIV, 513 (1856)	—	—	—	—	N.S.W.	—	—	B.fl.VI,32							
P. altior; F. v. M., fragm. I, 84 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,32	M.fr.VII,7.						
P. octophylla; R. Brown, prodr. 361 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,32	M.fr.VII,6.						
P. petraea; Meissner in Schlecht. Linnaea XXVI, 347 (1853)	—	S.A.	—	—	N.S.W.	—	—	B.fl.VI,33							
P. phylicoides; Meissner in Lehm. pl. Preiss. II, 271 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,33	M.fr.VII,6.						
P. Eyrei; F. v. M., fragm. V, 109 (1865)	W.A.	—	—	—	—	—	—	B.fl.VI,34	M.fr.V,109.						
P. longiflora; R. Brown, prodr. 361 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,34	M.fr.VII,5.						
P. stricta; Meissner in Schlecht. Linnaea XXVI, 348 (1853)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,34	M.fr.VII,4.						
P. Preissi; Meissner in Lehm. pl. Preiss. I, 601 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,35	M.fr.VII,3.						
P. Holroydii; F. v. M., fragm. VI, 159, t. 59 (1868)	—	—	—	—	—	—	N.A.	B.fl.VI,35	M.fr.250.						
DRAPETES; Lamarck, Journ. d'hist. nat. I, 119, t. 10 (1792). (Kelleria, Daphnobryon.)															
D. Tasmanica; J. Hooker in Kew Misc. V, 299, t. 7 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,36	M.fr.VII,1.						
WICKSTROEMIA; Endlicher, prodr. fl. insul. Norfolk. 47 (1833). (Subgenus Stellerae (1747) Daphne partly.)															
W. Indica; C. A. Meyer in Bull. de l'Acad. de Petersb. I, 357 (1843)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,37	M.fr.VII,1	...	As., P.				
PHALERIA; Jack in Hook. Comp. to Bot. Mag. I, 156 (1835). (Drymispermum.)															
P. Blumei; Bentham, Fl. Austr. VI, 38 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,38	...	...	As.				
P. Neumanni; F. v. M., fragm. VIII, 9 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,38	M.fr.V,26; VII,1.						
P. Clerodendron; F. v. M., fragm. VIII, 9 (1873)	—	—	—	—	—	Q.	—	B.fl.VI,38	M.fr.VII,1.						

LEGUMINOSAE.

HALLER, Enum. stirp. Helv. II, 565 (1742).—Genera 95, Species 1065.

JANSONIA; Kippist in Gard. Chron. 19 (1847). (Cryptosema.)															
J. formosa; Kippist in Transact. Linn. Soc. XX, 384, t. 16 (1851)	W.A.	—	—	—	—	—	—	B.fl.II,8							
BRACHYSEMA; R. Brown in Ait. hort. Kew. sec. edit. III, 10 (1811). (Leptosema, Kaleniczenkia, Burgesia.)															
B. praemorsum; Meissner in Lehm. pl. Preiss. I, 25 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,10							
B. lanceolatum; Meissner in Lehm. pl. Preiss. I, 24 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,10							
B. latifolium; R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,10							
B. undulatum; Ker in Bot. Regist. t. 642 (1822)	W.A.	—	—	—	—	—	—	B.fl.II,11	M.fr.IV,11.						

B. subcordatum ; Bentham, Fl. Austr. II, 11 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,11	
B. bracteolosum ; F. v. M., fragm. IV, 10 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,11	M.fr.IV,10.
B. bossiaeoides ; Bentham, Fl. Austr. II, 12 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,12	
B. oxylobioides ; Bentham, Fl. Austr. II, 12 (1864)...	—	—	—	—	—	Q.	N.A.	B.fl.II,12	
B. uniflorum ; R. Brown in Benth., Fl. Austr. II, 12 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,12	
B. aphyllum ; Hooker, Bot. Mag. t. 4481 (1849)	W.A.	—	—	—	—	—	—	B.fl.II,12	M.fr.I,222; II,182.
B. macrocarpum ; Bentham, Fl. Austr. II, 13 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,13	
B. tomentosum ; Bentham, Fl. Austr. II, 13 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,13	
B. Chambersii ; F. v. M. in Bentham, Fl. Austr. II, 13 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.II,13	
B. daviesioides ; Bentham, Fl. Austr. II, 13 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,13	

OXYLOBIUM ; Andrews, Bot. Reposit. t. 492 (1809). (Chorizematis subgenus, Callistachys, Podolobium.)

O. Callistachys ; Bentham, Fl. Austr. II, 16 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,16	M.fr.IV,18.
O. ellipticum ; R. Brown in Ait. hort. Kew. sec. ed. III, 10 (1811)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,16	
O. alpestre ; F. v. M. in Transact. phil. Soc. Vict. I, 38 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.II,17	
O. lineare ; Bentham, Fl. Austr. II, 17 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,17	M.fr.IV,17.
O. carinatum ; Bentham, Fl. Austr. II, 18 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. spathulatum ; Bentham, Fl. Austr. II, 18 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. obtusifolium ; Sweet, Fl. Austr. t. 5 (1827)	W.A.	—	—	—	—	—	—	B.fl.II,18	
O. cordifolium ; Andrews, Bot. Reposit. t. 492 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.II,19	M.fr.IV,17.
O. microphyllum ; Bentham, Fl. Austr. II, 19 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,19	
O. Pulteneae ; De Candolle, prodr. II, 104 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,19	M.fr.IV,19.
O. hamulosum ; Bentham in A. Gray, Bot. Wilk. Exped. t. 379	—	—	—	—	N.S.W.	—	—	B.fl.II,20	
O. scandens ; Bentham in Ann. Wien. Mus. II, 70 (1839)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,20	
O. procumbens ; F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.II,20	
O. tricuspidatum ; Meissner in Lehm. pl. Preiss. I. 30 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,21	
O. spectabile ; Endlicher, nov. stirp. dec. 2 (1839)	W.A.	—	—	—	—	—	—	B.fl.II.21	
O. atropurpureum ; Turczaninow in Bull. Soc. Mosc. XXVI, 250 (1853)... ...	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. retusum ; R. Brown in Bot. Regist. t. 913 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. virgatum ; Bentham, Fl. Austr. II, 22 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,22	
O. reticulatum ; Meissner in Lehm. in Preiss. I, 29 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. capitatum ; Bentham in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. cuneatum ; Bentham in Bot. Regist. XXV, App. XII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,23	
O. acutum ; Bentham, Fl. Austr. II, 24 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,24	
O. parviflorum ; Bentham in Bot. Regist. XXV, App. XII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,24	
O. heterophyllum ; Bentham, Fl. Austr. II, 25 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,25	
O. aciculiferum ; F. v. M., fragm. I, 75 implied (1859)	—	—	—	—	—	Q.	—	B.fl.II,25	M.fr.I,75.
O. trilobatum ; F. v. M., annual report 23 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,25	M.fr.IV,19.

CHORIZEMA ; Labillardière, Voy. I, 405 (1799). (Orthotropis.)

C. Dicksonii ; Graham in Maund's Botanist, t. 106 (1841)	W.A.	—	—	—	—	—	—	B.fl.II,27	
C. nervosum ; Th. Moore in Gard. Companion (1852)	W.A.	—	—	—	—	—	—	B.fl.II,27	
C. varium ; Bentham in Bot. Regist. XXV, t. 49 (1839)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. cordatum ; Lindley, Bot. Regist. t. 10 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. ilicifolium ; Labillardière, Voy. I, 405, t. 21 (1799)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. rhombeum ; R. Brown in Ait. hort. Kew. sec. ed. III, 9 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,28	
C. diversifolium ; A. de Candolle, Pl. rar. jard. Genèv. 44, t. 8 (1836)	W.A.	—	—	—	—	—	—	B.fl.II,29	
C. angustifolium ; Bentham in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,29	
C. reticulatum ; Meissner in Lehm. pl. Preiss. I, 34 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. trigonum ; Turczaninow in Bull. de la Soc. Mosc. XXVI, 254 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,30	
C. humile ; Turczaninow in Bull de la Soc. Mosc. XXVI, 254 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,30	

C. parviflorum ; Bentham in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,30	
C. cytisoides ; Turczaninow in Bull. Soc. Mosc. XXVI, 256 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,31	
C. Henchmanni ; R. Brown in Bot. Regist. t. 986 (1826)	W.A.	—	—	—	—	—	—	B.fl.II,31	
C. ericifolium ; Meissner in Lehm. pl. Preiss. II, 209 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,32	
GASTROLOBIUM ; R. Brown in Ait. hort. Kew. sec. edit. III, 16 (1811). (Chorizematis subgenus.)									
G. pyramidale ; Th. Moore in Gard. Comp. I, 81 (1852)	W.A.	—	—	—	—	—	—	B.fl II,98	
G. Lehmanni ; Meissner in Lehm. pl. Preiss. I, 70 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,98	M.fr.,X,35.
G. pulchellum ; Turczaninow in Bull. de la Soc. Mosc. XXVI, 274 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,98	
G. stipulare; Meissner in Lehm. pl. Preiss. II, 218 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,99	
G. Brownii ; Meissner in Lehm. pl. Preiss. I, 71 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,99	
G. reticulatum ; Bentham, Fl. Austr. II, 99 (1864)	W.A.	—	—	—	—	—	—	B.fl.II 99	M.fr.X,35.
G. truncatum ; Bentham, Fl. Austr. II, 99 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,99	
G. spathulatum ; Bentham in Bot. Regist. XX, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,100	
G. plicatum ; Turczaninow in Bull. Soc. Mosc. XXVI, 274 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,100	
G. tricuspidatum ; Meissner in Lehm. pl. Preiss. I, 66 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,100	
G. obovatum ; Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,101	
G. epacridioides ; Meissner in Lehm. Preiss. I, 72 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. trilobum ; Bentham in Bot. Regist. XXV, App XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,101	M.fr.X,35.
G. ilicifolium ; Meissner in Lehm. pl. Preiss I, 67 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,101	
G. villosum ; Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. polystachyum ; Meissner in Lehm. pl. Preiss. II, 217 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. ovalifolium ; Henfrey in Gard. Comp. I, 41 (1852)	W.A.	—	—	—	—	—	—	B.fl.II,102	
G. grandiflorum ; F. v. M., fragm. III, 17 (1862)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,103	M.fr.IV,174 ; X,35.
G. pycnostachyum ; Bentham, Fl. Austr. II, 103 (1864)	W.A.	—	—	—	—	—	—	B.fl.II.103	
G. spinosum ; Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,103	M.fr.X,35.
G. rotundifolium ; Meissner in Lehm. pl. Preiss. II, 216 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,104	
G. microcarpum ; Meissner in Lehm. pl. Preiss. I, 70 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,104	
G. oxylobioides ; Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,104	
G. calycinum ; Bentham in Bot. Regist. XXV, App. XIII (1839)...	W.A.	—	—	—	—	—	—	B.fl.II,104	M.fr.X,35.
G. Callistachys ; Meissner in Lehm. pl. Preiss. II, 216 (1846)	W.A.	—	—	—	—	—	—	B.fl.II,105	
G. stenophyllum ; Turczaninow in Bull. Soc. Mosc. XXVI, 275 (1853)	W.A.	—	—	—	—	—	—	B.fl.II.105	
G. crassifolium ; Bentham, Fl. Austr. II, 105 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,105	
G. parvifolium ; Bentham, Fl. Austr. II, 106 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,106	
G. hamulosum ; Meissner in Lehm. pl. Preiss. II, 218 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,106	
G. velutinum ; Lindley & Paxton, Flower-Gard. III, 76 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,106	M.fr.X,35.
G. bidens ; Meissner in Regensb. Bot. Zeit. 29 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,106	
G. bilobum ; R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,107	M.fr.X,35.
G. seorsifolium ; F. v. M., fragm. X, 45 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,35.
ISOTROPIS ; Bentham in Hueg. enum. 28 (1837).									
I. striata ; Bentham in Hueg. enum. 28 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,39	M.fr.III,16.
I. Drummondii ; Meissner in Lehm. pl. Preiss, I, 31 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,39	
I. juncea ; Turczaninow in Bull. Soc. Mosc. XXVI, 251 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,39	M.fr.X,51.
I. canescens ; F. v. M., fragm. X, 51 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,51.
I. atropurpurea ; F. v. M., fragm. III, 16 (1862)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,40	M.fr.X,51.
I. filicaulis ; Bentham in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	—	Q.	—	B.fl.II,40	M.fr.IV,20.
I. Forrestii ; F. v. M. in Wing's S. Sc. Rec. II, 252 (1882)	W.A.	—	—	—	—	—	—	—	
I. parviflora ; Bentham in Ann. des Wien. Mus. II, 71 (1838)	—	—	—	—	—	—	N.A.	B.fl.II,40	
I. Wheeleri ; F. v. M. in Benth. Fl. Austr. II, 40 (1864)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,40	
I. Winneckiana ; F. v. M. in Austr. Chemist (1884)	—	S.A.	—	—	—	—	—	—	

MIRBELIA; Smith in Koenig & Sims, Ann. of Bot. I, 511 (1805). (Dichosema, Oxycladium.)

M. dilatata; R. Brown in Ait. hort. Kew. sec. ed. III, 21 (1811) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,33	
M. racemosa; Turczaninow in Bull Soc. Mosc. XXVI, 282 (1853)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,33	M.fr.IV,18.
M. grandiflora; Aiton in Bot. Mag. t. 2771 (1827)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,34	
M. subcordata; Turczaninow in Bull. Soc. Mosc. XXVI, 282 (1853)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,34	
M. ovata; Meissner in Lehm. pl. Preiss. I, 77 (1844)	..	...	W.A.	—	—	—	—	—	—	B.fl.II,34	M.fr.II,154.
M. oxylobioides; F. v. M., fragm. II, 154 (1861)	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.II,34	M.fr.III,167; IV,12.
M. reticulata; Smith in Koen. & Sims, Ann. of Bot. I, 511 (1805)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,35	M.fr.IV,12.
M. aotoides; F. v. M. in Transact. phil. Inst. Vict. III, 53 (1859)	...	..	—	—	—	—	N.S.W.	Q.	—	B.fl.II,35	M.fr.IV,12.
M. pungens; Cunningham in G. Don, gen. syst. II, 126 (1832) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,35	
M. speciosa; Sieber in De Candolle, prodr. II, 115 (1824)	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,36	M.fr.IV,12.
M. floribunda; Bentham in Bot. Regist. XXV, App. XII (1839)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,36	
M. spinosa; Bentham, Fl. Austr. II, 36 (1864)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,36	
M. multicaulis; Bentham, Fl. Austr. II, 37 (1864)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,37	
M. microphylla; Bentham, Fl. Austr. II, 37 (1864)...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,37	
M. aphylla; F. v. M., fragm. IV, 11 (1863)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,37	M.fr.IV,11.
M. oxyclada; F. v. M., fragm. IV, 12 (1864)...	...	...	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,38	M.fr.I,168.

GOMPHOLOBIUM; Smith in Transact. Linn. Soc. IV, 220 (1798).

G. amplexicaule; Meissner in Lehm. pl. Preiss. I, 36 (1844)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,42	
G. latifolium; Smith in Koenig & Sims, Ann. of Bot. I, 505 (1805)	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,42	
G. pedunculare; Loddiges, Bot. Cabinet, t. 1639 (1830)	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.II,43	
G. polymorphum; R. Brown in Ait. hort. Kew sec. ed. III, 11 (1811)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,43	
G. obcordatum; Turczaninow in Bull. Soc. Mosc. XXVI, 258 (1853)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,44	
G. marginatum; R. Brown in Ait. hort. Kew. sec. ed. III, 11 (1811)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,44	
G. grandiflorum; Smith, Exot. Bot. t. 5 (1804)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,44	
G. virgatum; Sieber in De Candolle, prodr. II, 105 (1824)... ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,45	
G. minus; Smith in Transact. Linn. Soc. IX, 251 (1808)	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,45	
G. uncinatum; Cunningham in Annal. des Wien. Mus. II, 72 (1838)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,46	
G. Eatoniae; F. v. M. in Vict. Nat. VI (1889)	...	...	W.A.	—	—	—	—	—	—	—	
G. Baxteri; Bentham, Fl. Austr. II, 46 (1864)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,46	
G. aristatum; Bentham in Ann. des Wien. Mus. II, 72 (1838) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,46	
G. burtonioides; Meissner in Lehm. pl. Preiss. I, 37 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,46	
G. capitatum; Cunningham in Bot. Regist. t. 1563 (1839)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,47	
G. tomentosum; Labillardière, Nov. Holl. pl. sp. 1, 106, t. 134 (1804)	...	..	W.A.	—	—	—	—	—	—	B.fl.II,47	
G. Preissii; Meissner in Lehm. pl. Preiss. I, 40 (1844)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,47	
G. viscidulum; Meissner in Lehm. pl. Preiss. I, 39 (1844)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,48	
G. glabratum; De Candolle, prodr. II, 106 (1824)	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,48	
G. nitidum; Solander in Benth. Fl. Austr. II, 48 (1864)	...	...	—	—	—	—	—	Q.	—	B.fl.II,48	
G. pinnatum; Smith in Transact. Linn. Soc. IX, 251 (1808) ...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,48	
G. Shuttleworthii; Meissner in Lehm. pl. Preiss. I, 39 (1844) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,49	
G. venustum; R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,49	
G. Knightianum; Lindley, Bot. Regist. t. 1468 (1837)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,49	

BURTONIA; R. Brown in Ait. hort. Kew. ed. sec. III, 13 (1811). (Gompholobii subgenus.)

B. subulata; Bentham, Fl. Austr. II, 50 (1864)	...	...	—	—	—	—	—	—	N.A.	B.fl.II,50	M.fr.III,30; X,35.
B. foliolosa; Bentham, Fl. Austr. II, 50 (1864)	...	..	—	—	—	—	—	Q.	—	B.fl.II,50	
B. polyzyga; Bentham, Fl. Austr. II, 51 (1864)	...	...	—	S.A.	—	—	—	—	N.A.	B.fl.II,51	M.fr.III,29; X,34.
B. villosa; Meissner in Lehm. pl. Preiss. I, 41 (1844)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,51	M.fr.X,35.
B. Hendersonii; Bentham, Fl. Austr. II, 51 (1864)	...	...	W.A.	—	—	—	—	—	—	B.fl.II,51	
B. gompholobioides; F. v. M., fragm. X, 34 (1876)	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,34.

B. scabra; R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,51	M.fr.X,35.
B. conferta; De Candolle, prodr. II, 106 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,52	M.fr.X,35.
JACKSONIA; R. Brown in Ait. hort. Kew. sec. ed. III, 12 (1811). (Piptomeris.)									
J. dilatata; Bentham in Ann. des Wien. Mus. II, 74 (1838)	--	—	—	—	—	—	N.A.	B.fl.II,54	M.fr.X,37.
J. densiflora; Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,54	M.fr.X,38.
J. carduacea; Meissner in Bot. Zeit. 25 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,55	
J. floribunda; Endlicher in Ann. des Wien. Mus. II, 197 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,55	
J. odontoclada; F. v. M. in Benth. Fl. Austr. II, 55 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,55	M.fr.X,37.
J. ramosissima; Bentham in Mitch. Trop. Austr. 258 (1848)	—	—	—	—	—	Q.	—	B.fl.II,56	M.fr.X,37.
J. foliosa; Turczaninow in Bull. de la Soc. Imp. de Mosc. XXVI, 260 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,56	
J. spinosa; R. Brown in Ait. hort. Kew. sec. ed. III, 13 (1811)	W.A.	—	—	—	--	—	—	B.fl.II,56	M.fr.X,38&39.
J. stricta; Meissner in Regensb. Bot. Zeit. 27 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,56	
J. hakeoides; Meissner in Lehm. pl. Preiss. I, 45 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,57	M.fr.X,39.
J. furcellata; De Candolle, prodr. II, 107 (1824)	W.A.	—	—	—	—	—	—	B.fl.II,57	M.fr.X,39.
J. horrida; De Candolle, prodr. II, 107 (1824)	W.A.	—	—	—	—	—	—	B.fl.II,57	
J. sericea; Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,58	
J. Sternbergiana; Huegel, Bot. Archiv. t. 3 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,58	M.fr.X,39.
J. vernicosa; F. v. M. in Benth. Fl. Austr. II, 58 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,58	M.fr.X,37.
J. rhadinoclona; F. v. M., fragm. X, 37 (1876)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,37.
J. Forrestii; F. v. M. in proceed. L. S. of N.S.W. sec. ser. II, 194 (1887) ...	W.A.	—	—	—	—	—	—	—	
J. thesioides; Cunningham in den Ann. des Wien. Mus. II, 74 (1838)	—	—	—	—	—	Q.	N.A.	B.fl.II,59	M.fr.IV,161; X,37.
J. compressa; Turczaninow in Bull. de la Soc. de Mosc. XXVI, 260 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.II,59	M.fr.X,38.
J. Clarkei; F. v. M. in proceed. L. S. of N.S.W. sec. ser. II, 193 (1887) ...	—	--	—	V.	N.S.W.	--	—	—	
J. scoparia; R. Brown in Ait. hort. Kew. sec. ed. III, 13 (1811)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,59	M.fr.X,37.
J. cupulifera; Meissner in der Bot. Zeit. 27 (1855)	W.A.	—	—	—	—	—	—	—	M.fr.X,36.
J. restioides; Meissner in Lehm. pl. Preiss. I, 46 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. velutina; Bentham, Fl. Austr. II, 60 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. Lehmanni; Meissner in Lehm. pl. Preiss. I, 46 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. racemosa; Meissner in Lehm. pl. Preiss. II, 212 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,60	
J. umbellata; Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. nematoclada; F. v. M., fragm. X, 50 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,50.
J. rhadinoclada; F. v. M., fragm. X, 38 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,38.
J. capitata; Meissner in Lehm. pl. Preiss. I, 45 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. alata; Bentham in Hueg. enum. 30 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,61	M.fr.X,38.
J. angulata; Bentham, Fl. Austr. II, 62 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,62	
J. Stackhousii; F. v. M. in proc. Linn. Soc. N.S.W., VI, 791 (1881)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,138.
J. pteroclada; F. v. M., fragm. X, 37 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,37.
J. macrocalyx; Meissner in der Bot. Zeit. 26 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,62	
J. purpurascens; F. v. M., fragm. IV, 161 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,161.
J. piptomeris; Bentham, Fl. Austr. II, 62 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,62	
SPHAEROLOBIUM; Smith in Koenig & Sims, Ann. of Bot. I, 509 (1805). (Roea.)									
S. linophyllum; F. v. M., fragm. I, 167 (1859)	W.A.	—	--	—	—	—	—	B.fl.II,64	M.fr.I,167.
S. foliosum; F. v. M., fragm. I, 166 (1859)	W.A.	—	—	—	—	—	—	B.fl.II,64	M.fr.I,166.
S. gracile; Bentham, Fl. Austr. II, 64 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,64	
S. racemulosum; Bentham, Fl. Austr. II, 64 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,65	
S. alatum; Bentham in Hueg. enum. 32 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,65	
S. vimineum; Smith in Koen. & Sims, Ann. of Bot. I, 509 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,65	
S. grandiflorum; R. Brown in Benth. Fl. Austr. II, 66 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. fornicatum; Bentham in Hueg. enum. 32 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,66	

S. medium; R. Brown in Ait. hort. Kew. sec. ed. III, 14 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. scabriusculum; Meissner in Lehm. pl. Preiss. II, 214 (1849)	W.A.	—	—	—	—	—	—	B.fl.II,66	
S. macranthum; Meissner in Lehm. pl. Preiss. II, 213 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,67	
S. daviesioides; Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,67	
VIMINARIA; Smith, Exotic. Bot. 51, t. 27 (1804).									
V. denudata; Smith, Exot. Bot. 51, t. 27 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,68	
DAVIESIA; Smith in Transact. Linn. Soc. IV, 222 (1798).									
D. cordata; Smith in Transact. Linn. Soc. IX, 259 (1808)	W.A.	—	—	—	—	—	—	B.fl.II,72	
D. ovata; Bentham, Fl. Austr. II, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,72	
D. crenulata; Turczaninow in Bull. Soc. de Mosc. XXVI, 265 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,72	M.fr.IV,16.
D. oppositifolia; Endlicher in den Ann. des Wien. Mus. II, 199 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,73	
D. alternifolia; Endlicher in Ann. des Wien. Mus. II, 199 (1838)...	W.A.	—	—	—	—	—	—	B.fl.II,73	M.fr.II,105.
D. elongata; Bentham, Fl. Austr. II, 74 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,74	
D. pedunculata; Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,74	
D. arthropoda; F. v. M., fragm. VIII, 225 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,225.
D. mollis; Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,74	
D. concinna; R. Brown in Benth. Fl. Austr. II, 75 (1864)...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,75	
D. umbellulata; Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,75	
D. Wyattii; Bailey in pap. Garden. Assoc. of S. Austr. (1880)	—	—	—	V.	N.S.W.	Q.	—	—	M.fr.X,138.
D. latifolia; R. Brown in Ait. hort. Kew. sec. ed. III, 20 (1811)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,76	
D. corymbosa; Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,76	
D. horrida; Meissner in Lehm. pl. Preiss. I, 54 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,77	
D. reclinata; Cunningham in Benth. Fl. Austr. II, 77 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,77	
D. obtusifolia; F. v. M., fragm. II, 104 (1861)	W.A.	—	—	—	—	—	—	B.fl.II,77	M.fr.II,104.
D. obovata; Turczaninow in Bull. Soc. Mosc. XXVI, 261 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. longifolia; Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. chordophylla; Meissner in Lehm. plant. Preiss. I, 48 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. nematophylla; F. v. M. in Benth. Fl. Austr. II, 78 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,78	
D. daphnoides; Meissner in Lehm. plant. Preiss. I, 54 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,79	
D. nudiflora; Meissner in Lehm. plant. Preiss. I, 53 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,79	
D. rhombifolia; Meissner in Lehm. plant. Preiss. I, 56 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,79	
D. Drummondii; Meissner in Lehm. plant. Preiss. I, 53 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,80	M.fr.II,106.
D. filipes; Bentham in Mitch. Trop. Austr. 363 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,80	
D. cardiophylla; F. v. M., fragm. II, 105 (1861)	W.A.	—	—	—	—	—	—	B.fl.II,79	M.fr.II,105.
D. squarrosa; Smith in Koen. & Sims, Ann. of Bot. I, 507 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,80	
D. ulicina; Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,81	M.fr.II,106.
D. acicularis; Smith in Koen. & Sims, Ann. of Bot. I, 506 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,81	
D. pachyphylla; F. v. M., fragm. IV, 15 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,82	M.fr.IV,15.
D. teretifolia; R. Brown in Benth. Fl. Austr. II, 82 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,82	
D. genistifolia; Cunningham in Annal. Wien. Mus. II, 75 (1838)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,82	
D. hakeoides; Meissner in Lehm. plant. Preiss. I, 47 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,82	
D. colletioides; Meissner in Lehm. plant. Preiss. I, 48 (1844)	W.A.	—	—	—	—	—	—	B.fl.II.83	
D. reversifolia; F. v. M., fragm. I, 145 (1859)	W.A.	—	—	—	—	—	—	B.fl.II,83	M.fr.I,145.
D. incrassata; Smith in Transact. Linn. Soc. IX, 255 (1808)	W.A.	S.A.	—	—	—	—	—	B.fl.II,83	M.fr.I,145.
D. brevifolia; Lindley in Mitch. Three Exped. II, 201 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,84	
D. acanthoclona; F. v. M., fragm. X, 32 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,32.
D. Preissii; Meissner in Lehm. plant. Preiss. I, 50 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,84	
D. spinosissima; Meissner in Lehm. plant. Preiss. I, 51 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,85	
D. pachylina; Turczaninow in Bull. Soc. Mosc. XXVI, 263 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,85	

D. quadrilatera; Bentham in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,85	
D. striata; Turczaninow in Bull. Soc. Mosc. XXVI, 264 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,85	M.fr.II,105.
D. polyphylla; Bentham, in Bot. Regist. XXV, App. XIV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,86	
D. microphylla; Bentham, Fl. Austr. II, 86 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,86	
D. flexuosa; Bentham in Hueg. enum. 32 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,86	M.fr.I,145.
D. pectinata; Lindley in Mitch. Three Exped. II, 151 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,87	
D. trigonophylla; Meissner in Lehm. plant. Preiss. II, 213 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,87	
D. epiphylla; Meissner in Regensb. Bot. Zeit. 27 (1855)	W.A.	—	—	—	—	—	—	B.fl.II,87	
D. euphorbioides; Bentham, Fl. Austr. II, 88 (1864)	W.A.	—	—	—	—	—	—	B.fl.II.88	
D. divaricata; Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. paniculata; Bentham in Hueg. enum. 31 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. aphylla; F. v. M. in Benth. Fl. Austr. 88 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. juncea; Smith in Transact. Linn. Soc. IX, 260 (1808)	W.A.	—	—	—	—	—	—	B.fl.II,88	
D. alata; Smith in Transact. Linn. Soc. IX, 259 (1808)	—	—	—	—	N.S.W.	—	—	B.fl.II,89	
D. anceps; Turczaninow in Bull. Soc. Mosc. XXVI, 266 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,89	
AOTUS; Smith in Koenig & Sims Ann. of Bot. I, 504 (1805).									
A. gracillima; Meissner in Lehm. pl. Preiss. I, 59 (1844)	W.A.	—	—	—	—	—	—	B.fl.II.90	
A. villosa; Smith in Koen. & Sims, Ann. of Bot. I, 504 (1805)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,90	
A. mollis; Bentham in Mitch. Trop. Austr. 236 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,91	
A. Preissii; Meissner in Lehm. plant. Preiss. II, 214 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,91	
A. phylicoides; F. v. M. in Benth. Fl. Austr. II, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,92	
A. lanigera; Cunningham in Annal. des Wien. Mus. II, 78 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.II.92	
A. genistoides; Turczaninow in Bull. Soc. Mosc. XXVI, 268 (1853)	W.A.	—	—	—	—	—	—	B.fl.II.92	
A. carinata; Meissner in Lehm. plant. Preiss. II, 215 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,93	
A. passerinoides; Meissner in Lehm. plant. Preiss. I, 61 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,93	
A. Tietkensii; F. v. M., fragm. X, 33 (1876)...	W.A.	—	—	—	—	—	—	—	M fr.X.33.
A. cordifolia; Bentham in Hueg. enum. 33 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,93	
PHYLLOTA; De Candolle, prodr. II, 113 (1825). (Urodon.)									
P. barbata; Bentham in Hueg. enum. 33 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,94	
P. gracilis; Turczaninow in Bull. Soc. Mosc. XXVI, 267 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,94	
P. Sturtii; Bentham, Fl. Austr. II, 95 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,95	
P. phylicoides; Bentham in Ann. des Wien. Mus. II, 77 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,95	M.fr.X,33.
P. humifusa; Bentham, Fl. Austr. II, 95 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,95	
P. diffusa; F. v. M., fragm. I, 8 (1858)	—	—	T.	—	—	—	—	B.fl.II,119	M.fr.I,8.
P. pleurandroides; F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854)	—	S.A.	—	V.	—	—	—	B.fl.II,96	M.fr.X,33.
P. Luehmanni; F. v. M., fragm. X, 33 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,33.
P. Urodon; F. v. M. in papers of the Roy. Soc. of Tasm. 120 (1877)	W.A.	—	—	—	—	—	—	B.fl.II,124	
PULTENAEA; Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793). (Euchilus, Spadostyles, Bartlingia, Pulteneya.)									
P. daphnoides; Wendland, Bot. Beob. 49 (1798)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,112	
P. stricta; Sims, Bot. Mag. 1588 (1813)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,112	
P. retusa; Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,113	
P. Benthamii; F. v. M. in Transact. phil. Inst. Vict. I, 38 (1854)	—	—	—	V.	—	—	—	B.fl.II,113	
P. pycnocephala; F. v. M. in Benth. Fl. Austr. II, 114 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,114	
P. myrtoides; Cunningham in Annal. des Wien. Mus. II, 81 (1838)	—	—	—	—	—	Q.	—	B.fl.II,114	
P. mucronata; F. v. M., fragm. I, 8 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,114	M.fr.I,8.
P. polifolia; Cunningham in Field's N. S. Wales 346 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,115	
P. petiolaris; Cunningham in Ann. des Wien. Mus. II, 82 (1838)...	—	—	—	—	—	Q.	—	B.fl.II,115	
P. paleacea; Willdenow, spec. plant. II, 506 (1799)...	—	—	—	V.	N.S.W.	—	—	B.fl.II,115	

P. Gunnii; Bentham in Ann. des Wien. Mus. II, 82 (1839)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,116	
P. scabra; R. Brown in Ait. hort. Kew. sec. ed. III, 18 (1811)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,116	
P. Hartmanni; F. v. M., fragm. VIII, 166 (1874)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII,166.
P. microphylla; Sieber in De Candolle prodr. II, 11 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,117	
P. Skinneri; F. v. M., fragm. VIII, 166 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,166.
P. Drummondii; Meissner in Lehm. plant. Preiss. II, 219 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,117	
P. pinifolia; Meissner in Lehm. plant. Preiss. II, 220 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,118	
P. pedunculata; Hooker, Bot. Magaz. t. 2859 (1828)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,118	
P. conferta; Bentham, Fl. Austr. II, 118 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,118	
P. aciphylla; Bentham in Hueg. enum. 35 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,119	
P. ochreata; Meissner in Lehm. plant. Preiss. I, 73 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,120	
P. aspalathoides; Meissner in Lehm. plant. Preiss. I, 73 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,120	
P. obcordata; Bentham, Fl. Austr. II, 120 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,120	
P. rotundifolia; Bentham, Fl. Austr. II, 121 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,121	M.fr.I,145.
P. calycina; Bentham, Fl. Austr. II, 121 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,121	
P. spinulosa; Bentham, Fl. Austr. II, 121 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,121	
P. tenella; Bentham, Fl. Austr. II, 122 (1864)	—	—	—	V.	—	—	—	B.fl.II,122	
P. ternata; F. v. M., fragm. I, 8 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,122	M.fr.I,242; III,21; IV,10.
P. styphelioides; Cunningham in G. Don, gen. syst. II, 124 (1832)	—	—	—	V.	N.S.W.	—	—	B.fl.II,122	M.fr.III,21.
P. obovata; Bentham, Fl. Austr. II, 123 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,123	
P. incurvata; Cunningham in Field's N. S. Wales 346 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,124	
P. subumbellata; Hooker, Bot. Mag. t. 3254 (1833)...	—	—	T.	V.	N.S.W.	—	—	B.fl.II,124	
P. stipularis; Smith, Specim. of the Bot. of New Holl. 35, t. 12 (1793)	—	—	—	—	N.S.W.	—	—	B.fl.II,125	
P. glabra; Bentham, Fl. Austr. II, 124 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,125	
P. dentata; Labillardière, Nov. Holl. pl. spec. I, 103, t. 131 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,125	
P. aristata; Sieber in De Candolle, prodr. II, 112 (1825)	—	—	—	—	N.SW.	—	—	B.fl.II,126	
P. Baeuerlenii; F. v. M. in proceed. Linn. Soc. N.S.W. sec. ser. I, 1109 (1886)	—	—	—	—	N.S.W.	—	—	—	
P. plumosa; Sieber in De Candolle, prodr. II, 111 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,126	
P. viscosa; R. Brown in Benth. Fl. Austr. II, 127 (1864)	—	—	—	V.	N.S.W.	—	—	B.fl.II,127	
P. echinula; Sieber in De Candolle, prodr. II, 112 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,127	
P. hibbertioides; J. Hooker, Fl. Tasm. I, 89 (1860)...	—	—	T.	V.	N.S.W.	—	—	B.fl.II,127	
P. rosea; F. v. M., fragm. II, 15 (1860)	—	—	—	V.	—	—	—	B.fl.II,128	M.fr.II,15.
P. mollis; Lindley in Mitch. Three Exped. II, 260 (1838)	—	S.A.	—	V.	—	—	—	B.fl.II,128	
P. strobilifera; Meissner in Lehm. pl. Preiss. I, 75 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,128	
P. ericifolia; Bentham in Bot. Regist. XXV, App. XIII (1839)	W.A.	—	—	—	—	—	—	B.fl.II,129	
P. verruculosa; Turczaninow in Bull. Soc. Mosc. XXVI, 278 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,129	
P. empetrifolia; Meissner in Lehm. plant. Preiss. I, 76 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,129	
P. adunca; Turczaninow in Bull. Soc. Mosc. XXVI, 279 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,130	
P. neurocalyx; Turczaninow in Bull. Soc. Mosc. XXVI, 297 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,130	
P. rigida; R. Brown in Benth. Fl. Austr. II, 130 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,130	
P. juniperina; Labillardière, Nov. Holl. pl. spec. I, 102, t. 130 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,131	
P. acerosa; R. Brown in Benth. Fl. Austr. II, 131 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,131	
P. humilis; Bentham in J. Hook. Fl. Tasm. I, 91 (1860)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,131	
P. parviflora; Sieber in De Candolle, prodr. II, 111 (1852)...	—	—	—	V.	N.S.W.	—	—	B.fl.II,132	
P. setulosa; Bentham, Fl. Austr. II, 132 (1864)	—	—	—	—	—	Q.	—	B.fl.II,132	
P. vestita; R. Brown in Ait. hort. Kew. sec. ed. III, 19 (1811)	W.A.	S.A.	—	—	—	—	—	B.fl.II,132	
P. procumbens; Cunningham in Field's N. S. Wales 347 (1838)	—	—	—	—	N.S.W.	—	—	B.fl.II,133	
P. hispidula; R. Brown in Bentham, Fl. Austr. II, 133 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,133	
P. laxiflora; Bentham, Fl. Austr. I, 133 (1864)	—	S.A.	—	V.	—	—	—	B.fl.II,133	
P. largiflorens; F. v. M. in Benth. Fl. Austr. II, 134 (1864)	—	S.A.	—	V.	—	—	—	B.fl.II,134	

P. villosa ; Willdenow, spec. plant. II, 507 (1799)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,134	
P. foliolosa ; Cunningham in Ann. des Wien. Mus. II, 83 (1838)	—	—	—	V.	N.S.W.	—	—	B.fl.II,135	
P. flexilis ; Smith in Koen. & Sims, Ann. of Bot. I, 502 (1805)	—	—	—	V.	N.S.W.	—	—	B.fl.II,135	
P. euchila ; De Candolle, prodr. II, 112 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II.135	
P. selaginoides ; J. Hooker, Fl. Tasm. I, 87 (1860)	—	—	T.	—	—	—	—	B.fl.II,136	
P. densifolia ; F. v. M. in Transact. Vict. Inst. 119 (1854)...	—	S.A.	—	V.	—	—	—	B.fl.II,136	
P. elliptica ; Smith in Transact. Linn. Soc. IX, 246 (1808)..	—	—	—	—	N.S.W.	—	—	B.fl.II,136	
P. subspicata ; Bentham, Fl. Austr. II, 137 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,137	
P. villifera ; Sieber in De Candolle, prodr. II, 111 (1825)	—	S.A.	V.	—	N.S.W.	—	—	B.fl.II,137	
P. involucrata ; Bentham, Fl. Austr. II, 138 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,138	
P. Muelleri ; Bentham, Fl. Austr. II, 138 (1864)	—	—	—	V.	—	—	—	B.fl.II,138	
P. polyota ; F. v. M. in J. Hooker, Fl. Tasm. I, 89 (1860)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,138	
P. canaliculata ; F. v. M. in Transact. Vict. Inst. 119 (1854)	—	S.A.	—	V.	—	—	—	B.fl.II,139	
P. fasciculata ; Bentham in Ann. des Wien. Mus. II, 82 (1838)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,139	
P. tenuifolia ; R. Brown in Bot. Mag. t. 2086 (1819)	W.A.	S.A.	T.	V.	—	—	—	B.fl.II,139	M.fr.I,9.
P. viscidula ; Tate in Transact. Roy. Soc. of S. Austr. VI, 140 (1883)	—	S.A.	—	—	—	—	—	—	
LATROBEA ; Meissner in Lehm. plant. Preiss. II, 219 (1847). (Leptocytisus.)									
L. pungens ; Bentham, Fl. Austr. II, 140 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,140	M.fr.106.
L. genistoides ; Meissner in Lehm. plant. Preiss. II, 219 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,141	
L. Brunonis ; Meissner in Lehm. plant. Preiss. II, 219 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,141	
L. tenella ; Bentham, Fl. Austr. II, 141 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,141	
L. hirtella ; Bentham, Fl. Austr. II, 142 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,142	
L. diosmifolia ; Bentham, Fl. Austr. II, 142 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,142	
EUTAXIA ; R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811). (Sclerothamnus.)									
E. cuneata ; Meissner in Lehm. plant. Preiss. I, 65 (1844)...	W.A.	—	—	—	—	—	—	B.fl.II,143	
E. myrtifolia ; R. Brown in Ait. hort. Kew. sec. ed. III, 16 (1811)	W.A.	—	—	—	—	—	—	B.fl.II,144	
E. epacridoides ; Meissner in Lehm. plant. Preiss. I, 64 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,144	
E. virgata ; Bentham in Hueg. enum. 34 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,144	
E. densifolia ; Turczaninow in Bull. Soc. Mosc. XXVI, 271 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,145	
E. dillwynioides ; Meissner in Lehm. plant. Preiss. I, 63 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,145	
E. parvifolia ; Bentham in Hueg. enum. 34 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,145	
E. empetrifolia ; Schlechtendal, Linnaea XX, 667 (1847)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,145	M.fr.I,7.
DILLWYNIA ; Smith in Koenig & Sims, Ann. of Bot. I, 510 (1805).									
D. hispida ; Lindley in Mitch. Three Exped. II, 251 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,147	
D. ericifolia ; Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805)..	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,147	
D. floribunda ; Smith in Koen. & Sims, Ann. of Bot. I, 510 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,149	
D. Preissii ; Bentham, Fl. Austr. II, 149 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,149	
D. brunioides ; Meissner in Lehm. plant. Preiss. I, 62 (1844)	—	—	—	—	N.S.W.	—	—	B.fl.II,150	
D. juniperina ; Sieber in Hueg. enum. plant. N. Holl. occ. 33 (1837)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,150	
D. pungens ; Mackay in den Annal. des Wien. Mus. II, 79 (1838)	W.A.	—	—	—	—	—	—	B.fl.II,150	
D. cinerascens ; R. Brown in Bot. Mag. t. 2247 (1821)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,151	
D. divaricata ; Bentham, Fl. Austr. II, 151 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,151	
D. patula ; F. v. M., fragm. IV, 16 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,151	M.fr.IV,16.
EUCHILOPSIS ; F. v. M. in Melbourne Chemist, June (1882).									
E. linearis ; F. v. M. in Melb. Chem., June (1882)	W.A.	—	—	—	—	—	—	B.fl.II,67	
PLATYLOBIUM ; Smith in Transact. Linn. Soc. II, 350 (1794).									
P. formosum ; Smith in Transact. Linn. Soc. II, 350 (1794)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,153	M.fr.IX,157.
P. obtusangulum ; Hooker, Bot. Mag. t. 3258 (1833)	—	S.A.	T.	V.	—	—	—	B.fl.II,153	

P. triangulare; R. Brown in Ait. hort. Kew. sec. ed. IV, 266 (1812)	—	S.A.	T.	V.	—	—	—	B.fl.II,152	M.fr.IX,158.
P. alternifolium; F. v. M. in Wing's S. Sc. Rec. III, 99 (1883)	—	—	—	V.	—	—	—	—	
BOSSIAEA; Ventenat, Jardin de Cels. I, 7, t. 7 (1800). (Lalage, Scottia.)									
B. dentata; Bentham, Fl. Austr. II, 156 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,156	
B. Aquifolium; Bentham, Fl. Austr. II, 157 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,157	
B. Webbii; F. v. M. in Melb. Chemist No. 56, p. 65 (1882)	W.A.	—	—	—	—	—	—	—	
B. strigillosa; Bentham, Fl. Austr. II, 157 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,157	
B. cordigera; Bentham in J. Hook. Fl. Tasm. I, 95, t. 16 (1860)...	—	—	T.	V.	—	—	—	B.fl.II,157	M.fr.III,100; IX,45.
B. lenticularis; Sieber in De Candolle, prodr. II, 117 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,157	
B. Kiamensis; Bentham, Fl. Austr. II, 158 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,158	
B. ornata; F. v. M., fragm. IV, 13 (1863)	W.A.	—	—	—	—	—	—	B.fl.II,158	M.fr.IX,45.
B. eriocarpa; Bentham in Hueg. enum. 36 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,159	M.fr.IX,45.
B. divaricata; Turczaninow in Bull. Soc. Mosc. XXVI, 285 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,159	M.fr.IV,12.
B. calycina; Bentham, Fl. Austr. II, 159 (1864)	W.A.	—	—	—	—	—	—	B.fl II,159	
B. foliosa; Cunningham in Field's N. S. Wales, 347 (1825)...	—	—	—	V.	N.S.W.	—	—	B.fl.II,160	M.fr.III,100; IX,45.
B. cinerea; R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,160	M.fr.III,98.
B. biloba; Bentham in Hueg. enum. 36 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,160	M.fr.IX,45.
B. Preissii; Meissner in Lehm. pl. Preiss. I, 82 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,161	M.fr.IX,45.
B. concinna; Bentham, Fl. Austr. II, 161 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,161	
B. carinalis; Bentham in Mitch. Trop. Austr. 290 (1848)	—	—	—	—	—	Q.	—	B.fl.II,161	M.fr.IX,45.
B. rupicola; Cunningham in Benth. Fl. Austr. II, 162 (1864)	—	—	—	—	—	Q.	—	B.fl.II,162	
B. linophylla; R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812)	W.A.	—	—	—	—	—	—	B.fl.II,162	M.fr.IX,45.
B. disticha; Lindley, Bot. Regist. t. 55 (1841)	W.A.	—	—	—	—	—	—	B.fl.II,162	
B. prostrata; R. Brown in Ait. hort. Kew. sec. ed. IV, 268 (1812)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,162	M.fr.I,9; III,97.
B. Scortechinii; F. v. M. in Wing's S. Sc. Record III, 1 (1883)	—	—	—	—	N.S.W.	—	—	—	
B. Neo-Anglica; F. v. M., fragm. V, 106 (1866)	—	—	—	—	N.S.W.	—	—	—	M.fr.V,106.
B. buxifolia; Cunningham in Field's N. S. Wales, 348 (1825)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,163	M.fr.I,9; III,98; IX,45.
B. Brownii; Bentham, Fl. Austr. II, 163 (1864)	—	—	—	—	—	Q.	—	B.fl.II,163	
B. rhombifolia; Sieber in De Candolle, prodr. II, 117 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,164	M.fr.III,99; IX,45.
B. pulchella; Meissner in Lehm. pl. Preiss. I, 84 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,164	
B. microphylla; Smith in Transact. Linn. Soc. IX, 303 (1808)	—	—	—	V.	N.S.W.	—	—	B.fl.II,164	M.fr.III,99.
B. peduncularis; Turczaninow in Bull. Soc. Mosc. XXVI, 287 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,165	
B. Stephensoni; F. v. M. in proceed. L. S. of N.S.W. sec. ser. I, 1107 (1886) ...	—	—	—	—	N.S.W.	—	—	—	
B. heterophylla; Ventenat, Jardin de Cels. t. 7 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,165	M.fr.I,10; III,97.
B. rufa; R. Brown in Ait. hort. Kew. sec. ed. IV, 267 (1812)	W.A.	—	—	—	—	—	—	B.fl.II,165	M.fr.III,97; IX,45.
B. bracteosa; F. v. M. in Benth. Fl. Austr. II, 166 (1864)	—	—	—	V.	—	—	—	B.fl.II,166	
B. riparia; Cunningham in Benth. Fl. Austr. II, 166 (1864)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,166	
B. ensata; Sieber in De Candolle, prodr. II, 117 (1825)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,167	M.fr.II,120; XI,27&123.
B. scolopendria; Smith in Transact. Linn. Soc. IX, 307 (1808)	—	—	—	—	N.S.W.	—	—	B.fl.II,167	M.fr.II,120; III,95.
B. Walkeri; F. v. M., fragm. II, 120 (1861)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,167	M.fr.III,97,166; IX,45; XI,27.
B. Armitii; F. v. M., fragm. IX, 44 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,44.
B. phylloclada; F. v. M. in Transact. phil. Inst. Vict. III, 52 (1859)	—	—	—	—	—	—	N.A.	B.fl.II,168	M.fr.II,120.
TEMPLETONIA; R. Brown in Ait. hort. Kew. sec. ed. 269 (1812).									
T. retusa; R. Brown in Ait. hort. Kew. sec. ed. IV, 269 (1812)	W.A.	S.A.	—	—	—	—	—	B.fl.II,169	M.fr.IX,158.
T. Drummondii; Bentham, Fl. Austr. II, 169 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,169	
T. Muelleri; Bentham, Fl. Austr. II, 169 (1864)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,169	M.fr.I,9; II,178; IX,158.
T. aculeata; Bentham, Fl. Austr. II, 170 (1864)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.II,170	M.fr.II,120.
T. egena; Bentham, Fl. Austr. II, 170 (1864)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,170	M.fr.III,94; IX,158.
T. Battii; F. v. M. in Melb. Chemist new ser. II, 31 (1887)	W.A.	S.A.	—	—	—	—	—	—	
T. sulcata; Bentham, Fl. Austr. II, 171 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,171	M.fr.III,94,163.

HOVEA; R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812). (Poiretia, Plagiolobium, Platychilum.)

H. linearis; R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,172					
H. heterophylla; Cunningham in J. Hook. Fl. Tasm. I, 93, t. 15 (1860)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,172	M.fr.IX,158.				
H. longifolia; R. Brown in Ait. hort. Kew. sec. ed. IV, 275 (1812)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,172	M.fr.IX,158.				
H. acutifolia; Cunningham in G. Don, gen. syst. II, 126 (1832)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,174	M.fr.IX,158.				
H. longipes; Bentham in Hueg. enum. 37 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,174	M.fr.IX,158.				
H. acanthoclada; F. v. M., fragm. IV, 15 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,174	M.fr.IV,15.				
H. chorizemifolia; De Candolle, prodr. II, 116 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,174	M.fr.IX,158.				
H. Celsi; Bonpland, plant. cult. à Malmais. t. 51 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,175	M.fr.IX,158.				
H. trisperma; Bentham in Hueg. enum. 37 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,176	M.fr.IX,158.				
H. stricta; Meissner in Lehm. pl. Preiss. I, 79 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,176	M.fr.IX,158.				
H. pungens; Bentham in Hueg. enum. 37 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,176					

NEMATOPHYLLUM; F. v. M. in Hook. Kew. miscell. IX, 20 (1857).

N. Hookeri; F. v. M. in Hook. Kew. misc. IX, 20 (1857)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,170	M.fr.I,169; IV,170; IX,157.				

GOODIA; Salisbury, paradis. Londin. t. 41 (1806).

G. lotifolia; Salisbury, parad. Lond. t. 41 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,177	M.fr.I,10; IX,157.				
G. medicaginea; F. v. M., Second Gen. Rep. 11 (1854)	W.A.	S.A.	—	V.	—	—	—	—	M.fr.I,10,242; IX,157.				

CROTALARIA; Hermann, hort. acad. Lugd. Bat. 196—202, t. 197—203 (1687). (Pentadynamis.)

C. verrucosa; Linné, spec. plant. 715 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,179	M.fr.III,54; IX,156.	As., Afr., Am., P.			
C. crispata; F. v. M. in Benth. Fl. Austr. II, 179 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,179	M.fr.III,55.				
C. juncea; Linné, spec. plant. 714 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,179	M.fr.III,51; IX,156.	As., P.			
C. linifolia; Linné, fl. suppl. 322 (1781)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,180	M.fr.III,55; IX,156.	As., P.			
C. humifusa; Graham in Wall. numeric. list 5421 (1828)	—	—	—	—	—	Q.	N.A.	—	M.fr.VI,225; IX,156.	As.			
C. calycina; Schranck, pl. rar. hort. Monac. t. 12 (1819)	—	—	—	—	—	Q.	N.A.	B.fl.II,180	M.fr.III,56; IX,156.	As., P.			
C. retusa; Linné, spec. plant. 715 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,181	M.fr.III,51; IX,156.	As., Afr., Am.			
C. Mitchelli; Bentham in Mitch. Trop. Austr. 120 (1848)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,181	M.fr.III,50; IX,156.				
C. Novae-Hollandiae; De Candolle, prodr. II, 127 (1825)	W.A.	—	—	—	—	Q.	N.A.	B.fl.II,181	M.fr.III,55; IX,156.				
C. crassipes; Hooker, icon. plant. t. 830 (1852)	—	—	—	—	—	—	N.A.	B.fl.II,182					
C. Cunninghamii; B. Brown, App. to Sturt's Centr. Austr. 8 (1849)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.II,182	M.fr.III,52; IX,156.				
C. medicaginea; Lamarck, encycl. meth. II, 201 (1786)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,183	M.fr.III,56; IX,157.	As., Afr., Am., P.			
C. incana; Linné, spec. plant. 716 (1753)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,183	M.fr.III,53; IX,157.	As., Afr., Am., P.			
C. dissitiflora; Bentham in Mitch. Trop. Austr. 386 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,184	M.fr.III,56; IX,156.				
C. striata; De Candolle, prodr. II, 137 (1825)	—	—	—	—	—	Q.	—	—	...	...	...	As., Afr.	
C. laburnifolia; Linné, spec. plant. 715 (1753)	—	—	—	—	—	Q.	—	B.fl.II,184	M.fr.III,52; IX,157.	As., P.			
C. quinquefolia; Linné, spec. plant. 716 (1753)	—	—	—	—	—	Q.	—	B.fl.II,184	M.fr.IX,157	...	As., P.		
C. alata; Hamilton in D. Don, prodr. Fl. Nepal. 241 (1825)	—	—	—	—	—	Q.	N.A.	—	M.fr.XI,138	...	As.		

WESTONIA; Sprengel, syst. veg. III, 152 (1826).

W. trifoliata; Sprengel, syst. veg. III, 230 (1826)	—	—	—	—	—	—	N.A.	B.fl.II,185	...	...	...	As.	

TRIGONELLA; Linné, gen. pl. 351 (1737).

T. suavissima; Lindley in Mitch. Three Exped. I, 255 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,187					

LOTUS; Tournefort, inst. 402, t. 227 (1700) from C. Bauhin (1623).

L. corniculatus; Linné, spec. plant. 775 (1753)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,188	...	...	...	E., As., Afr.	
L. australis; Andrews, Botanist's Reposit. t. 624 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,188	M.fr.XI,20&27	...	P.		

PSORALEA; Linné, gen. ed. sec. 358 (1742).

P. badocana; Bentham, Fl. Austr. II, 190 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,190	M.fr.IX,156	...	As., P.		
P. cephalantha; F. v. M., fragm. IV, 35 (1863)	—	—	—	—	—	Q.	N.A.	—	M.fr.IV,35.				

P. Martini; F. v. M., fragm. V, 11 (1865)	—	—	—	—	—	—	N.A.	—	M.fr.V,11.	
P. Archeri; F. v. M., fragm. IV, 21 (1863)	—	—	—	—	—	—	N.A.	B.fl.II,190	M.fr.IV,21.	
P. balsamica; F. v. M. in Transact. Vict. Inst. III, 55 (1858)	—	S.A.	—	—	—	—	N.A.	B.fl.II,191		
P. plumosa; F. v. M., fragm. IV, 22 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,191	M.fr.IV,22.	
P. pustulata; F. v. M. in Transact. Vict. Inst. III, 54 (1858)	W.A.	—	—	—	—	—	N.A.	B.fl.II,191		
P. lachnostachys; F. v. M., fragm. III, 105 (1863)	—	—	—	—	—	—	N.A.	B.fl.II,191	M.fr III,105.	
P. criantha; Bentham in Mitch. Trop. Austr. 131 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,192	M.fr.III,45.	
P. patens; Lindley in Mitch. Three Exped. II, 9 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,192	M.fr.IX,155; III,45; IV,22.	
P. cinerea; Lindley in Mitch. Three Exped. II, 65 (1838)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,192	M.fr.III,45; IV,22.	
P. leucantha; F. v. M. in Transact. Vict. Inst. III, 54 (1859)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.II,193	M.fr.IX,156.	
P. parva; F. v. M. in Transact. Vict. Inst. I, 40 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,194	M.fr.IX,155.	
P. tenax; Lindley in Mitch. Three Exped. II, 10 (1838)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,193	M.fr.III,45; IX,155.	
P. adscendens; F. v. M. in Transact. Vict. Inst. I, 40 (1854)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,193	M.fr.III,45; IX,155.	
P. Testariae; F. v. M., fragm. V, 45 (1865)	—	—	—	—	—	Q.	—	—	M.fr.V,45,189.	
INDIGOFERA; Royen, pl. hort. Lugd. 372 (1740), from Isnard (1724).										
I. linifolia; Retzius, observ. bot. IV, 29 (1786)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,195	M.fr.III,101; IX,44	As., Afr., P.
I. cordifolia; Heyne in Roth, nov. pl. spec. 357 (1821)	—	—	—	—	—	—	N.A.	B.fl.II,196	... ...	As., Afr.
I. enneaphylla; Linné, mantissa altera 272 (1771)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,196	M.fr.III,102; IX,44	As., Afr. P.
I. glandulosa; Willdenow, spec. plant. III, 1227 (1800)	—	—	—	—	—	Q.	—	B.fl.II,196	M.fr.IV,22; VII,106; IX,44.	As.
I. haplophylla; F. v. M., fragm. III, 102 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.II,196	M.fr.IX,44.	
I. trifoliata; Linné, amoen. acad. IV, 327 (1759)	—	—	—	—	—	Q.	—	B.fl.II,197	M.fr.III,164 ...	As., P.
I. trita; Linné fil., suppl. plant. 335 (1781)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,197	M.fr.III,103 ...	As., Afr., P.
I. parviflora; Heyne in Wall. numeric. list 5457 (1828)	—	—	—	—	—	Q.	N.A.	B.fl.II,197	M.fr.III,103; IX,44	As., Afr.
I. viscosa; Lamarck, Encycl. méth. III, 247 (1789)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,198	M.fr.III,104; IX,44	As., Afr.
I. hirsuta; Linné, spec. plant. 751 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,198	M.fr.III,105; IX,44	As., Afr., Am., P.
I. pratensis; F. v. M., essay on plants of Burdek. Exped. 10 (1860)	—	—	—	—	—	Q.	—	B.fl.II,198		
I. Schultziana; F. v. M., fragm. VII, 105 (1870)	—	—	—	—	—	—	N.A.	—	M.fr.VII,105.	
I. monophylla; De Candolle, prodr. II, 222 (1825)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,199	M.fr.III,44; VII,106; IX,44.	
I. rugosa; Bentham, Fl. Austr. II, 199 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,199		
I. saxicola; F. v. M. in Benth. Fl. Austr. II, 199 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,199		
I. Australis; Willdenow, spec. plant. III, 1235 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,199		
I. brevidens; Bentham in Mitch. Trop. Austr. 385 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,200		
I. efoliata; F. v. M. in Melb. Chemist. Nov. (1883)	—	—	—	—	N.S.W.	—	—	—		
I. Baileyi; F. v. M., fragm. IX, 43 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,43.	
I. coronillifolia; Cunningham in Benth. Fl. Austr. II, 201 (1864)	—	S.A.	—	—	N.S.W.	—	—	B.fl.II,201		
PTYCHOSEMA; Bentham in Lindl. Bot. Regist. XXV. App. XVI (1839).										
P. pusillum; Bentham in Lindl. Bot. Regist. XXV, App. XVI (1839)	W.A.	—	—	—	—	—	—	B.fl.II,201		
P. anomalum; F. v. M., fragm. IX, 62 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,62.	
P. trifoliolatum; F. v. M. in Wing's S. Science Record II, 72 (1882)	W.A.	S.A.	—	—	N.S.W.	—	—	—		
LAMPROLOBIUM; Bentham, Fl. Austr. II, 202 (1864).										
L. fruticosum; Bentham, Fl. Austr. II, 202 (1864)	—	—	—	—	—	Q.	—	B.fl.II,202	M.fr.IX,68.	
L. megalophyllum; F. v. M., fragm. IX, 67 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.IX,67.	
TEPHROSIA; Persoon, synops. pl. II, 328 (1807).										
T. coriacea; Bentham, Fl. Austr. II, 204 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,204	M.fr.IX,65&66.	
T. lamprolobioides; F. v. M., fragm. IX, 64 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.IX,64.	
T. Forrestiana; F. v. M., fragm. XI, 98 (1880)	—	—	—	—	—	—	N.A.	—	M.fr.XI,98.	
T. flammea; F. v. M. in Bentham Fl. Austr. II, 204 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,204		
T. reticulata; Bentham, Fl. Austr. II, 205 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,205	M.fr.IX,64.	

T. crocea; Bentham, Fl. Austr. II, 205 (1864) ... ... — — — — — — N.A. B.fl.II,205
T. oblongata; Bentham, Fl. Austr. II, 205 (1864) ... ... — — — — — — N.A. B.fl.II,205
T. porrecta; Bentham, Fl. Austr. II, 206 (1864) ... ... — — — — — — N.A. B.fl.II,206
T. polyzyga; F. v. M. in Benth. Fl. Austr. II, 206 (1864) ... ... — — — — — — N.A. B.fl.II,206
T. graminifolia; F. v. M. in Benth. Fl. Austr. II, 206 (1864) ... ... — — — — — — N.A. B.fl II,206 M.fr.IX,6
T. simplicifolia; F. v. M. in Benth. Fl. Austr. II, 206 (1864) ... ... — — — — — — N.A. B.fl.II,206
T. nematophylla; F. v. M., fragm. IX, 63 (1879) ... ... — — — — — — N.A. — M.fr.IX,63.
T. leptoclada; Bentham, Fl. Austr. II, 207 (1864) ... ... — — — — — Q. N.A. B.fl.II.207
T. brachycarpa; F. v. M. in Benth. Fl. Austr. II, 207 (1864) ... ... — — — — — — N.A. B.fl.II,207
T. uniovulata; F. v. M., fragm. XI, 70 (1879) ... ... — — — — — — N.A. — M.fr.IX,70.
T. Stuartii; Bentham, Fl. Austr. II, 207 (1864) ... ... — — — — — — N.A. B.fl.II,207
T. conferta; F. v. M., fragm. IX, 65 (1879) ... ... — — — — — — N.A. — M fr.IX,75.
T. eriocarpa; Bentham, Fl. Austr. II, 207 (1864) ... ... — — — — — — N.A. B.fl.II,207
T. phaeosperma; F. v. M. in Benth. Fl. Austr. II, 207 (1864) ... ... — — — — — — N.A. B.fl.II,208
T. astragaloides; Bentham, Fl. Austr. II, 208 (1864) ... ... — — — — — Q. — B.fl.II,208
T. juncea; Bentham, Fl. Austr. II, 208 (1864) ... ... — — — — — Q. N.A. B.fl.II,208 M.fr.IX,64
T. filipes; Bentham, Fl. Austr. II, 208 (1864) ... ... — — — — — Q. N.A. B.fl.II,208
T. sphaerospora; F. v. M. in Wing's S. Sc. Record III, 128 (1883) ... ... — S.A. — — — — N.A. —
T. remotiflora; F. v. M. in Benth. Fl. Austr. II, 209 (1864) ... ... — — — — — — N.A. B.fl.II,209
T. oligophylla; Bentham, Fl. Austr. II, 209 (1864) ... ... — — — — — Q. N.A. B.fl.II,209
T. macrocarpa; Bentham, Fl. Austr. II, 209 (1864)... ... ... — — — — — — N.A. B.fl.II,209
T. purpurea; Persoon, synops. plant. II, 329 (1807)... ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.II,209 ... .. As, Afr., P.
T. Bidwilli; Bentham, Fl. Austr. II, 210 (1864) ... ... — — — — N.S.W. Q. — B.fl.II,210
T. rosea; F. v. M. in Benth. Fl. Austr. II, 211 (1864) ... ... — — — — — — N.A. B.fl.II,211

WISTARIA; Nuttall, Gen. of N. Amer. pl. II, 115 (1818). (Milletia.)

W. megasperma; F. v. M., fragm. I, 10 (1858) ... ... — — — — N.S.W. Q. — B.fl.II,211 M.fr.I,10.
W. australis; F. v. M. in Britten's Journ. of Bot. XXII, 290 (1884) ... ... — — — — N.S.W. — — —

SESBANIA; Scopoli, Introd. 308 (1777). (Agati from Alpino 1592.)

S. grandiflora; Persoon, synops. pl. II, 316 (1807) ... ... W.A. — — — — Q. N.A. B.fl.II,212 M.fr.II,88; XI,33 ... As.
S. Aegyptiaca; Persoon, synops. pl. II, 316 (1807) ... ... — — — — — Q. N.A. B.fl.II,212 M.fr.XI,33 ... ... As., Afr
S. aculeata; Persoon, synops. pl. II, 316 (1807) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.II,213 M.fr.XI,33 ... ... As., Afr., P.
S. simpliciuscula; F. v. M. in Benth. Fl. Austr. II, 213 (1864) ... . — — — — — — N.A. B.fl.II,213
S. brachycarpa; F. v. M., fragm. XI, 32 (1878) ... ... — — — — — — N.A. — M.fr.IX,32.

CARMICHHAELIA; R. Brown in Bot. Regist. XI, 912 (1825).

C. exsul; F. v. M., fragm. VII, 126 (1871) ... ... — — — — N.S.W. — — — M.fr.VII,121,193.

CLIANTHUS; Banks & Solander in G. Don, dichlam. pl. II, 468; indicative (1832). (Donia.)

C. Dampieri; Cunningham in Trans. hort. soc. Lond. sec. ser. I, 522 (1835) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.II,214

STREBLORRHIZA; Endlicher, prodr. fl. Norfolk. 92 (1833).

S. speciosa; Endlicher, prodr. fl. Norf. 93 (1833) ... ... — — — — N.S.W. — — —

SWAINSONIA; Salisbury, parad. Londin. t. 28 (1806). (Cyclogyne, Diplolobium.)

S. Greyana; Lindley, Bot. Regist. t. 66 (1846) ... ... — S.A. — V. N.S.W. Q. — B.fl.II,216 M.fr.IX,154; X,7.
S. galegifolia; R. Brown in Ait. hort. Kew. sec. ed. III, 327 (1811) ... ... — — — — N.S.W. Q. — B.fl.II,217
S. coronillifolia; Salisbury, paradis. Lond. t. 28 (1806) ... ... — S.A. — — N.S.W. Q. — B.fl.II,217 M.fr.XI,70.
S. brachycarpa; Bentham, Fl. Austr. II, 217 (1864)... ... ... — — — — N.S.W. Q. — B.fl.II,217 M.fr.IX,155.
S. phacoides; Bentham in Mitch. Trop. Austr. 363 (1848) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.II,217
S. Burkittii; F. v. M. in Benth. Fl. Austr. II, 218 (1864) ... ... — S.A. — — N.S.W. — — B.fl.II,218
S. Burkei; F. v. M. in Benth. Fl. Austr. II, 218 (1864) ... ... W.A. S.A. — — — — N.A. B.fl.II,218 M.fr.IX,155.

S. oligophylla; F. v. M. in Benth. Fl. Austr. II, 219 (1864)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.II,219		
S. campylantha; F. v. M. in Gregory's pap. rel. to Leichh. Exp. 6 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,219	M.fr.III,46; IX,154.	
S. occidentalis; F. v. M., fragm. III, 46 (1862)	W.A.	—	—	—	—	—	N.A.	B.fl.II,219	M.fr.III,46.	
S. stenodonta; F. v. M., fragm. XI, 70 (1879)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,70.	
S. oncinotropis; F. v. M. in Melb. Chem. Oct. (1884)	—	—	—	V.	N.S.W.	—	—	—		
S. gracilis; Bentham, Fl. Austr. II, 220 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,220		
S. plagiotropis; F. v. M., fragm. IX, 153 (1875)	—	—	—	V.	N.S.W.	—	—	—	M.fr.IX,143.	
S. procumbens; F. v. M., fragm. III, 46 (1862)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,220	M.fr.I,76; III,46; IX,154; XI,27.	
S. Drummondii; Bentham, Fl. Austr. II, 220 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,220		
S. Oliveri; F. v. M. in Wing's S. Sc. Rec. II, 152 (1882)	W.A.	S.A.	—	—	N.S.W.	—	—	—		
S. canescens; F. v. M., fragm. III, 46 (1862)	W.A.	—	—	—	—	—	—	B.fl.II,221	M.fr.III,46; IX,154; XI,70.	
S. Beasleyana; F. v. M. in Melb. Chemist, new ser. II, 84 (1882)	W.A.	—	—	—	—	—	—	—		
S. stipularis; F. v. M. in Linnaea XXV, 393 (1852)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,221	M.fr.I,76; IX,154.	
S. oroboides; F. v. M. in Benth. Fl. Austr. II, 222 (1864)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,222		
S. Kingii; F. v. M. in Transact. Roy. Soc. Vict. XXIII, 53 (1886)	W.A.	—	—	—	—	—	N.A.	—		
S. unifoliolata; F. v. M., fragm. VIII, 226 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,226.	
S. lessertiifolia; De Candolle, prodr. II, 271 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,222	M.fr.IX,154.	
S. luteola; F. v. M., fragm. I, 75 (1858)	—	—	—	—	—	Q.	—	B.fl.II,223	M.fr.XI,70.	
S. parviflora; Bentham, Fl. Austr. II, 223 (1864)	—	—	—	—	—	Q.	—	B.fl.II,223		
S. monticola; Cunningham in A. Gray's Bot. Wilk. Exp. I, 411 (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,223	M.fr.IX,155.	
S. microphylla; A. Gray, Bot. Wilk. Exped. I, 410 (1854)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,223	M.fr.IX,155.	
S. laxa; R. Brown in Sturt's Centr. Austr. App. 18 (1849)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,224		
S. colutoides; F. v. M., fragm. X, 6 (1876)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,6.	
S. Fraseri; Bentham, Fl. Austr. II, 224 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,224		
S. Macullochiana; F. v. M., fragm. VII, 25 (1869)	W.A.	—	—	—	—	—	N.A.	—	M.fr.IX,154.	
GLYCYRRHIZA; Tournefort, inst. 389, t. 210 (1700) from Dioscorides and Plinius. (Clidanthera.)										
G. psoraleoides; Bentham, Fl. Austr. II, 225 (1864)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,225	M.fr.III,45.	
ORMOCARPUM; Palisot, Fl. d'Ovar. I. 95, t. 58 (1805).										
O. scnnoides; De Candolle, prodr. II, 315 (1825)	—	—	—	—	—	Q.	—	B.fl.II,226	...	... As.
AESCHYNOMENE; Linné, gen. pl. 350 (1737).										
A. Indica; Linné, spec. plant. 713 (1753)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,227	M.fr.XII,18	... As., Afr., P.
A. falcata; De Candolle, prodr. II, 322 (1825)	—	—	—	—	—	Q.	—	B.fl.II,227	M.fr.XII,18	... Afr., Am.
A. Americana; Linné, spec. plant. 713 (1753)	—	—	—	—	—	Q.	—	—	M.fr.XII,18	... As., Am.
SMITHIA; Aiton, hort. Kew. III, 496 (1789).										
S. capitata; Desvaux, journ. de Bot. I, (1813)	—	—	—	—	—	Q.	N.A.	B.fl.II,228	...	... As., P.
S. sensitiva; Aiton, hort. Kew. III, 496, t. 13 (1789)	—	—	—	—	—	Q.	—	—	M.fr.VII,27; IX,193.	As., Afr.
ZORNIA; Gmelin, Syst. Nat. 1076 (1791).										
Z. diphylla; Persoon, synops. II, 318 (1807)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,228	...	... ... As., Afr., Am., P.
Z. chaetophora; F. v. M. in Transact. phil. Inst. Vict. III, 56 (1857)	—	—	—	—	—	—	N.A.	B.fl.II,229		
DESMODIUM; Desvaux, Journ. de Bot. III, 122 (1813). (Dicerma, Phyllodium, Pteroloma.)										
D. umbellatum; De Candolle, prodr. II, 325 (1825)	—	—	—	—	—	Q.	—	B.fl.II,230	M.fr.IX,66	... As., Afr., Am.
D. pulchellum; Bentham, Fl. Hongk. 83 (1861)	—	—	—	—	—	—	N.A.	B.fl.II,231	...	... ... As., P.
D. biarticulatum; F. v. M., fragm. II, 121 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.II,231	M.fr.IX,66	... As.
D. acanthocladum; F. v. M., fragm. II, 122 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,231	M.fr.IX,66.	
D. triquetrum; De Candolle, prodr. II, 326 (1825)	—	—	—	—	—	Q.	—	—	M.fr.IX,66	... As.
D. Gangeticum; De Candolle, prodr. II, 327 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.II,232	...	... ... As., As., P.
D. dependens; Blume in Miq. Fl. Ind. Batav. I, 249 (1855)	—	—	—	—	—	Q.	—	—	M.fr.VIII,225,284; IX,66&195; X,120.	As., P.

D. brachypodum; A. Gray, Bot. Wilk. Exped. 434 (1854)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,232						
D. varians; Endlicher in Ann. des Wien. Mus. I, 185 (1839)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,232						P.
D. flagellare; Bentham, Fl. Austr. II, 233 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,233						
D. rhytidophyllum; F. v. M. in Benth. Fl. Austr. II, 223 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,233						
D. campylocaulon; F. v. M. in Benth. Fl. Austr. II, 233 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,233						
D. nemorosum; F. v. M. in Benth. Fl. Austr. II, 234 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,234						
D. neurocarpum; Bentham, Fl. Austr. II, 234 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,234						
D. trichostachyum; Bentham, Fl. Austr. II, 234 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,234						
D. polycarpum; De Candolle, prodr. II, 334 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,235	M.fr.IX,66					As., Afr., P.
D. trichocaulon; De Candolle, prodr. II, 335 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,235						
D. Muelleri; Bentham, Fl. Austr. II, 235 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,235						
D. reniforme; De Candolle, prodr. II, 327 (1825)	—	—	—	—	—	Q.	N.A.	—	M.fr.XII					As.
D. parvifolium; De Candolle, prodr. II, 334 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,235	M.fr.IX,66					As., P.
PYCNOSPORA; R. Brown in Wight & Arnott, prodr. I, 197 (1834).														
P. hedyaaroides; R. Brown in Wight & Arnott, prodr. 197 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.II,236	M.fr.V,30					As., P.
URARIA; Desvaux, Journ. de Bot. III, 122 (1813).														
U. picta; Desvaux, Journ. de Bot. III, 122 (1813)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,237						As., Afr., P.
U. cyliudracea; Bentham, Fl. Austr. II, 237 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,237						
U. Iagopoides; De Candolle, prodr. II, 324 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.II,237						As., P.
LOUREA; Necker, elem. Bot. III, 17 (1790).														
L. obcordata; Desvaux, Journ. de Bot. III, 122 (1813)	—	—	—	—	—	—	N.A.	B.fl.II,238						As., P.
ALYSICARPUS; Necker, elem. Bot. III, 15 (1790).														
A. vaginalia; De Candolle, prodr. II, 353 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.II,239						As., Afr., P.
A. longifoliua; Wight & Arnott, prodr. 223 (1834)	—	—	—	—	—	—	N.A.	B.fl.II,239						As.
A. cylindricus; Desvaux in Annal. Soc. Linn. Par. 301 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.II,239						As., Afr.
LESPEDEZA; Cl. Richard in Michaux, fl. bor. am. II, 70 (1803).														
L. cuneata; G. Don, gen. syst. II, 307 (1832)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,240						As.
L. lanata; Bentham, Fl. Austr. II, 241 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.II,241						
CLITORIA; Linné, gen. plant. 216 (1737), from Petiver.														
C. Ternatea; Linné, sp. pl. 753 (1753)	—	—	—	—	—	Q.	—	—						As., Afr., P.
C. Australis; Bentham, Fl. Austr. II, 242 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,242						
GLYCINE; Linné, gen. pl. 349 (1737). (Leptolobium, Leptocyamus.)														
G. falcata; Bentham, Fl. Austr. II, 243 (1864)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,243						
G. clandestina; Wendland, Bot. Beob. 54 (1798)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl,II,243						
G. Latrobeana; Bentham, Fl. Austr. II, 224 (1864)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,244						
G. tabacina; Bentham, Fl. Austr. II, 244 (1864)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,244						P.
G. sericea; Bentham, Fl. Austr. II, 245 (1864)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,245						
G. tomentosa; Bentham, Fl. Austr. II, 245 (1864)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,245						As., P.
KENNEDYA; Ventenat, Jardin de la Malmaison II, t. 104—106 (1804). (Caulina, Moench 1802, Hardenbergia, Physolobium, Zichya, Amphodus.)														
K. nigricans; Lindley, Bot. Regist. t. 1715 (1835)	W.A.	—	—	—	—	—	—	B.fl.II,249	M.fr.VII,127.					
K. Beckxiana; F. v. M., fragm. XI, 98 (1880)	W.A.	—	—	—	—	—	—	—	M.fr.XI,98.					
K. rubicunda; Ventenat, Jard. de la Malm. t. 104 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,249	M.fr.VII,127.					
K. procurrens; Bentham in Mitch. trop. Austr. 365 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,249						
K. prorepens; F. v. M., fragm. VIII, 225 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,225.					
K. prostrata; R. Brown in Ait. hort. Kew. sec. ed. IV, 299 (1812)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,250	M.fr.VII,128.					
K. bracteata; Gaudichaud in Freyc. voy. 286, t. 113 (1826)	W.A.	—	—	—	—	—	—	B.fl.II,250	M.fr.IV,79.					

K. eximia; Lindley in Paxt. Mag. XVI, 35 (1849)	W.A.	—	—	—	—	—	—	B.fl.II,250			
K. microphylla; Meissner in Lehm. pl. Preiss. I, 91 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,251			
K. parviflora; Meissner in Lehm. pl. Preiss. I, 91 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,251			
K. Stirlingii; Lindley, Bot. Regist. t. 1845 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,252			
K. glabrata; Lindley Bot. Regist. t. 1838 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,252	M.fr.VII,128.		
K. macrophylla; F. v. M., fragm. IV, 79 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,252	M.fr.IV,79.		
K. coccinea; Ventenat, Jard. de la Malm. t. 105 (1804)	W.A.	—	—	—	—	—	—	B.fl.II,250	M.fr.VII,128.		
K. monophylla; Ventenat, Jard. de la Malm. t. 106 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,246	M.fr.VII,128.		
K. Comptoniana; Link, enum. pl. hort. bot. Berol. II, 235 (1822)	W.A.	—	—	—	—	—	—	B.fl.II,247	M.fr.VII,128.		
K. retusa; F. v. M., fragm. V, 106 (1866)	—	—	—	—	—	Q.	—	B.fl.II,247	M.fr.VII,128	...	P.
ERYTHRINA; Linné, gen. pl. 216 (1737).											
E. vespertilio; Bentham in Mitch. Trop. Austr. 218 (1848)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.II,253			
E. Indica; Lamarck, Encyclop. méthod. II, 391 (1786)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,253	...	...	As., P.
STRONGYLODON; T. Vogel in Linnaea X, 585 (1836).											
S. ruber; T. Vogel in Linnaea X, 585 (1836)	—	—	—	—	—	Q.	—	—	...	...	As., P.
MUCUNA; Marcgraf, hist. nat. Brazil. 18 (1648).											
M. gigantea; De Candolle, prodr. II, 405 (1825)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,254	...	...	As., P.
GALACTIA; P. Browne, Civil & Nat. Hist. of Jamaica 298 (1756).											
G. tenuiflora; Wight & Arnott, prodr. fl. penins. Ind. 206 (1834)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,255	...	...	As., Afr., Am., P.
G. Muelleri; Bentham, Fl. Austr. II, 255 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,255			
CANAVALIA; De Candolle, prodr. II, 403 (1825). (Canavali.)											
C. obtusifolia; De Candolle, prodr. II, 404 (1825)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,256	...	...	As., Afr., Am., P.
PHASEOLUS; Tournefort, inst. 412, t. 232 (1700), from Dioscorides & Columella.)											
P. Truxillensis; Humb. Bonpl. & Kunth, nov. gen. Amer. VI, 451 (1823)	—	—	—	—	—	Q.	N.A.	B.fl.II,257	...	...	As., Afr., Am., P.
P. Max; Linné, sp. pl. 725 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,257	...	...	As., Afr., P.
VIGNA; Savi, Dissert. 6 (1824). (Callicystus.)											
V. vexillata; Bentham in Mart. Fl. Bras. Papil. 193, t. 50 (1862)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,258	...	...	As., Afr., Am., P.
V. lutea; A. Gray, Bot. Wilk. Exped. I, 454 (1854)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,259	...	...	As., Afr., Am., P.
V. luteola; Bentham in Mart. Fl. Bras. Papil. 194, t. 50 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,260	...	...	As., Afr., Am.
V. lanceolata; Bentham in Mitch. trop. Austr. 350 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,260			
DOLICHOS; Linné, gen. plant. 222 (1737).											
D. biflorus; Linné, spec. plant. 727 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,261	...	...	As., Afr.
DUNBARIA; Wight & Arnott, prodr. I, 258 (1834).											
D. conspersa; Bentham in plant. Junghuhn. (1851)	—	—	—	—	—	Q.	—	B.fl.II,262	...	...	As.
CAJANUS; De Candolle, Catal. hort. Monsp. 85 (1813). (Cajan, Atylosia.)											
C. marmoratus; F. v. M., Cens. 41 (1882) from Benth. (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,263			
C. scarabaeoides; F. v. M., Cens. 41 (1882) from Benth. (1851)	—	—	—	—	—	Q.	—	B.fl.II,263	...	...	As., Afr.
C. reticulatus; F. v. M., Cens. 41 (1882) from Benth. (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,263			
C. cinereus; F. v. M., Cens. 41 (1882)	W.A.	—	—	—	—	—	N.A.	B.fl.II,264			
C. grandifolius; F. v. M., Essay on pl. Burdekin Exped. 9 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.II,264			
C. confertiflorus; F. v. M., Essay on pl. Burdekin Exped. 9 (1860)	—	—	—	—	—	Q.	—	B.fl.II,264			
RHYNCHOSIA; Loureiro, Fl. Cochinch. II, 460 (1790).											
R. rhomboidea; F. v. M. in Benth. Fl. Austr. II, 265 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,265			
R. acutifolia; Bentham, Fl. Austr. II, 266 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,266			

R. rostrata ; Bentham, Fl. Austr. II, 266 (1864) ... ... — — — — — — N.A. B.fl.II,266
R. Cunninghamii ; Bentham, Fl. Austr. II, 266 (1864) ... ... — — — — — Q. — B.fl.II,266 .. .. P.
R. minima ; De Candolle, prodr. II, 385 (1825) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.II,267 ... . As., Afr., Am., P.

ERIOSEMA ; De Candolle, prodr. II, 388 (1825).
E. Chinense ; T. Vogel in Meyen's Beitr. zur Bot. 31 (1843) ... ... — — — — — Q. N.A. B.fl.II,268 ... ... As., P.

FLEMINGIA ; Roxburgh in Ait. hort. Kew. sec. ed. IV, 349 (1812).
F. lineata ; Roxburgh in Ait. hort. Kew. sec. ed. IV, 350 (1812) ... . — — — — — Q. N.A. B.fl.II,268 ... ... As., P.
F. pauciflora ; Bentham, Fl. Austr. II, 269 (1864) ... ... — — — — — Q. N.A. B.fl.II,269
F. parviflora ; Bentham, Fl. Austr. II, 269 (1864) ... ... — — — — — Q. — B.fl.II,269
F. involucrata ; Bentham in pl. Jungh. I, 246 (1851) ... ... — — — — — Q. — B.fl.II,269 ... ... As.

ABRUS ; Dalechamps, hist. general. plant. append. 193 (1587), from Rauwolf (1582).
A. precatorius ; Linné, syst. natur. ed. XII, 533 (1767) ... ... — — — — — Q. N.A. B.fl.II,270 ... ... As., Afr., Am., P.

DALBERGIA ; Linné, fil. suppl. 52 (1781).
D. densa ; Bentham in Hook. Lond. Journ. II, 217 (1843) ... ... — — — — — Q. — B.fl.II,271 ... ... P.

LONCHOCARPUS ; Humboldt, Bonpland & Kunth, nov. gen. VI, 383 (1823).
L. Blackii ; Bentham, Fl. Austr. II, 271 (1864) ... ... — — — — N.S.W. Q. — B.fl.II,271 M.fr.II,123.

DERRIS ; Loureiro, Fl. Cochinch. II, 432 (1790). (Brachypterum.)
D. scandens ; Bentham in Journ. Linn. Soc. IV, suppl. 103 (1860) ... ... — — — — N.S.W. Q. — B.fl.II,272 ... ... As.
D. uliginosa ; Bentham in pl. Jungh. I, 252 (1851) ... ... — — — — — Q. N.A. B.fl.II,272 ... ... As., Afr., P.

PONGAMIA ; Lamarck in Ventenat, Jardin de la Malmaison t. 28 (1803) from Adanson (1763).
P. glabra ; Ventenat, Jard. Malm. t. 28 (1803) ... ... — — — — — Q. N.A. B.fl.II,273 ... . As., P.

SOPHORA ; Linné, gen. plant. 125 (1737). (Calpurnia partim.)
S. tomentosa ; Linné, spec. plant. 373 (1753) .. .. — — — — N.S.W. Q. N.A. B.fl.II,274 M.fr.VII,27 ... As., Afr., Am., P.
S. Fraseri ; Bentham, Fl. Austr. II, 274 (1864) ... ... — — — — N.S.W. Q. — B.fl.II,274 M.fr.V,31 ; VII,27.
S. tetraptera ; J. Miller, icon. plant. t. 1 (1780) ... ... — — — — N.S.W. — — — M.fr.VII,26 ... N.Z.

PODOPETALEM ; F. v. M., in Melbourne Chemist, June (1882).
P. Ormondi ; F. v. M. in Melb. Chem. June (1882) ... ... — — — — — Q. — —

CASTANOSPERMUM ; Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830).
C. Australe ; Cunningham & Fraser in Hook. bot. misc. I, 241, t. 51 (1830) .. — — — — N.S.W. Q. — B.fl.II,275 M.fr.VII,27.

BARKLYA ; F. v. M., fragm. I, 109 (1859).
B. syringifolia ; F. v. M. in Journ. Linn. Soc. III, 158 (1859) ... ... — — — — N.S.W. Q. — B.fl.II,275 M.fr.I,109; III,163; VIII,27.

CAESALPINA ; Plumier, nov. pl. Amer. gen. 28, t. 9 (1703) from Ammand. (Guilandina, Caesalpinia 1810.)
C. Bonducella ; Fleming in Asiat. Research XI, 159 (1801) ... ... — — — — N.S.W. Q. N.A. B.fl.II,276 ... ... As., Afr., Am., P.
C. Nuga ; W. T. Aiton, hort. Kew. sec. ed. III, 32 (1811) ... ... — — — — — Q. — B.fl.II,277 M.fr.X,7 ... ... As., P.
C. sepiaria ; Roxburgh, Fl. Indic. II, 360 (1832) ... ... — — — — — Q. — B.fl.II,277 M.fr.X,7 ... ... As.

MEZONEURON ; Desfontaines in Mém. du Mus. IV, 245, t. 10 et 11 (1818).
M. brachycarpum ; Bentham, Fl. Austr. II, 278 (1864) ... ... — — — — N.S.W. Q. — B.fl.II,278 M.fr.X,7.
M. Scortechinii ; F. v. M. in Wing's South. Sc. Record II. 73 (1882) ... ... — — — — N.S.W. Q. — —

PTEROLOBIUM ; R. Brown in Salt. Abyssin. 65 (1814).
P. nitens ; F. v. M. in Bentham, Fl. Austr. II, 279 (1864) ... ... — — — — — Q. — B.fl.II,279 M.fr.X,7.

PELTOPHORUM ; Vogel in Linnaea XI, 406 (1837).
P. ferrugineum ; Bentham, Fl. Austr. II, 279 (1864) ... ... — — — — — Q. N.A. B.fl.II,279 M.fr.X,7 ... ... As.

CASSIA; Tournefort, inst. 619, t. 392 (1700), from Plumier (1693). (Cathartocarpus.)

C. Brewsteri; F. v. M., fourth ann. Rep. 17 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,282	M.fr.I,110; X,8.			
C. laevigata; Willdenow, enum. pl. hort. bot. Berol. I, 441 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,282	M.fr.IV,14; X,8	...	Am.	
C. Sophera; Linné, spec. plant. 379 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,283	M.fr.III,49; IV,14; X,9.		As., P.	
C. laxiflora; Bentham, Fl. Austr. II, 283 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,283				
C. magnifolia; F. v. M., fragm. I, 166 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.II,283	M.fr.X,9.			
C. venusta; F. v. M., fragm. I, 165 (1859)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.II,284	M.fr.X,9.			
C. notabilis; F. v. M., fragm. III, 28 (1862)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,284	M.fr.X,9.			
C. pleurocarpa; F. v. M., fragm. I, 223 (1859)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,284	M.fr.X,9.			
C. glauca; Lamarck, Encycl. méth. I, 647 (1783)	—	—	—	—	S.W.	Q.	N.A.	B.fl.II,285	M.fr.IV,13; X,9	...	As., P.	
C. retusa; Solander in Linnaea XI, 72 (1842)	—	—	—	—	—	Q.	—	B.fl.II,285				
C. Australis; Sims in Bot. Mag. t. 2676 (1826)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.II,285	M.fr.X,9.			
C. Chatelainiana; Gaudichaud in Freyc. voy. 485, t. 111 (1826)	W.A.	—	—	—	—	—	N.A.	B.fl.II,286				
C. glutinosa; De Candolle, prodr. II, 495 (1825)	—	S.A.	—	—	—	—	N.A.	B.fl.II,286				
C. Cuthbertsoni; F. v. M. in Vict. Nat. V, 75 (1888)	W.A.	—	—	—	—	—	—	—				
C. pruinosa; F. v. M., fragm. III, 48 (1862)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,286	M.fr.III,48.			
C. circinata; Bentham in Mitch. trop. Austr. 384 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,286	M.fr.I,165.			
C. phyllodinea; R. Brown, App. to Sturt's Centr. Austr. 15 (1849)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,287				
C. eremophila; Cunningham in T. Vogel, gen. Cass. synops. 47 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,287				
C. artemisioides; Gaudichaud in De Candolle, prodr. II, 495 (1825)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,288				
C. Sturtii; R. Brown, App. to Sturt's Centr. Austr. 14 (1849)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,288				
C. desolata; F. v. M. in Linnaea XXV, 387 (1852)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,289				
C. oligophylla; F. v. M., fragm. III, 49 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,289	M.fr.III,49.			
C. cardiosperma; F. v. M., fragm. X, 50 (1877)	W.A.	—	—	—	—	—	—	—	M.fr.X,50.			
C. oligoclada; F. v. M., fragm. III, 49 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,289	M.fr.X,9.			
C. heptanthera; F. v. M., fragm. X, 8 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.X,8.			
C. leptoclada; Bentham, Fl. Austr. II, 290 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,290				
C. Absus; Linné, spec. plant. 376 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.II,290	M.fr.III,50; X,9	...	As., Afr.	
C. pumila; Lamarck, Encycl. méth. I, 620 (1783)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,290	M.fr.III,47; X,10	...	As.	
C. concinna; Bentham, Fl. Austr. II, 291 (1864)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,291	M.fr.X,10.			
C. mimosoides; Linné, spec. plant. 379 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,291	M.fr.III,48; X,10	...	As., Afr., Am., P.	

PETALOSTYLIS; R. Brown, App. to Sturt's Centr. Austr. 79 (1849). (Petalostylos, Petalogyne.)

P. labichoides; R. Brown, App. to Sturt's Centr. Austr. 17 (1849)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,292	M.fr.X,7.			

LABICHEA; Gaudichaud in De Candolle, prodr. II, 507 (1825).

L. cassioides; Gaudichaud in De Candolle, prodr. II, 507 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,292				
L. nitida; Bentham, Fl. Austr. II, 293 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.II,293	M.fr.X,7.			
L. lanceolata; Bentham in Hueg. enum. 41 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,293	M.fr.X,7.			
L. rupestris; Bentham in Mitch. Trop. Austr. 342 (1848)	—	—	—	—	—	Q.	—	B.fl.II,293	M.fr.X,7; XII,19.			
L. punctata; Bentham in Bot. Regist. XXV, App. XV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,294	M.fr.X,8; XII,19.			
L. Buettneriana; F. v. M. in Melb. Chemist, June (1882)	—	—	—	—	—	Q.	—	—	M.fr.XII,18.			

TAMARINDUS; Tournefort, inst. 660, t. 445, App. (1700) from C. Bauhin (1623).

T. Indicus; Linné, spec. plant. 34 (1753)	—	—	—	—	—	—	N.A.	B.fl.II,294	...	...	...	As., Afr.

BAUHINIA; Plumier, nov. gen. pl. Amer. 22, t. 13 (1703).

B. acuminata; Linné, spec. plant. 375 (1753)	—	—	—	—	—	Q.	—	—	...	...	...	As.
B. Leichhardtii; F. v. M. in Transact. Vict. Inst. III, 50 (1858)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,295	M.fr.X,8.			
B. Carronii; F. v. M. in Transact. Vict. Inst. III, 49 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,295	M.fr.X,8.			
B. Hookeri; F. v. M. in Transact. Vict. Inst. III, 51 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.II,295	M.fr.X,8.			

B. Gilesii ; F. v. M. & Bailey in Wing's S. Scc. Record, II, 151 (1882) ...	...	—	—	—	—	—	—	N.A.	—					
B. monandra ; Kurz, Forest Flora of Burma I, 39 (1877) ...	...	...	...	—	—	—	—	—	Q	—	—	...	...	... As.
AFZELIA ; Smith in Transact. Linn. Soc. IV, 221 (1798).														
A. bijuga ; A. Gray, Bot. Wilk. Expl. Exped. 467, t. 51 (1854) ...	...	...	—	—	—	—	—		Q.	—	—	...	...	... As., P.
CYNOMETRA ; Linné in Act. Soc. Ups. 78 (1741).														
C. ramiflora ; Linné, spec. plant. 382 (1753) ...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.II,296	... ... As., P.
ERYTHROPHLAEUM ; Afzelius in Tuckey's Congo 430 (1818). (Laboucheria.)														
E. Laboucherii ; F. v. M. in Benth. Fl. Austr. II, 297 (1864) ...	...	...	—	—	—	—	—		Q.	N.A.	B.fl.II,297			
ENTADA ; Adanson, Familles des plantes II, 318 (1763).														
E. Pursaetha ; De Candolle, mémoir. sur la fam. des Légum. 12 (1825) ...	...	—	—	—	—	—		Q.	—	B.fl.II,298	M.fr.V,179	...	As., Afr., Am., P.	
ADENANTHERA ; Royen in Linné, coroll. 7 (1737).														
A. pavonina ; Linné, spec. plant. 384 (1753) ...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.V,30 ... As.
A. abrosperma ; F. v. M., fragm. V, 30 (1865) ...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,298	M.fr.V,31.
NEPTUNIA ; Loureiro, Fl. Cochinch. II, 653 (1790). (Dichrostachys.)														
N. cinerea ; F. v. M., Cens. 43 (1882) from Wight & Arnott (1834) ...	...	—	—	—	—	—		—	N.A.	B.fl.II,299	...	...	...	As.
N. spicata ; F. v. M., fragm. III, 151 (1862) ...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,299	M.fr.III,151.
N. gracilis ; Bentham in Hook. Journ. of Bot. IV, 355 (1842) ...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,300			
N. monosperma ; F. v. M. in Benth. Fl. Austr. II, 299 (1864) ...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,300			
ACACIA ; Tournefort, inst. 605, t. 375 (1700) from Dioscorides & Plinius. (Vachellia, Chithonanthus, Tetracheilus, Mimosa partly.)														
I. Alatae.														
A. glaucoptera ; Bentham in Schlecht. Linnaea XXVI, 604 (1853) ...	...	W.A.	—	—	—	—	—	—	B.fl.II,320					
A. alata ; R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,300				
A. diptera ; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,321			
A. stenoptera ; Bentham in Hook. Lond. Journ. I, 325 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,321				
II. Continuae.														
A. incurva ; Bentham in Hook. Lond. Journ. I, 325 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,322				
A. trigonophylla ; Meissner in Lehm. pl. Preiss. II, 199 (1846) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,322	M.fr.IV,3.			
A. continua ; Bentham, Fl. Austr. II, 322 (1864) ...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,322		
A. Peuce ; F. v. M., fragm. III, 151 (1863) ...	...	...	...	...	...	—	S.A.	—	—	—	Q.	—	B.fl.II,323	M.fr.III,151.
III. Pungentes.														
A. spinescens ; Bentham in Hook. Lond. Journ. I, 323 (1842) ...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,323				
A. cochlearis ; H. L. Wendland, comm. de Acac. aphyll. 15 (1820) ...	...	W.A.	S.A.	—	—	—	—	—	B.fl.II,324					
A. lanigera ; Cunningham in Field's N.S.W. 345 (1825) ...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,324			
A. phlebocarpa ; F. v. M. in Journ. Linn. Soc. III, 119 (1858) ...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,325				
A. trinervata ; Sieber in De Candolle, prodr. II, 451 (1825) ...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.II,325				
A. colletioides ; Cunningham in Hook. Lond. Journ. I, 336 (1842) ...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,325	M.fr.IV,4.				
A. striatula ; Bentham in Hook. Lond. Journ. I, 336 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,326				
A. campylophylla ; Bentham in Schlecht. Linnaea XXVI, 605 (1853) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,326				
A. teretifolia ; Bentham in Hook. Lond. Journ. I, 326 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,326				
A. sulcata ; R. Brown in Ait. hort. Kew. sec. ed. V, 460 (1813) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,327				
A. costata ; Bentham in Hook. Lond. Journ. I, 339 (1842) ...	...	...	...	W.A.	—	—	—	—	—	B.fl.II,327				
A. barbinervis ; Bentham in Hook. Lond. Journ. I, 326 (1842) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,327				
A. ataxiphylla ; Bentham in Schlecht. Linnaea XXVI, 605 (1853) ...	...	W.A.	—	—	—	—	—	—	B.fl.II,328					
A. Baxteri ; Bentham in Hook. Lond. Journ. I, 327 (1842) ...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,328			
A. aureo-nitens ; Lindley, Bot. Regist. XXV, App. XV (1839) ...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,328				
A. quadrisulcata ; F. v. M., fragm. III, 127 (1863) ...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,328	M.fr.III,127.	

A. erioclada; Bentham in Schlecht. Linnaea XXVI, 606 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,329	
A. siculiformis; Cunningham in Hook. Lond. Journ. I, 337 (1842)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,329	
A. patens; F. v. M. in Journ. Linn. Soc. III, 120 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,329	M.fr.III,46.
A. laricina; Meissner in Lehm. pl. Preiss. I, 6 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,330	
A. tetragonophylla; F. v. M. in Journ. Linn. Soc. III, 121 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,330	M.fr.IV,3.
A. genistoides; Cunningham in Benth. Fl. Austr. II, 330 (1864)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,330	
A. sphacelata; Bentham in Hook. Lond. Journ. I, 338 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,331	
A. ingrata; Bentham, Fl. Austr. II, 331 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,331	
A. juniperina; Willdenow, spec. plant. IV, 1049 (1805)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,331	
A. asparagoides; Cunningham in Field's N. S. Wales 343 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,332	
A. tenuifolia; F. v. M. in Transact. phil. Soc. Vict. I, 37 (1854)	—	—	—	V.	—	—	—	B.fl.II,332	
A. diffusa; Edwards, Bot. Regist. t. 634 (1822)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,332	
A. rupicola; F. v. M. in Schlecht. Linnaea XXVI, 610 (1853)	—	S.A.	—	V.	—	—	—	B.fl.II,333	
IV. Calamiformes.									
A. volubilis; F. v. M., fragm. X, 98 (1877)	W.A.	—	—	—	—	—	—	—	M.fr.X,98.
A. tetragonocarpa; Meissner in Lehm. pl. Preiss. I, 4 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,336	
A. restiacea; Bentham in Hook. Lond. Journ. I, 3 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,336	
A. squamata; Lindley, Bot. Regist. XXV, App. XV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,336	
A. brachyphylla; Bentham in Schlecht. Linnaea XXVI, 615 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,337	
A. Bynoeana; Bentham in Schlecht. Linnaea XXVI, 614 (1853)	—	—	—	—	—	—	N.A.	B.fl.II,337	M.fr.IV,9.
A. triptycha; F. v. M. in Benth. Fl. Austr. II, 337 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,337	
A. leptoneura; Bentham in Hook. Lond. Journ. I, 341 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,337	
A. rigens; Cunningham in Loudon, hort. Brit. II, 406 (1830)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,337	
A. papyrocarpa; Bentham, Fl. Austr. II, 338 (1864)	W.A.	S.A.	—	—	—	—	—	B.fl.II,338	
A. Gilesiana; F. v. M. in Melb. Chemist, July (1882)	—	S.A.	—	—	—	—	—	—	
A. sessiliceps; F. v. M. in Melb. Chemist, July (1882)	—	S.A.	—	—	—	—	—	—	
A. pugioniformis; Wendland, comm. de Acac. aphyll. 38, t. 9 (1820)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,338	
A. juncifolia; Bentham in Hook. Lond. Journ. I, 341 (1842)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,339	
A. calamifolia; Sweet in Bot. Regist. t. 839 (1824)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,339	
A. spodiosperma; F. v. M. in proc. L. Soc. of N.S.W. sec. ser. III, 164 (1888) ...	W.A.	—	—	—	—	—	—	—	
A. scirpifolia; Meissner in Bot. Zeitung 10 (1855)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.II,339	
A. extensa; Lindley, Bot. Regist. XXV, App. XV (1839)	W.A.	—	—	—	—	—	—	B.fl.II,340	
A. gonophylla; Bentham in Schlecht. Linnaea XXVI, 613 (1853)...	W.A.	S.A.	—	—	—	—	—	B.fl.II,340	
A. ericifolia; Bentham in Hook. Lond. Journ. I, 345 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,340	
A. uncinella; Bentham in Schlecht. Linnaea XXVI, 613 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,341	
V. Brunioidae.									
A. cedroides; Bentham in Schlecht. Linnaea XXVI, 615 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,341	
A. lycopodifolia; Cunningham in Hook. icon. plant. t. 172 (1837)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,342	
A. spondylophylla; F. v. M., fragm. VIII, 243 (1874)	—	S.A.	—	—	—	—	N.A.	—	M.fr.243.
A. galioides; Bentham in Hook. Lond. Journ. I, 344 (1842)	—	—	—	—	—	Q.	N.A.	B.fl.II,342	
A. Baueri; Bentham in Hook. Lond. Journ. I, 344 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,342	M.fr.XI,33.
A. subternata; F. v. M. in Journ. Linn. Soc. III, 124 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,343	
A. minutifolia; F. v. M., fragm. VIII, 243 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.X,243.
A. brunioides; Cunningham in G. Don, gen. syst. II, 404 (1832)	—	—	—	—	—	Q.	—	B.fl.II,343	
A. conferta; Cunningham in Hook. Lond. Journ. I, 345 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,343	M.fr.I,6.
VI. Uninerves.									
A. scabra; Bentham in Schlecht. Linnaea XXVI, 605 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,344	
A. nodiflora; Bentham in Schlecht. Linnaea XXVI, 621 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,344	
A. spinosissima; Bentham in Schlecht. Linnaea XXVI, 621 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,344	
A. ulicina; Meissner in Lehm. pl. Preiss. II, 202 (1846)	W.A.	—	—	—	—	—	—	B.fl.II,345	

A. erinacea; Bentham in Hook. Lond. Journ. I, 360 (1842)	W.A.	S.A.	—	—	—	—	—	B.fl.II,345	
A. Huegelii; Bentham in Hueg. enum. 42 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,345	
A. nervosa; De Candolle, Mém. sur la fam. des Légum. 444 (1825)	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. obovata; Bentham in Hook. Lond. Journ. I, 329 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. congesta; Bentham in Hook. Lond. Journ. I, 327 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. dermatophylla; Bentham, Fl. Austr. II, 346 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,346	
A. aspera; Lindley in Mitch. Three Exped. II, 139 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,347	
A. armata; R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,347	
A. idiomorpha; Cunningham in Hook. Lond. Journ. I, 329 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,348	
A. Shuttleworthii; Meissner in Lehm. pl. Preiss. I, 7 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,348	
A. Gregorii; F. v. M., fragm. III, 47 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,348	M.fr.III,47.
A. pilosa; Bentham in Schlecht. Linnaea XXVI, 607 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,348	
A. crispula; Bentham in Schlecht. Linnaea XXVI, 607 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,349	
A. crassistipulea; Bentham in Hook. Lond. Journ. I, 326 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,349	
A. hastulata; Smith in Rees' Cyclopaed. XXXIX, Suppl. (1819)	W.A.	—	—	—	—	—	—	B.fl.II,349	
A. horridula; Meissner in Lehm. pl. Preiss. I, 9 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,350	
A. divergens; Bentham in Hook. Lond. Journ. I, 331 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,350	
A. vomeriformis; Cunningham in Hook. Lond. Journ. I, 332 (1842)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,350	
A. plagiophylla; F. v. M. in Journ. Linn. Soc. III, 131 (1858)	—	—	—	—	N.S.W.	Q.	—	—	
A. biflora; R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,351	
A. decipiens; R. Brown in Ait. hort. Kew. sec. ed. V, 463 (1813)	W.A.	—	—	—	—	—	—	B.fl.II,351	
A. cuneata; Bentham in Hueg. enumer. 42 (1837)	W.A.	—	—	—	—	—	—	B.fl.II,351	
A. dilatata; Bentham in Schlecht. Linnaea XXVI, 608 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,352	
A. bidentata; Bentham in Hook. Lond. Journ. I, 333 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,352	
A. acanthoclada; F. v. M., fragm. III, 127 (1863)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,352	M.fr.III,127.
A. obliqua; Cunningham in Hook. Lond. Journ. I, 334 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,353	
A. acinacea; Lindley in Mitch. Three Exped. II, 267 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,353	
A. runciformis; Cunningham in G. Don, gen. syst. II, 404 (1832)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,353	M.fr.I,5; II,177.
A. lachnophylla; F. v. M. in Wing's S. Sc. Record II, 150 (1882)	W.A.	—	—	—	—	—	—	—	
A. triquetra; Bentham in Hook. Lond. Journ. I, 358 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,354	
A. ligustrina; Meissner in Lehm. pl. Preiss. II, 203 (1846)	W.A.	—	—	—	—	—	—	B.fl.II,354	
A. Meissneri; Lehmann, delect. sem. hort. bot. Hamb. (1842)	W.A.	—	—	—	—	—	—	B.fl.II,354	
A. anceps; De Candolle, Mém. sur la fam. des Légum. 446 (1825)	W.A.	S.A.	—	—	—	—	—	B.fl.II,355	
A. hispidula; Willdenow, spec. plant. IV, 1054 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,355	
A. undulifolia; Fraser in Loddiges, Bot. Cabinet XVI, 1544 (1829)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,355	
A. flexifolia; Cunningham in Hook. Lond. Journ. I, 359 (1842)	—	—	—	—	N.S.W.	—	—	B.fl.II,356	
A. dura; Bentham in Schlecht. Linnaea XXVI, 622 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,356	
A. spathulata; F. v. M. in Benth. Fl. Austr. II, 356 (1864)	W.A.	—	—	—	—	—	N.A.	B.fl.II,356	
A. microcarpa; F. v. M., second gen. Report 7 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,357	M.fr.I,6.
A. montana; Bentham in Hook. Lond. Journ. I, 360 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,357	
A. verniciflua; Cunningham in Field's N. S. Wales 344 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,358	
A. leprosa; Sieber in De Candolle, prodr. II, 450 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.II,358	
A. stricta; Willdenow, spec. plant. IV, 1052 (1805)	—	—	T.	V.	N.S.W.	—	—	B.fl.II,358	
A. dodonaeifolia; Willdenow, enum. pl. hort. Berol. suppl. 68 (1813)	—	S.A.	—	—	—	—	—	B.fl.II,359	
A. Gnidium; Bentham, Fl. Austr. II, 359 (1864)	—	—	—	—	—	Q.	—	B.fl.II,359	
A. ramosissima; Bentham in Hook. Lond. Journ. I, 356 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,360	
A. Dempsteri; F. v. M., fragm. XI, 65 (1879)	W.A.	—	—	—	—	—	—	—	M.fr.XI,65.
A. Sentis; F. v. M., second gen. report 11 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,360	M.fr.XI,65.
A. dentifera; Bentham in Maund, Botanist IV, t. 179 (1839)	W.A.	—	—	—	—	—	—	B.fl.II,361	
A. sclerosperma; F. v. M. in Wing's South. Sc. Rec. II, 150 (1882)	W.A.	—	—	—	—	—	N.A.	—	

A. fasciculifera; F. v. M. in Benth. Fl. Austr. II, 361 (1864)	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,361	
A. falcata; Willdenow, spec. plant. IV, 1053 (1805)	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.II,361	
A. macradenia; Bentham in Mitch. Trop. Austr. 360 (1848)	—	—	—	—	—	—	—	Q.	—	B.fl.II,362	
A. penninervis; Sieber in De Candolle, prodr. II, 452 (1825)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,362			
A. amblyphylla; F. v. M. in Wing's S. Sc. Record, II, 149 (1882)	W.A.	—	—	—	—	—	—	—			
A. retinodes; Schlechtendal, Linnaea XX, 664 (1847)	—	S.A.	—	V.	—	—	—	B.fl.II,362			
A. neriifolia; Cunningham in Hook. Lond. Journ. I, 357 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,363			
A. microbotrya; Bentham in Hook. Lond. Journ. I, 353 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,363			
A. leiophylla; Bentham in Hook. Lond. Journ. I, 351 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,364			
A. cyanophylla; Lindley, Bot. Regist. Misc. 49 (1835)	W.A.	—	—	—	—	—	—	B.fl.II,364			
A. pycnantha; Bentham in Hook. Lond. Journ. I, 351 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,365			
A. notabilis; F. v. M., fragm. I, 6 (1858)	W.A.	S.A.	—	—	N.S.W.	—	N.A.	B.fl.II,365	M.fr.I,6.		
A. gladiiformis; Cunningham in Hook. Lond. Journ. I, 354 (1842)	—	—	—	—	N.S.W.	—	—	B.fl.II,365			
A. obtusata; Sieber in De Candolle, prodr. II, 453 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.II,366			
A. rubida; Cunningham in Field's N.S.W. 344 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.II,366			
A. amoena; H. L. Wendland, comm. de Acac. aphyll. 10 t. 4 (1820)	—	—	—	V.	N.S.W.	—	—	B.fl.II,366			
A. hakeoides; Cunningham in Hook. Lond. Journ. I, 354 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,367			
A. salicina; Lindley in Mitch. Three Exped. II, 20 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,367			
A. rostellifera; Bentham in Hook. Lond. Journ. I, 356 (1842)	W.A.	—	—	—	—	—	—	B.fl.II,368			
A. crassiuscula; H. L. Wendland, comm. de Acac. 31, t. 8 (1820)	W.A.	—	—	—	—	—	—	B.fl.II,368			
A. Harveyi; Bentham, Fl. Austr. II, 368 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,368			
A. iteaphylla; F. v. M. in Schlecht. Linnaea XXVI, 617 (1853)	—	S.A.	—	—	—	—	—	—			
A. suaveolens; Willdenow, spec. plant. IV, 1050 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,369			
A. subcoerulea; Lindley, Bot. Regist. t. 1075 (1827)	W.A.	—	—	—	—	—	—	B.fl.II,369			
A. Graffiana; F. v. M. in Melb. Chemist, new ser. II, 118 (1887)	W.A.	—	—	—	—	—	—	—			
A. Lindleyi; Meissner in Lehm. pl. Preiss. I, 14 (1844)	W.A.	—	—	—	—	—	—	B.fl.II,370			
A. leptopetala; Bentham in Schlecht. Linnaea XXVI, 619 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,370			
A. Dietrichiana; F. v. M. in Wing's S. Sc. Rec. II, 149 (1882)	—	—	—	—	—	Q.	—	—			
A. Murrayana; F. v. M. in Benth. Fl. Austr. II, 370 (1864)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.II,370			
A. linifolia; Willdenow, spec. plant. IV, 1051 (1805)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,371			
A. Leichhardtii; Bentham, Fl. Austr. II, 372 (1864)	—	—	—	—	—	Q.	—	B.fl.II,372			
A. buxifolia; Cunningham in Field's N.S.W. 344 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,372			
A. lunata; Sieber in De Candolle, prodr. II, 452 (1825)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,373			
A. brachybotrya; Bentham in Hook. Lond. Journ. I, 347 (1842)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,373			
A. Wattsiana; F. v. M. in Benth. Fl. Austr. II, 374 (1864)	—	S.A.	—	—	—	—	—	B.fl.II,374			
A. podalyrifolia; Cunningham in Loudon, hort. Brit. 707 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,374			
A. uncifera; Bentham in Mitch. Trop. Austr. 341 (1848)	—	—	—	—	—	Q.	—	B.fl.II,374			
A. vestita; Edwards in Bot. Regist. t. 698 (1823)	—	—	—	V.	N.S.W.	—	—	B.fl.II,375			
A. cultriformis; Cunningham in G. Don. gen. syst. II, 406 (1832)	—	—	—	—	N.S.W.	—	—	B.fl.II,375	M.fr.XI,123.		
A. pravissima; F. v. M., first gen. report 12 (1853)	—	—	—	V.	N.S.W.	—	—	B.fl.II,375	M.fr.I,5,242.		
A. strongylophylla; F. v. M., fragm. VIII, 226 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,226.		
A. pyrifolia; De Candolle, Mém. sur la fam. des Légum. 447 (1825)	W.A.	—	—	—	—	—	N.A.	B.fl.II,376	M.fr.III,17.		
A. myrtifolia; Willdenow, spec. plant. IV, 1054 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,376			
VII. Plurinerves.											
A. scalpelliformis; Meissner in Lehm. pl. Preiss. II, 200 (1847)	W.A.	—	—	—	—	—	—	B.fl.II,377			
A. urophylla; Bentham in Bot. Reg. Misc. 24 (1841)	W.A.	—	—	—	—	—	—	B.fl.II,377			
A. Luehmanni; F. v. M., fragm. XI, 116 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.XI,116.		
A. sublanata; Bentham in Hueg. enum. 42 (1837)	—	S.A.	—	—	—	—	—	B.fl.II,378	M.fr.XI,117.		
A. pravifolia; F. v. M., fragm. I, 4 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.XI,117.		
A. amblygona; Cunningham in Hook. Lond. Journ. I, 332 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,378	M.fr.IV,3; XI,117.		

A. adnata; F. v. M. in Melb. Chemist, July (1882)	W.A.	—	—	—	—	—	—	—			
A. deltoidea; Cunningham in G. Don, gen. syst. II, 401 (1832)	—	—	—	—	—	—	N.A.	B.fl.II,378			
A. stipulosa; F. v. M. in Journ. Linn. Soc. III, 119 (1858)	W.A.	—	—	—	—	—	N.A.	B.fl.II,379	M.fr.XI,117.		
A. loxophylla; Bentham in Schlecht. Linnaea XXVI, 625 (1853)...	W.A.	—	—	—	—	—	—	B.fl.II,379			
A. setulifera; Bentham in Schlecht. Linnaea XXVI, 625 (1853)	—	—	—	—	—	—	N.A.	B.fl.II,379			
A. translucens; Cunningham in Hook. icon. plant. t. 160 (1837)	—	—	—	—	—	—	N.A.	B.fl.II,379			
A. impressa; F. v. M. in Journ. Linn. Soc. III, 133 (1858)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.II,380			
A. bivenosa; De Candolle, prodr. II, 452 (1825)	W.A.	—	—	—	—	—	N.A.	B.fl.II,380	M.fr.III,47.		
A. trincura; F. v. M., pl. of Vict. II, 25 (1860)	—	S.A.	—	V.	—	—	N.A.	B.fl.II,381	M.fr.IV,5.		
A. nitidula; Bentham, Fl. Austr. II, 381 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,381			
A. estrophiolata; F. v. M. in Wing's South. Sc. Rec. II, 150 (1882)	—	S.A.	—	—	—	—	—	—			
A. heteroclita; Meissner in Lehm. pl. Preiss. I, 318 (1845)	W.A.	—	—	—	—	—	—	B.fl.II,381	M.fr.IV,6.		
A. elongata; Sieber in De Candolle, prodr. II, 451 (1825)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,381			
A. subporosa; F. v. M., pl. of Vict. II, 24 (1860)	—	—	—	V.	N.S.W.	—	—	B.fl.II,382	M.fr.IV,5.		
A. Simsii; Cunningham in Hook. Lond. Journ. I, 368 (1842)	—	—	—	—	—	Q.	N.A.	B.fl.II,382	...	...	... P.
A. leptospermoides; Bentham in Schlecht. Linnaea XXVI, 626 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,383			
A. homalophylla; Cunningham in Hook. Lond. Journ. I, 365 (1842)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,383			
A. pendula; Cunningham in Loudon, hort. Brit. 490 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,383			
A. Osswaldi; F. v. M., pl. of Vict. II, 27 (1860)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,384	M.fr.IV,5.		
A. lineolata; Bentham in Schlecht. Linnaea XXVI, 626 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,384			
A. quadrimarginea; F. v. M., fragm. X, 31 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,31.		
A. coriacea; De Candolle, Mém. sur la fam. des Légum. 446 (1825)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.II,385	M.fr.III,47.		
A. praelongata; F. v. M. in Melb. Chem. Aug. (1883)	—	—	—	—	—	—	N.A.	—			
A. stenophylla; Cunningham in Hook. Lond. Journ. I, 366 (1842)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,385			
A. hemignosta; F. v. M. in Journ. Linn. Soc. III, 134 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,385	M.fr.XI,67.		
A. sclerophylla; Lindley in Mitch. Three Exped. II, 138 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,386			
A. farinosa; Lindley in Mitch. Three Exped. II, 145 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,386			
A. heteroneura; Bentham in Schlecht. Linnaea XXVI, 624 (1853)	W.A.	—	—	—	—	—	—	B.fl.II,387			
A. viscidula; Cunningham in Hook. Lond. Journ. I, 366 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,387			
A. ixiophylla; Bentham in Hook. Lond. Journ. I, 364 (1842)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.II,387	M.fr.IV,6.		
A. dictyophleba; F. v. M., fragm. III, 128 (1863)	—	S.A.	—	—	—	Q.	N.A.	B.fl.II,388	M.fr.III,128.		
A. cyclops; Cunningham in Loudon, hort. Brit. 407 (1830)	W.A.	S.A.	—	—	—	—	—	B.fl.II,388			
A. melanoxylon; R. Brown in Ait. hort. Kew. sec. ed. V, 462 (1813)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,388			
A. oraria; F. v. M., fragm. XI, 66 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,66.		
A. implexa; Bentham in Hook. Lond. Journ. I, 368 (1842)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,389			
A. harpophylla; F. v. M. in Benth. Fl. Austr. II, 389 (1864)	—	—	—	—	—	Q.	—	B.fl.II,389			
A. excelsa; Bentham in Mitch. Trop. Austr. 225 (1848)	—	—	—	—	—	Q.	—	B.fl.II,390	M.fr.IV,6		
A. homaloclada; F. v. M., fragm. XI, 34 (1878)	—	—	—	—	—	Q.	—	—	M.fr.XI,34.		
A. complanata; Cunningham in Hook. Lond. Journ. I, 369 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,390	M.fr.XI,34.		
A. dineura; F. v. M. in Journ. Linn. Soc. III, 130 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,391	M.fr.XI,65.		
A. binervata; De Candolle, prodr. II, 452 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,390	M.fr.XI,66.		
A. latescens; Bentham in Hook. Lond. Journ. I, 380 (1842)	—	—	—	—	—	—	N.A.	B.fl.II,391			
A. flavescens; Cunningham in Hook. Lond. Journ. I, 381 (1842)	—	—	—	—	—	Q.	N.A.	B.fl.II,391	M.fr.XI,67,68.		
A. retivenea; F. v. M., fragm. III, 128 (1863)	—	S.A.	—	—	—	—	N.A.	B.fl.II,392	M.fr.III,128.		
VIII. Juliferae.											
A. triptera; Bentham in Hook. Lond. Journ. I, 325 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,323			
A. rhigiophylla; F. v. M. in Schlecht. Linnaea XXVI, 611 (1853)	—	S.A.	—	—	—	—	—	B.fl.II,333			
A. Oxycedrus; Sieber in De Candolle, prodr. II, 453 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,334			
A. verticillata; Willdenow, spec. plant. IV, 1049 (1805)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,334			
A. Riceana; Henslow in Maund, Botanist III, t. 135 (1839)	—	—	T.	—	—	—	—	B.fl.II,335			

A. amentifera; F. v. M. in Journ. Linn. Soc. III, 141 (1858)	—	—	—	—	—	—	—	—	N.A.	B.fl.II,392	
A. Wickhami; Bentham in Hook. Lond. Journ. I, 379 (1842)	—	—	—	—	—	—	—	—	N.A.	B.fl.II,392	
A. lysiphloia; F. v. M. in Journ. Linn. Soc. III, 137 (1858)	—	S.A.	—	—	—	—	Q.	N.A.	B.fl.II,393		
A. craspedocarpa; F. v. M. in Melb. Chem. new ser. II, 73 (1887)	W.A.	—	—	—	—	—	—	—	—		
A. linarioides; Bentham in Hook. Lond. Journ. I, 371 (1842)	—	—	—	—	—	—	—	N.A.	B.fl.II.393		
A. stipuligera; F. v. M. in Journ. Linn. Soc. III, 144 (1858)	—	S.A.	—	—	—	—	Q.	N.A.	B.fl.II,393		
A. ptychophylla; F. v. M. in Journ. Linn. Soc. III, 142 (1858)	—	—	—	—	—	—	—	N.A.	B.fl.II,394		
A. stigmatophylla; Cunningham in Hook. Lond. Journ. I, 377 (1842)	—	—	—	—	—	—	—	N.A.	B.fl.II.394		
A. acradenia; F. v. M. in Journ. Linn. Soc. III (1858)	—	—	—	—	—	—	Q.	N.A.	B.fl.II,394		
A. leptophleba; F. v. M. in Journ. Linn. Soc. III, 143 (1858)	—	—	—	—	—	—	—	N.A.	B.fl.II,395		
A. limbata; F. v. M. in Journ. Linn. Soc. III, 145 (1858)	—	—	—	—	—	—	—	N.A.	B.fl.II,395		
A. brevifolia; Bentham, Fl. Austr. II, 395 (1864)	—	—	—	—	—	—	Q.	N.A.	B.fl.II,395		
A. megalantha; F. v. M. in Journ. Linn. Soc. III, 143 (1858)	—	—	—	—	—	—	—	N.A.	B.fl.II,395		
A. gonoclada; F. v. M. in Journ. Linn. Soc. III, 140 (1858)	—	—	—	—	—	—	Q.	N.A.	B.fl.II,396		
A. pycnostachya; F. v. M., pl. of Vict. II, 33 (1860)	—	—	—	—	N.S.W.	—	—		B.fl.II,396		
A. subtilinervis; F. v. M., pl. of Vict. II, 32 (1860)	—	—	—	V.	N.S.W.	—	—		B.fl.II,396	M.fr.IV,8.	
A. cochliocarpa; Meissner in Bot. Zeitung 10 (1855)	W.A.	—	—	—	—	—	—		B.fl.II,397		
A. Dallachiana; F. v. M., fragm. I, 7 (1858)	—	—	—	V.	—	—	—		B.fl.II,397	M.fr.I,7.	
A. alpina; F. v. M., fragm. III, 129 (1863)	—	—	—	V.	—	—	—		B.fl.II.397	M.fr.III,129.	
A. longifolia; Willdenow, spec. pl. IV, 1052 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—		B.fl.II,397		
A. linearis; Sims, Bot. Mag. t. 2156 (1820)	—	—	T.	V.	N.S.W.	—	—		B.fl.II,399		
A. aciphylla; Bentham in Schlecht. Linnaea, XXVI, 627 (1853)	W.A.	—	—	—	—	—	—		B.fl.II,399		
A. ephedroides; Bentham in Hook. Lond. Journ. I, 370 (1842)	W.A.	—	—	—	—	—	—		B.fl.II.399		
A. Burkittii; F. v. M. in Benth. Fl. Austr. II, 400 (1864)	—	S.A.	—	—	N.S.W.	—	—		B.fl.II,400		
A. microneura; Meissner in Lehm. pl. Preiss. I, 19 (1844)	W.A.	—	—	—	—	—	—		B.fl.II,400		
A. cyperophylla; F. v. M. in Benth. Fl. Austr. II, 400 (1864)	—	S.A.	—	—	N.S.W.	Q.	—		B.fl.II,400		
A. multispicata; Bentham, Fl. Austr. II, 400 (1864)	W.A.	—	—	—	—	—	—		B.fl.II,400		
A. pityoides; F. v. M. in Journ. Linn. Soc. III, 135 (1858)	—	—	—	—	—	Q.	N.A.		B.fl.II,400		
A. xylocarpa; Cunningham in Hook. Lond. Journ. I, 370 (1842)	—	—	—	—	—	—	N.A.		B.fl.II,401		
A. gonocarpa; F. v. M. in Journ. Linn. Soc. III, 136 (1858)	—	—	—	—	—	—	N.A.		B.fl.II,401		
A. oncinophylla; Lindley, Bot. Regist. XXV, App. XV (1839)	W.A.	—	—	—	—	—	—		B.fl.II,401		
A. conjunctifolia; F. v. M., fragm. XI, 68 (1879)	—	—	—	—	—	—	N.A.		—	M.fr.XI,68.	
A. drepanocarpa; F. v. M. in Journ. Linn. Soc. III, 137 (1858)	—	—	—	—	—	—	N.A.		B.fl.II,402		
A. arida; Bentham in Hook. Lond. Journ. I, 370 (1842)	—	—	—	—	—	—	N.A.		B.fl.II,402		
A. aneura; F. v. M. in Schlecht. Linnaea XXVI, 627 (1853)	W.A.	S.A.	—	V.	N.S.W.	Q.	—		B.fl.II,402	M.fr.IV,8.	
A. cibaria; F. v. M. in Melb. Chemist, July (1882)	W.A.	S.A.	—	—	N.S.W.	Q.	—		—		
A. Kempeana; F. v. M. in Melb. Chemist, July (1882)	—	S.A.	—	—	N.S.W.	Q.	—		—		
A. conspersa; F. v. M. in Journ. Linn. Soc. III, 140 (1858)	—	—	—	—	—	—	N.A.		B.fl.II,403		
A. Doratoxylon; Cunningham in Field's N.S.W. 345 (1825)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.		B.fl.II,403		
A. acuminata; Bentham in Hook. Lond. Journ. I, 373 (1862)	W.A.	—	—	—	—	—	—		B.fl.II,404	M.fr.IV,7.	
A. stereophylla; Meissner in Lehm. pl. Preiss. II, 203 (1847)	W.A.	—	—	—	—	—	—		B.fl.II,404		
A. signata; F. v. M., fragm. IV, 7 (1864)	W.A.	—	—	—	—	—	—		B.fl.II,404	M.fr.IV,7.	
A. delibrata; Cunningham in Hook. Lond. Journ. I, 374 (1842)	—	—	—	—	—	Q.	N.A.		B.fl.II,404		
A. oligoneura; F. v. M. in Journ. Linn. Soc. III, 139 (1858)	—	—	—	—	—	—	N.A.		B.fl.II,405		
A. torulosa; Bentham in Journ. Linn. Soc. III, 139 (1858)	—	—	—	—	—	Q.	N.A.		B.fl.II,405		
A. julifera; Bentham in Hook. Lond. Journ. I, 374 (1842)	—	—	—	—	—	Q.	—		B.fl.II,405		
A. Solandri; Bentham, Fl. Austr. II, 406 (1864)	—	—	—	—	—	Q.	—		B.fl.II,406		
A. leptostachya; Bentham, Fl. Austr. II, 406 (1864)	—	—	—	—	—	Q.	N.A.		B.fl.II,406		
A. glaucescens; Willdenow, spec. plant. IV, 1052 (1805)	—	—	—	V.	N.S.W.	Q.	—		B.fl.II,406		
A. Cunninghamii; Hooker, icon. plant. t. 165 (1837)	—	—	—	—	N.S.W.	Q.	—		B.fl.II,407		

A. leptocarpa; Cunningham in Hook. Lond. Journ. I, 376 (1842)			—	—	—	—	—	Q.	—	B.fl.II,407	
A. polystachya; Cunningham in Hook. Lond. Journ. I, 376 (1842)			—	—	—	—	—	Q.	N.A.	B.fl.II,407	
A. holcocarpa; Bentham, Fl. Austr. II, 408 (1864) ...			—	—	—	—	—	Q.	—	B.fl.II,408	M.fr.XI,69.
A. plectocarpa; Cunningham in Hook. Lond. Journ. I, 375 (1842)			—	—	—	—	—	Q.	N.A.	B.fl.II,408	
A. pachycarpa; F. v. M. in Journ. Linn. Soc. III, 139 (1858)			—	—	—	—	—	—	N.A.	B.fl.II,408	
A. tumida; F. v. M. in Journ. Linn. Soc. III, 144 (1858) ...			—	—	—	—	—	—	N.A.	B.fl.II,409	
A. loxocarpa; Bentham in Hook. Lond. Journ. I, 377 (1842)			—	—	—	—	—	—	N.A.	B.fl.II,409	
A. oncinocarpa; Bentham in Hook. Lond. Journ. I, 378 (1842)			—	—	—	—	—	—	N.A.	B.fl.II,409	
A. retinervis; Bentham in Hook. Lond. Journ. I, 379 (1842)			—	—	—	—	—	—	N.A.	B.fl.II,410	
A. aulacocarpa; Cunningham in Hook. Lond. Journ. I, 378 (1842)			—	—	—	—	—	Q.	N.A.	B.fl.II,410	M.fr.XI,69.
A. calyculata; Cunningham in Hook. Lond. Journ. I, 379 (1842)...			—	—	—	—	—	Q.	—	B.fl.II,410	
A. auriculiformis; Cunningham in Hook. Lond. Journ. I, 377 (1842)			—	—	—	—	—	Q.	N.A.	B.fl.II,411	
A. latifolia; Bentham in Hook. Lond. Journ. I, 382 (1842)			—	—	—	—	—	Q.	N.A.	B.fl.II,411	
A. Mangium; Willdenow, spec. plant. IV, 1053 (1805)			—	—	—	—	—	Q.	—	—	M.fr.XI,36 ... As
A. holosericea; Cunningham in G. Don, gen. syst. II, 407 (1832)...			—	—	—	—	—	Q.	N.A.	B.fl.II,411	... ... P.
A. cincinnata; F. v. M., fragm. XI, 35 (1878)			—	—	—	—	—	Q.	—	—	M.fr.XI,35.
A. dimidiata; Bentham in Hook. Lond. Journ. I, 381 (1842)			—	—	—	—	—	—	N.A.	B.fl.II,412	
A. humifusa; A. Cunningham in Lond. Journ. I, 382 (1842)			—	—	—	—	—	Q.	N.A.	B.fl.II,412	
A. denticulosa; F. v. M., fragm. X, 32 (1876)		W.A.	—	—	—	—	—	—	—	—	M.fr.X,32.
IX. Bipinnatae.											
A. Drummondii; Lindley, Bot. Regist. XXV, App. XV (1839)		W.A.	—	—	—	—	—	—	—	B.fl.II,419	
A. pulchella; R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813)		W.A.	—	—	—	—	—	—	—	B.fl.II,416	
A. Mitchelli; Bentham in Hook. Lond. Journ. I, 387 (1842)		—	S.A.	—	V.	—	—	—	—	B.fl.II,417	
A. pentadenia; Lindley, Bot. Regist. t. 1521 (1832)...		W.A.	—	—	—	—	—	—	—	B.fl.II,417	
A. Gilberti; Meissner in Lehm. pl. Preiss. II, 204 (1847)		W.A.	—	—	—	—	—	—	—	B.fl.II,417	
A. nigricans; R. Brown in Ait. hort. Kew. sec. ed. V, 465 (1813)		W.A.	—	—	—	—	—	—	—	B.fl.II,418	
A. Tayloriana; F. v. M. in Wing's S. Sc. Record, July (1882)		W.A.	—	—	—	—	—	—	—	—	
A. strigosa; Link, enum. pl. hort. Berol. 444 (1822)		W.A.	—	—	—	—	—	—	—	B.fl.II,418	
A. elata; Cunningham in Hook. Lond. Journ. I, 383 (1842)		—	—	—	—	N.S.W.	—	—	—	B.fl.II,413	
A. pruinosa; Cunningham in Hook. Lond. Journ. I, 383 (1842)		—	—	—	—	N.S.W.	—	—	—	B.fl.II,413	
A. spectabilis; Cunningham in Hook. Lond. Journ. I, 383 (1842)...		—	—	—	—	N.S.W.	Q.	—	—	B.fl.II,413	
A. Baileyana; F. v. M. in Transact. Roy. Soc. of Vict. XXIV, 168 (1887)		—	—	—	—	N.S.W.	—	—	—	—	
A. polybotrya; Bentham in Hook. Lond. Journ. I, 384 (1842)		—	—	—	—	N.S.W.	Q.	—	—	B.fl.II,414	
A. discolor; Willdenow, spec. plant. IV, 1068 (1805)		—	—	T.	V.	N.S.W.	—	—	—	B.fl.II,414	
A. decurrens; Willdenow, spec. plant. IV, 1872 (1805)		—	—	—	V.	N.S.W.	Q.	—	—	B.fl.II,414	
A. mollissima; Willdenow, enum. pl. hort. Berol. 1053 (1809)		—	S.A.	T.	V.	N.S.W.	—	—	—	—	
A. dealbata; Link, enum. pl. hort. Berol. 445 (1822)		—	—	—	V.	N.S.W.	Q.	—	—	B.fl.II,415	
A. cardiophylla; Cunningham in Hook. Lond. Journ. I, 385 (1842)		—	—	—	—	N.S.W.	—	—	—	B.fl.II,415	
A. leptoclada; Cunningham in Hook. Lond. Journ. I, 385 (1842)...		—	—	—	—	N.S.W.	Q.	—	—	B.fl.II,416	
A. pubescens; R. Brown in Ait. hort. Kew. sec. ed. V, 464 (1813)		—	—	—	—	N.S.W.	—	—	—	B.fl.II,416	
A. Farnesiana; Willdenow, spec. plant. IV, 1083 (1805)		W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	—	B.fl.II,419	... ... E., As., Afr., Am., P.
A. suberosa; Cunningham in Hook. Lond. Journ. I, 499 (1842)		—	—	—	—	—	—	N.A.	—	B.fl.II,420	M.fr.XI,36.
A. Bidwilli; Bentham in Schlecht. Linnaea XXVI, 629 (1853)		—	—	—	—	—	Q.	N.A.	—	B.fl.II,420	
A. pallida; F. v. M. in Journ. Linn. Soc. III, 14 (1858)		—	—	—	—	—	—	N.A.	—	B.fl.II,421	
A. Sutherlandi; F. v. M., inconogr. of Austr. Acac. twelfth decade (1888)		—	—	—	—	—	Q.	—	—	—	M.fr.VI,22; IX,179.
ALBIZZIA; Durazzini in Magazzino Toscano III, part IV, 13 (1772). (Zygia 1756, Pithecolobium, Cathormion, Calliandra, Enterolobium, Serianthes, Archidendron.)											
A. lophantha; Bentham in Hook. Lond. Journ. III, 86 (1844)		W.A.	—	—	—	—	—	—	—	B.fl.II,421	
A. basaltica; Bentham, Fl. Austr. II, 422 (1864)		—	—	—	—	—	Q.	—	—	B.fl.II,422	
A. Thozetiana; F. v. M. in Benth. Fl. Austr. II, 422 (1864)		—	—	—	—	—	Q.	—	—	B.fl.II,422	M.fr.IV,9&176.

A. procera; Bentham in Hook. Lond. Journ. III, 88 (1844)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.II,422	...	...	As.	
A. pruinosa; F. v. M. in Trimen's Journ. of Bot. 9 (1872)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,423	M.fr.IX,179	...	As.	
A. Lebbek; Durazzini in Magaz. Tosc. III, pt. IV, 13 (1772)	...	...	...	—	—	—	—	—	Q.	—	—	...	...	As., Afr.	
A. canescens; Bentham, Fl. Austr. II, 423 (1864)	...	...	...	...	...	—	—	—	—	Q.	—	B.fl.II,423	M.fr.IX,179.		
A. Tozeri; F. v. M. in Trimen's Journ. of Bot. 10 (1872)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.II,424	M.fr.V,106; IX,179.		
A. Hendersoni; F. v. M. in Trimen's Journ. of Bot. 10 (1872)	...	...	...	...	...	—	—	—	N.S.W.	Q.	—	—	M.fr.V,191; VI,202; IX,178.		
A. monilifera; F. v. M. in Trimen's Journ. of Bot. 10 (1872)	...	...	...	...	...	—	—	—	—	—	N.A.	B.fl.II,424	M.fr.V,191	...	.. As.
A. ramiflora; F. v. M., fragm. IX, 178 (1875)	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.V,10; IX,178.		
A. Vaillantii; F. v. M., fragm. V, 9 (1865)	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.V,60; IX,178.		
A. Lucyi; F. v. M., iconogr. of Austr. Acai. 13, Dec. 6 (1888)	...	...	...	...	...	—	—	—	—	Q.	—	—	M.fr.V,201; IX,178.		

CONNARACEAE.

R. Brown, Narr. Exped. Cong. App. V, 431 (1818).—Genera 2, Species 2.

ROUREA; Aublet, Hist. des pl. de la Guian. I, 467, t. 187 (1775).

R. brachyandra; F. v. M., fragm. VIII, 6 (1872)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.X,119.	

TRICHOLOBUS; Blume, Mus. Bot. Lugd. I, 237 (1850).

T. connaroides; F. v. M., fragm. VIII, 224 (1874)	..	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.V,105; X,120.	

ROSACEAE.

A. L. de Jussieu, Gen. pl. 334 (1789), from B. de Jussieu (1759).—Genera 8, Species 17.

PARINARIUM; A. L. de Jussieu, gen. pl. 342 (1789).

P. Nonda; F. v. M. in Benth. Fl. Austr. II, 426 (1864)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,426			
P. Griffithianum; Bentham in Hook. Niger-Flora 334 (1849)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.II,426	M.fr.VIII,34	...	As., P.	

STYLOBASIUM; Desfontaines in Mém. du Mus. V, 37, t. 2 (1819). (Macrostigma.)

S. spathulatum; Desfontaines in Mém. du Mus. V, 37, t. 2 (1819)	...	...	W.A.	S.A.	—	—	—	—	N.A.	B.fl.II,427					
S. lineare; Nees in Lehm. pl. Preiss. I, 95 (1844)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,427		

GEUM; Linné, gen. plant. 148 (1737) from Plinius. (Sieversia.)

G. urbanum; Linné, spec. plant. 501 (1753)	...	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,428	...	...	...	E., As.
G. renifolium; F. v. M. in Transact. phil. Inst. Vict. II, 66 (1857)	...	...	...	—	—	T.	—	—	—	—	B.fl.II,428				

POTENTILLA; Linné, gen. plant. 147 (1737) from Camerarius (1586).

P. anserina; Linné, spec. plant. 495 (1753)	...	...	...	...	...	...	—	S.A.	T.	V.	—	—	—	B.fl.II,429	...	...	...	E., As., Af., Am., N.Z.

RUBUS; Tournefort, Inst. 614, t. 385 (1700) from Plinius.

R. Gunnianus; Hooker, icon. plant. t. 291 (1840)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.II,430	M.fr.IV,32			
R. parvifolius; Linné, spec. plant. 1197 (1753)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,430	M.fr.IV,30	...	As.
R. rosifolius; Smith, plantar. icon. ex herb. Linn. t. 60 (1790)	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,431	M.fr.IV,32; VIII,34.	As., Afr.		
R. Moluccanus; Linné, spec. plant. 1197 (1753)	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.II,430	M.fr.IV,31; VIII,34.	As., P.	
R. Moorei; F. v. M. in Transact. phil. Inst. Vict. II, 67 (1857)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,431	M.fr.IV,29&176; VIII,34.			

AGRIMONIA; Tournefort, inst. 301, t. 155 (1700) from C. Bauhin (1623).

A. Eupatoria; Linné, spec. plant. 448 (1753)	...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,34	...	As., Afr., Am.

ALCHEMILLA; Linné, gen. pl. 30 (1737) from C. Bauhin (1623). (Alchimilla.)

A. vulgaris; C. Bauhin, pinax 319 (1623)	...	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.II,432	...	...	...	E., As., Am.

ACAENA; Mutis in Linné, mantiss. II, 200 (1771). (Ancistrum.)

A. ovina; Cunningham in Field's N. S. Wales, 358 (1825)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,433						Am., N.Z.
A. sanguisorbae; Vahl, enum. plant. I, 294 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,434						N.Z.
A. montana; J. Hooker in Hook. Lond. Journ. VI, 276 (1847)	—	—	T.	—	—	—	—	B.fl.II,434						

SAXIFRAGEAE.

VENTENAT, Tabl. III, 277 (1799).—Genera 22, Species 36.

ARGOPHYLLUM; R. et G. Forster, charact. gen. 29, t. 15 (1776).

A. Lejourdanii; F. v. M., fragm. IV, 33 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,437	M.fr.IV,177; VI,188.					

ABROPHYLLUM; J. Hooker in Benth. Fl. Aust. II, 437 (1864). (Brachynema.)

A. ornans; J. Hooker in Benth. Fl. Austr. II, 437 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,437	M.fr.III,90; VI,189; VII,150.					

CUTTSIA; F. v. M., fragm. V, 47 (1865).

C. viburnea; F. v. M., fragm. V, 47 (1865)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,47&189; VI,189.					

COLMEIROA; F. v. M., fragm. VII, 149 (1871).

C. carpodetoides; F. v. M., fragm. VII, 149 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,149.					

QUINTINIA; A. de Candolle, Mon. Campan. 92 (1830).

Q. Sieberi; A. De Candolle, Monogr. Camp. 92 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,438	M.fr.II,126; III,165; VI,189.					
Q. Verdonii; F. v. M., fragm. II, 125 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,438	M.fr.VI,189.					
Q. Fawkneri; F. v. M., fragm. VI, 92 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VII,150.					

POLYOSMA; Blume, Bijdr. 658 (1825).

P. Cunninghamii; J. J. Bennett, pl. Jav. rar. 196 (1840)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,438	M.fr.VI,189.					
P. alangiacea; F. v. M., fragm. VIII, 8 (1872)	—	—	—	—	—	Q.	—	—	M.fr.VIII,8.					

ANOPTERUS; Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804).

A. glandulosus; Labillardière, Nov. Holl. pl. spec. I, 86, t. 112 (1804)	—	—	T.	—	—	—	—	B.fl.II,439	M.fr.II,126.					
A. Macleayanus; F. v. M. in Journ. Pharm. Soc. Vict. 43 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,439	M.fr.VI,188; VII,150.					

CALLICOMA; Andrews, Bot. Reposit. t. 566 (1809). (Calycomis partly.)

C. serratifolia; Andrews, Bot. Reposit. t. 566 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,440	M.fr.V,32.					
C. Stutzeri; F. v. M., fragm. V, 34 (1865)	—	—	—	—	—	Q.	—	—	M.fr.VI,188&252; VII,150.					

ANODOPETALUM; Cunningham in Endl. gen. 818 (1839).

A. biglandulosum; Cunningham in J. Hook. Fl. Tasm. I, 148 (1860)	—	—	T.	—	—	—	—	B.fl.II,440	M.fr.VI,189.					

APHANOPETALUM; Endlicher, nov. stirp. decad. I, 34 (1839). (Platyptelea.)

A. resinosum; Endlicher, nov. stirp. dec. I, 35 (1839)	—	—	—	V.	N.S.W.	Q.	—	B.fl.II,441	M.fr.I,228.					
A. occidentale; F. v. M. fragm. I, 228 (1858)	W.A.	—	—	—	—	—	—	B.fl.II,441	M.fr.XI,20.					

CERATOPETALUM; Smith, specim. of the Bot. of New Holland, 9 (1793).

C. gummiferum; Smith, specim. of the Bot. of New Holl. t. 3 (1793)	—	—	—	—	N.S.W.	—	—	B.fl.II,442	M.fr.II,126.					
C. apetalum; D. Don in Edinb. new phil. journ. (1830)	—	—	—	—	N.S.W.	—	—	B.fl.II,442	M.fr.II,126; VI,189; VII,150.					

SCHIZOMERIA; D. Don in Edinb. new phil. journ. IX, 94 (1830).

S. ovata; D. Don in Edinb. new phil. journ. IX, 94 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,443	M.fr.VI,189.					

ACROPHYLLUM; Bentham in Maund & Henslow, Botanist II, 95 (1840). (Calycomis partly.)

A. venosum; Bentham in Maund & Henslow, Bot. II, 95 (1840)	—	—	—	—	N.S.W.	—	—	B.fl.II,443	M.fr.VI,189; VII,150.					

WEINMANNIA; Linné, syst. ed. X, 1005 (1759). (Ackama, Geissois.)

W. paniculosa; F. v. M., fragm. II, 126 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,444	M.fr.II,83,175; VI,188; VII,150.					

W. Biagiana; F. v. M., fragm. V, 16 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr.VII,150.						
W. lachnocarpa; F. v. M., fragm. VIII, 7 (1782) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VIII,7,281.						
W. rubifolia; Bentham, Fl. Aust. II, 445 (1864) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,445	M.fr.II,82&175; V,180; VI,181.						
W. Benthami; F. v. M., Docum. intercol. Exhibit. 31 (1866) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,446	M.fr.V,180; VI,188.						
SPIRAEANTHEMUM; Asa Gray in proceed. Amer. Acad. of Science III (1853).															
S. Davidsoni; F. v. M. in Austr. Journ. of Pharm. new ser. II, 85 (1887) ...	—	—	—	—	—	Q.	—	—							
GILLBEEA; F. v. M., fragm. V, 17 (1865).															
G. adenopetala; F. v. M., fragm. V, 17 (1865) ...	—	—	—	—	—	Q.	—	—	M.fr.VI,188.						
DAVIDSONIA; F. v. M., fragm. VI, 4 (1867).															
D. pruriens; F. v. M., fragm. VI, 4 (1867) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr,VI,249.						
TETRACARPAEA; J. Hooker, icon. plant. t. 264 (1840).															
T. Tasmanica; J. Hooker, icon. plant. t. 264 (1840)...	—	—	T.	—	—	—	—	B.fl.II,445							
EUCRYPHIA; Cavanilles, icon. IV, 48, t. 372 (1797). (Carpodontos.)															
E. Billardieri; Spach, veg. phan. V, 344 (1836) ...	—	—	T.	—	—	—	—	B.fl.II,446	M.fr.IV,2; VI,189; VII,150.						
E. Moorei; F. v. M., fragm. IV, 2 (1864) ...	—	—	—	V.	N.S.W.	—	—	B.fl.II,447	M.fr.IV,176; VI,189; VII,150.						
BAUERA; Banks & Kennedy in Andrews, Bot. Rep. t. 198 (1793).															
B. rubioides; Andrews, Bot. Reposit. t. 198 (1793) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,447	M.fr.IV,23.						
B. capitata; Séringe in De Candolle, prodr. IV, 13 (1830) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,448	M.fr.IV,24.						
B. sessiliflora; F. v. M. in Transact. phil. Soc. Vict. I, 41 (1854)...	—	—	—	V.	—	—	—	B.fl.II,448	M.fr.IV,24; VII,150.						
EREMOSYNE; Endlicher in Hueg. enum. pl. Nov. Holl. Austr. occ. 53 (1837).															
E. pectinata; Endlicher in Hueg. enum. pl. 53 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.II,449	M.fr.VI,189.						
CEPHALOTUS; Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806).															
C. follicularis; Labillardière, Nov. Holl. pl. spec. II, 7, t. 145 (1806) ...	W.A.	—	—	—	—	—	—	B.fl.II,449	M.fr.VI,189.						

NEPENTHACEAE.

Reichenbach, Conspectus regni vegetabilis 45 (1828).—Genus 1, Species 2.

NEPENTHES; Linné, syst. nat. 9 (1735); Linné, gen. pl. 273 (1737).															
N. Kennedyana; F. v. M., fragm. V, 154 (1866) ...	—	—	—	—	—	Q.	—	B.fl.VI,40	M.fr.V,154.						
N. phyllamphora; Willdenow, sp. plant. IV, 874 (1805) ...	—	—	—	—	—	Q.	—	—	...	...	...	As., P.			

ARISTOLOCHIEAE.

A. L. de Jussieu in Ann. du Mus. V, 221 (1804).—Genus 1, Species 5.

ARISTOLOCHIA; Tournefort, inst. 162, t. 71 (1700), from Hippocrates, Theophrastos & Dioscorides.															
A. deltantha; F. v. M., fragm. VI, 179 (1869) ...	—	—	—	—	—	Q.	—	B.fl.VI,208	M.fr.VI,179&255.						
A. praevenosa; F. v. M., fragm. II, 166 (1861) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,208	M.fr.VI,180.						
A. pubera; R. Brown, prodr. 349 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,208							
A. Thozetii; F. v. M., fragm. II, 167 (1861) ...	—	—	—	—	—	Q.	—	B.fl.VI,208	M.fr.II,167.						
A. Indica; Linné, spec. plant. 960 (1753) ...	—	—	—	—	—	Q.	—	B.fl.VI,209	...	...	As., P.				

CRASSULACEAE.

De Candolle, Bull. de la Soc. Philom. n. 49 (1801).—Genus 1, Species 6.

TILLAEA; Micheli, nov. gen. 22, t. 20 (1729). (Bulliarda, Crassulae subgenus.)															
T. verticillaris; De Candolle, prodr. III, 382 (1828)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,451	M.fr.XI,118	...	Am., N.Z.				
T. purpurata; J. Hooker in Lond. Journ. of Bot. VI, 472 (1847)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,451	M.fr.XI,118.						

T. macrantha; J. Hooker in Hook. icon. plant. t. 310 (1841)				S.A.	T.	V.	N.S.W.	—	—	B.fl.II,452	M.fr.XI,117.				
T. pedicellosa; F. v. M., fragm. XI, 118 (1881)				W.A.	—	—	—	—	—	—	M.fr.XI,118.				
T. intricata; Nees in Lehm. pl. Preiss. I, 278 (1844)				W.A.	—	—	—	—	—	—	M.fr.XI,117.				
T. recurva; J. Hooker, Fl. Tasm. I, 146 (1860)				W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,452	M.fr.XI,118.			

HAMAMELIDEAE.

R. Brown in Abel, Narr. journ. Chin. 374 (1818).

Genus and species undertermined								—	—	—	—	—	Q.	—	—

ONAGREAE.

Adanson, Familles des plantes II, 81 (1763), from B. de Jussieu (1759).—Genera 4, Species 5.

OENOTHERA; Linné, syst. nat. 8 (1735); Linné, gen. plant. 112 (1737).

O. Tasmanica; J. Hooker, Fl. Tasm. I, 119 (1860)								—	—	T.	—	—	—	—	B.fl.III,303

EPILOBIUM; Dillenius in Linné, syst. nat. 8 (1735), from Gesner (1542).

E. glabellum; G. Forster, florul. ins. Austr. prodr. 27 (1786)				W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,305				N.Z.

JUSSIEUA; Linné, gen. plant. 126 (1737). (Jussiaea.)

J. diffusa; Forskael, fl. Aeg. Arab. 210 (1775)						—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,306		As., Afr.
J. suffrnticosa; Linné, spec. plant. 388 (1753)						—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,307	M.fr.III,13	As., Afr., Am., P.

LUDWIGIA; Linné, coroll. 3 (1737).

L. parviflora; Roxburgh, Fl. Ind. edit. Carey. I, 440 (1820)						—	—	—	—	—	Q.	N.A.	B.fl.III,307	M.fr.III,129; IV,175	As., Afr.

SALICARIEAE.

Adanson, Familles des plantes II, 232 (1763), from B. de Jussieu (1759).—Genera 7, Species 19.

ROTALA; Linné, mantissa altera 143 (1771).

R. verticillaris; Linné, mantissa altera 143 (1771)						—	—	—	—	—	—	N.A.	B.fl.III,295	M.fr.III,108	As.
R. Roxburghiana; Wight, Icon. t. 260 (1840)						—	—	—	—	—	Q.	N.A.	B.fl.III,296	M.fr.XI,118	As.
R. Mexicana; Chamisso & Schlechtendal in Linnaea V, 567 (1830)						—	—	—	—	—	Q.	N.A.	—		As., Am.
R. occultiflora; Koehne in Engler's Jahrb. I, 152 (1880)						—	—	—	—	—	Q.	N.A.	—		
R. diandra; F. v. M. in Transact. Roy. Soc. of N.S.W. 89 (1880)						—	S.A.	—	—	—	Q.	N.A.	B.fl.III,296	M.fr.III,108,168.	

AMMANNIA; Houston in Linné, gen. plant. 337 (1737). (Koehnea, Crinopodium.)

A. crinipes; F. v. M. in Transact. phil. Soc. Vict. III, 49 (1858)						—	—	—	—	—	—	N.A.	B.fl.III,296		
A. triflora; R. Brown in Benth. Fl. Austr. III, 297 (1866)						—	—	—	—	—	Q.	N.A.	B.fl.III,297		
A. baccifera; Linné, spec. plant. ed. sec. 175 (1762)						—	S.A.	—	—	—	Q.	N.A.	B.fl.III,297		As., Afr.
A. auriculata; Willdenow, hort. Berol. t. 7 (1806)						—	—	—	—	—	Q.	—	B.fl.III,297		As., Afr.
A. multiflora; Roxburgh, Fl. Ind. edit. Carey. I, 447 (1820)				W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,298				As., Afr.

LAGERSTROEMIA; Linné, syst. ed. X, 1076 (1759).

L. Indica; Linné, spec. plant. ed. sec. 734 (1762)						—	—	—	—	—	Q.	—	—	M.fr.VIII,35	As.
L. Flos Reginae; Retzius, observ. botan. V, 25 (1789)						—	—	—	—	—	Q.	—	—	M.fr.XII,20	As.
L. Archeriana; Bailey, Synop. of Queensland Flora 196 (1883)						—	—	—	—	—	Q.	—	—		

LYTHRUM; Linné, syst. nat. 8 (1735); Linné, gen. plant. 136 (1737).

L. Salicaria; Linné, spec. plant. 446 (1753)						—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,298	M.fr.III,110; VII,147	E., As., Afr., Am.
L. Hyssopifolia; Linné, spec. plant. 447 (1753)						—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,299	M.fr.III,109; VII,147	E., As., Afr., Am.

NESAEA; Commerçon in A. L. de Jussieu, gen. plant. 332 (1789). (Decadon.)

N. Robertsii; F. v. M., fragm. VII, 145 (1871)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VII,145.	
N. Arnhemica; F. v. M., fragm. VII, 146 (1871)	..	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,299	M.fr.III,109,168.	

PEMPHIS; R. et G. Forster, charact. gen. 67, t. 34 (1776).

P. acidula; R. & G. Forster, charact. gen. 67, t. 34 (1776)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,300	M.fr.III,109; VII,147		As., Afr., P.

LAWSONIA; Linné, gen. plant. 111 (1737).

L. alba; Lamarck, Encycl. méth. III, 106 (1789)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,301	...	...	As., Afr.

HALORAGEAE.

R. Brown in Flind. Voy. II, 549 (1814).—Genera 4, Species 58.

GUNNERA; Linné, mantiss. plant. 16 (1767). (Milligania.)

G. cordifolia; J. Hooker, Fl. Tas. I, 125 (1860)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.II,491		

LOUDONIA; Lindley, Bot. Regist. XXV, App. XLII (1839). (Glischrocaryon.)

L. aurea; Lindley, Bot. Regist. XXV, App. XLII (1839)	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.II,472	M.fr.VIII,162; XI,20.		
L. Behrii; Schlechtendal, Linnaea XX, 648 (1847)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,472	M.fr.VIII,162.	
L. Roei; Schlechtendal, Linnaea XX, 648 (1847)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,472	M.fr.VIII,162.	

HALORAGIS; R. et G. Forster, charact. gen. 61, t. 31 (1776). (Cercodia, Goniocarpus, Meionectes.)

H. Gossei; F. v. M., fragm. VIII, 161 (1874)	...	...	...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	—	M.fr.XI,134.	
H. trigonocarpa; F. v. M., fragm. X, 84 (1876)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,84.	
H. digyna; Labillardière, Nov. Holl. pl. spec. I, 101, t. 129 (1804)	...	...	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.II,475		
H. mucronata; Bentham, Fl. Austr. II, 475 (1864)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.II,475		
H. pityoides; Bentham, Fl. Austr. II, 476 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,476		
H. cordigera; Fenzl in Hueg. enum. 45 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,476		
H. elata; Cunningham in Hueg. enum. 45 (1837)	...	..	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.II,476	M.fr.VIII,162.	
H. tenuifolia; Bentham, Fl. Austr. II, 477 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,477		
H. scoparia; Fenzl in Hueg. enum. 45 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,477	M.fr.X,54.	
H. aculeolata; Bentham, Fl. Austr. II, 477 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,477	M.fr.X,54.	
H. foliosa; Bentham, Fl. Austr. II, 477 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,477		
H. pycnostachya; F. v. M. in Transact. R. S. Vict. XXIV, p. 135 (1887)					...	W.A.	—	—	—	—	—	—	—		
H. confertifolia; F. v. M., fragm. X, 53 (1876)	...	...	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,54.	
H. platycarpa; Bentham, Fl. Austr. II, 478 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,478		
H. acutangula; F. v. M. in Transact. Vict. Inst. 125 (1854)	...	...	...		...	—	S.A.	—	—	—	—	—	B.fl.II,478		
H. hexandra; F. v. M., fragm. III, 31 (1862)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,478	M.fr.III,31.	
H. odontocarpa; F. v. M., fragm. I, 108 (1859)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.II,479	M.fr.I,108; X,54.	
H. serra; Brongniart in Duperr. Voy. sur la Coquille, t. 69 (1829)		...	...		...	—	—	—	—	N.S.W.	—	—	B.fl.II,479		
H. aspera; Lindley in Mitch. Trop. Austr. 306 (1848)	...	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,479		
H. exalata; F. v. M. in. Transact. R. Soc. of Vict. XXIV. 132 (1887)		...	...		...	—	—	—	—	N.S.W.	—	—	B.fl.II,479		
H. racemosa; Labillardière, Nov. Holl. pl. spec. I, 100, t. 128 (1804)		...	...		...	W.A.	—	—	—	—	—	—	B.fl.II,480	M.fr.VIII,162.	
H. Bauerlenii; F. v. M. in Transact. R. Soc. Vict. XXIV, p. 132 (1887)					...	—	—	—	V.	N.S.W.	—	—	—		
H. rotundifolia; Bentham, Fl. Austr. II, 480 (1864)	...	...	...		...	W.A.	—	—	—	—	—	—	B.fl.II,480	M.fr.VIII,162.	
H. rudis; Bentham, Fl. Austr. II, 480 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,480		
H. paniculata; R. Brown in Benth. Fl. Austr. II, 481 (1864)		...	...		...	W.A.	—	—	—	—	—	—	B.fl.II,481		
H. pusilla; R. Brown in Benth. Fl. Austr. II, 481 (1864)	...	...	..		...	W.A.	—	—	—	—	—	—	B.fl.II,481		
H. intricata; Bentham, Fl. Austr. II, 481 (1864)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,481		
H. nodulosa; Walpers, repert. V, 671 (1846)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.II,481		
H. trichostachya; Bentham, Fl. Austr. II, 481 (1864)	...	...	...		...	W.A.	—	—	—	—	—	—	B.fl.II,481		

H. lanceolata ; R. Brown in Benth. Fl. Austr. II, 481 (1864)	W.A.	—	—	—	—	—	—	B.fl.II,482			
H. micrantha ; R. Brown in Flind. Voy. II, 550 (1814) implied	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,482	M.fr.VIII,162; X,54	...	As., N.Z.
H. stricta ; R. Brown in Benth. Fl. Austr. II, 482 (1864)	—	—	—	—	—	Q.	—	B.fl.II,482			
H. heterophylla ; Brongniart in Duperr. Voy. t. 68 (1829)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.II,483			
H. leptotheca ; F. v. M., fragm. III, 32 (1862)	—	—	—	—	—	—	N.A.	B.fl.II,483	M.fr.VIII,162.		
H. tetragyna ; R. Brown in Flind. Voy. II, 550 (1814) implied	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,484	M.fr.III,32; IV,26	...	N.Z.
H. teucrioides ; Schlechtendal, Linnaea XX, 648 (1847)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,484			
H. scordioides ; Bentham, Fl. Austr. II, 485 (1864)...	W.A.	—	—	—	—	—	—	B.fl.II,485			
H. depressa ; Walpers, repertor. II, 99 (1843)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.II,485	M.fr.IV,26 ...	...	N.Z.
H. salsoloides ; Bentham, Fl. Austr. II, 485 (1864)	—	—	—	—	N.S.W.	—	—	B.fl.II,485			
H. monosperma ; F. v. M. in proceed. Linn. Soc. N.S.W., X, 197 (1885) ...	—	—	—	V.	N.S.W.	—	—	—			
H. Meionectes ; F. v. M., Key to the syst. of Vict. pl. I, 261 (1888)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,486	M.fr.VIII,163.		
MYRIOPHYLLUM ; l'Ecluse, rar. stirp. hist. II, 252 (1853) from Dioscorides. (Pelonastes).											
M. variifolium ; J. Hooker in Hook. icon. plant. t. 289 (1840)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,487	...	...	N.Z.
M. elatinoides ; Gaudichaud in Ann. des sc. nat. V, 105 (1825)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,487	...	...	Am., N.Z.
M. verrucosum ; Lindley in Mitch. Trop. Austr. 384 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.II,488	M.fr.VIII,163.		
M. latifolium ; F. v. M., fragm. II, 87 (1860)...	—	—	—	—	N.S.W.	Q.	—	B.fl.II,488	M.fr.II,87; IV,172; X,54.		
M. Muelleri ; Sonder in Linnaea XXVIII, 233 (1855)	W.A.	S.A.	—	V.	—	—	—	B.fl.II,488	M.fr.VIII,162.		
M. amphibium ; Labillardière, Nov. Holl. pl. spec. II, 70, t. 220 (1806)... ...	—	S.A.	T.	V.	—	—	—	B.fl.II,489	M.fr.VIII,163.		
M. pedunculatum ; J. Hooker in Lond. Journ. VI, 474 (1847)	W.A.	S.A.	T.	V.	—	—	—	B.fl.II,489	M.fr.VIII,162	...	N.Z.
M. dicoccum ; F. v. M. in Transact. phil. Inst. Vict. III, 41 (1858)	—	—	—	—	—	—	N.A.	B.fl.II,489	M.fr.VIII,162.		
M. trachycarpum ; F. v. M., fragm. II, 87 (1860)	—	—	—	—	—	—	N.A.	B.fl.II,489	M.fr.II,87.		
M. gracile ; Bentham, Fl. Austr. II, 489 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.II,489			
M. filiforme ; Bentham, Fl. Austr. II, 489 (1864)	—	—	—	—	—	—	N.A.	B.fl.II,489			
M. integrifolium ; J. Hooker, Fl. Tasm. I, 123, t. 23 (1860)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.II,490	M.fr.VIII,162.		
M. Drummondii ; J. Hooker, Fl. Tasm. I, 123 implied (1860)	W.A.	—	—	—	—	—	—	B.fl.II,490			

CALLITRICHINAE.

LINK, Enumer. plant. hort. reg. Berol. I, 7 (1821).—Genus 1, Species 2.

CALLITRICHE ; Linné, syst. nat. ed. sext. 82 (1748) from Colonna (1616).											
C. verna ; Linné, flor. Suec. edit. sec. 32 (1755)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.II,492	...	...	E., As., Afr., Am., P., N.Z.
C. Muelleri ; Sonder in Linnaea XXVIII, 229 (1855)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,163.		

RHIZOPHOREAE.

R. BROWN in Flind. Voy. II, 549 (1814).—Genera 4, Species 7.

RHIZOPHORA ; Linné, gen. plant. 137 (1737).											
R. mucronata ; Lamarck, Encycl. méth. VI, 169 (1804)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,493	M.fr.IX,159 ...		As., Afr., P.
CERIOPS ; Arnott in Jardine, Ann. of nat. hist. I, 363 (1838).											
C. Candolleana ; Arnott in Ann. of nat. hist. I, 364 (1838)...	—	—	—	—	—	Q.	N.A.	B.fl.II,494	M.fr.IX,159 ...		As., Afr.
BRUGUIERA ; Lamarck, Encycl. méth. IV, 696, t. 397 (1796).											
B. Rhèedii ; Blume, enum. pl. Jav. 92 (1827)...	—	—	—	—	—	Q.	N.A.	B.fl.II,494	...	...	As., P.
B. caryophylloides ; Blume, enum. pl. Jav. 93 (1827)	—	—	—	—	—	Q.	—	—	M.fr.IX,159 ...		As.
B. parviflora ; Wight & Arnott, prodr. fl. penins. Ind. 311 (1834)	—	—	—	—	—	Q.	—	—	...	...	As.
B. gymnorrhiza ; Lamarck, Encycl. méth. IV, 696, t. 397 (1796)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.II,495	M.fr.IX,159.		As., Afr., P.
CARALLIA ; Roxburgh in Flind. Voy. II, 549 (1814).											
C. integerrima ; De Candolle, prodr. III, 33 (1828)	—	—	—	—	—	Q.	N.A.	B.fl.II,495	M.fr.IX,159.		As.

COMBRETACEAE.

R. Brown, prodr. 351 (1810).—Genera 4, Species 27.

TERMINALIA; Linné, mantiss. plant. 21 (1767). (Chuncoa.)

T. platyptera; F. v. M., fragm. II, 151 (1861)		—	—	—	—	—	Q.	N.A.	B.fl.II,498	M.fr.II,151; IX,160.		
T. volucris; R. Brown in Benth. Fl. Austr. II, 498 (1864)		—	—	—	—	—	—	N.A.	B.fl.II,498			
T. oblongata; F. v. M., fragm. II, 152 (1861)		—	—	—	—	—	Q.	—	B.fl.II,499	M.fr.III,92,167; IX,160.		
T. bursarina; F. v. M., fragm. II, 149 (1861)		—	—	—	—	—	Q.	N.A.	B.fl.II,499	M.fr.II,149.		
T. circumalata; F. v. M., fragm. III, 91 (1862)	W.A.	—	—	—	—	—	—	N.A.	B.fl.II,499	M.fr.III,91.		
T. pterocarya; F. v. M., fragm. II, 152 (1861)		—	—	—	—	—	—	N.A.	B.fl.II,500	M.fr.II,152.		
T. Thozetii; Bentham, Fl. Austr. II, 500 (1864)		—	—	—	—	—	Q.	—	B.fl.II,500	M.fr.IX,160.		
T. Catappa; Linné, mantiss. pl. 128 (1767)		—	—	—	—	—	Q.	—	—	M.fr.IX,160.	...	As., P.
T. melanocarpa; F. v. M., fragm. III, 92 (1862)		—	—	—	—	—	Q.	N.A.	B.fl.II,500	M.fr.III,92; IX,160.		
T. Muelleri; Bentham, Fl. Austr. II, 500 (1864)		—	—	—	—	—	Q.	—	B.fl.II,500	M.fr.IX,160.		
T. latipes; Bentham, Fl. Austr. II, 501 (1864)		—	—	—	—	—	—	N.A.	B.fl.II,501			
T. edulis; F. v. M., fragm. II, 151 (1861)		—	—	—	—	—	—	N.A.	B.fl.II,501	M.fr.II,151; IX,160.		
T. discolor; F. v. M., fragm. III, 92 (1862)		—	—	—	—	—	—	N.A.	B.fl.II,501	M.fr.III,92.		
T. porphyrocarpa; F. v. M. in Benth. Fl. Austr. II, 501 (1864)		—	—	—	—	—	Q.	—	B.fl.II,501			
T. platyphylla; F. v. M., fragm. II, 150 (1861)		—	—	—	—	—	Q.	N.A.	B.fl.II,502	M.fr.II,150.		
T. sericocarpa; F. v. M., fragm. IX, 159 (1875)		—	—	—	—	—	Q.	—	—	M.fr.IX,159.		
T. microcarpa; Decaisne in Ann. du Mus. III (1834)		—	—	—	—	—	—	N.A.	B.fl.II,502	M.fr.II,151; III,92.		As.
T. petiolaris; Cunningham in Benth. Fl. Austr. II, 502 (1864)		—	—	—	—	—	—	N.A.	B.fl.II,502			
T. erythrocarpa; F. v. M., fragm. II, 150 (1861)		—	—	—	—	—	—	N.A.	B.fl.II,503	M.fr.II,150.		
T. grandiflora; Bentham, Fl. Austr. II, 503 (1864)		—	—	—	—	—	Q.	N.A.	B.fl.II,503	M.fr.IX,160.		

LUMNITZERA; Willdenow in den Verh. der Ges. Nat. Freunde zu Berl. IV, 186 (1803).

L. coccinea; Wight & Arnott, prodr. 316 (1834)		—	—	—	—	—	Q.	—	B.fl.II,503	M.fr.IX,160	...	As., P.
L. racemosa; Willdenow in Verh. der Nat. Freunde zu Berl. IV, 186 (1803)		—	—	—	—	—	Q.	N.A.	B.fl.II,504	M.fr.III,92; IX,160.		As., Afr., P.

MACROPTERANTHES; F. v. M., fragm. III, 91 (1862).

M. montana; F. v. M., fragm. III, 91 (1862)		—	—	—	—	—	Q.	—	B.fl.II,504	M.fr.II,149; III,91&166.		
M. Kekwickii; F. v. M., fragm. III, 151 (1863)		—	—	—	—	—	—	N.A.	B.fl.II,504	M.fr.III,151.		
M. Leichhardtii; F. v. M., fragm. III, 91 (1862)		—	—	—	—	—	Q.	—	B.fl.II,505	M.fr.III,91.		
M. Fitzalani; F. v. M., fragm. VIII, 160 (1874)		—	—	—	—	—	Q.	—	—	M.fr.VIII,160.		

GYROCARPUS; N. Jacquin, select. stirp. Amer. hist. 282 (1763).

G. Americanus; N. Jacquin, select. stirp. Amer. hist. 282, t. 178 (1763)		—	—	—	—	—	Q.	N.A.	B.fl.II,505	M.fr.IX,160	...	As., Am., P.

MYRTACEAE.

Adanson, Familles des plantes II, 86 (1763), from B. de Jussieu (1759).—Genera 40, Species 663.

ACTINODIUM; Schauer in Schlecht. Linnaea X, 311 (1835). (Triphelia.)

A. Cunninghamii; Schauer in Lindl. introd. to the nat. syst., sec. ed. 440 (1835)	W.A.	—	—	—	—	—	—	—	B.fl.III,5			

DARWINIA; Rudge in Transact. Linn. Soc. XI, 299 (1813). (Genetyllis, Chamaelaucium, Homoranthus, Hedaroma, Polyzone, Francissia, Decalophium, Schuermannia, Cryptostemon.)

D. macrostegia; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.III,8			
D. Hookeriana; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.III,9	M.fr.VIII,182.		
D. fimbriata; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.III,9	M.fr.VIII,182; IX,177.		
D. speciosa; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.III,9	M.fr.XI,10.		
D. Meissneri; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	—	B.fl.III,9	M.fr.VIII,182; IX,177.		

D. helichrysoides; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,10.	
D. Neildiana; F. v. M., fragm. IX, 177 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,177.
D. oederoides; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,10	M.fr.IX,177; XI,9.
D. virescens; Bentham in Journ. Linn. Soc. IX, 179 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,10	M.fr.XI,10.
D. Oldfieldii; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,10	
D. purpurea; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,11	M.fr.XI,9.
D. citriodora; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,11	M.fr.I,169; IV,174; IX,177; XI,9.
D. thymoides; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,11	M.fr.VIII,182; IX,176.
D. taxifolia; Cunningham in Field's N. S. Wales 352 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.III,12	M.fr.IX,176.
D. vestita; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,12	M.fr.VIII,182; IX,76.
D. pauciflora; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,12	M.fr.XI,9.
D. diosmoides; Bentham in Journ. Linn. Soc. IX, 180 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,13	M.fr.IX,176.
D. fascicularis; Rudge in Transact. Linn. Soc. XI, 299, t. 22 (1813)	—	—	—	—	N.S.W.	—	—	B.fl.III,13	M.fr.IX,176.
D. pinifolia; Bentham in Journ. Linn. Soc. IX, 181 (1865)...	W.A.	—	—	—	—	—	—	B.fl.III,14	
D. rhadinophylla; F. v. M., fragm. IX, 175 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,175.
D. sanguinea; Bentham in Journ. Linn. Soc. IX, 181 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,14	M.fr.IX,176.
D. micropetala; Bentham in Journ. Linn. Soc. IX, 181 (1865)	—	S.A.	—	V.	—	—	—	B.fl.III,14	M.fr.I,12; IV,58; IX,176.
D. Schuermanni; Bentham in Journ. Linn. Soc. IX, 181 (1865)	—	S.A.	—	—	—	—	—	B.fl.III,14	M.fr.I,12; IV,58.
D. verticordina; Bentham in Journ. Linn. Soc. IX, 181 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,15	M.fr.IV,57.
D. Thomasii; Bentham in Journ. Linn. Soc. IX, 181 (1865)	—	—	—	—	—	Q.	—	B.fl.III,15	M.fr.IV,137.
D. virgata; F. v. M., fragm. IX, 176 (1875) from Cunningham (1835)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,16	M.fr.IX,176.
D. ciliata; F. v. M., Cens. 51 (1882) from Desf. (1821)	W.A.	—	—	—	—	—	—	B.fl.III,36	M.fr.VIII,182.
D. gracilis; F. v. M., Cens. 51 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,36	M.fr.IV,62; VIII,182.
D. Forrestii; F. v. M., fragm. XI, 9 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,9.
D. heterandra; F. v. M., Cens. 51 (1882) from Benth. (1866)	W.A.	—	—	—	—	—	—	B.fl.III,36	
D. Drummondii; F. v. M., Cens. 51 (1882) from Meissn. (1857)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. Endlicheri; F. v. M., Cens. 51 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. brevifolia; F. v. M., Cens. 51 (1882) from Benth. (1866)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. uncinata; F. v. M., Cens. 51 (1882) from Schauer (1844)	W.A.	—	—	—	—	—	—	B.fl.III,37	
D. megalopetala; F. v. M., Cens. 51 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,38	M.fr.VIII,182.
D. Turczaninowii; F. v. M., Cens. 51 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,38	M.fr.IX,176; XI,9.
D. axillaris; F. v. M., Cens. 51 (1882)...	W.A.	—	—	—	—	—	—	B.fl.III,38	
VERTICORDIA; De Candolle in Dict. class XI, 400 (1826). (Chrysorrhoea, perhaps also Huegelia.)									
V. Wilhelmii; F. v. M. in Transact. Vict. Inst. 122 (1855)	—	S.A.	—	—	—	—	—	B.fl.III,19	M.fr.X,28.
V. densiflora; Lindley in Bot. Regist. XXV, App. VI (1839)	W.A.	—	—	—	—	—	—	B.fl.III,20	
V. stelluligera; Meissner in Journ. Linn. Soc. I, 38 (1857)...	W.A.	—	—	—	—	—	—	B.fl.III,20	
V. minutiflora; F. v. M., fragm. IV, 58 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,20	M.fr.IV,38.
V. Fontainesii; De Candolle, prodr. III, 209 (1828)...	W.A.	—	—	—	—	—	—	B.fl.III,21	M.fr.VIII,182; X,28.
V. helichrysantha; F. v. M. in Benth. Fl. Austr. III, 21 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,21	
V. Brownii; De Candolle, prodr. III, 209 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,21	
V. fastigiata; Turczaninow in Bullet. de l'Ac. de St. Petersb. 402 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,22	M.fr.VIII,122; X,28.
V. Harveyi; Bentham, Fl. Austr. III, 22 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,22	
V. fimbrilepis; Turczaninow in Bull. Mosc. XX, 158 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,22	M.fr.X,28.
V. serrata; Schauer in nov. act. acad. Caes. XIX, suppl. II, 70 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,23	
V. nitens; Schauer in nov. act. acad. Caes. XIX, suppl. II, 71, t. 4 (1841) ...	W.A.	—	—	—	—	—	—	B.fl.III,23	M.fr.X,28.
V. grandiflora; Endlicher in Ann. Wien. Mus. II, 195 (1838)	W.A.	—	—	—	—	—	—	B.fl.III,23	
V. chrysantha; Endlicher in Ann. Wien. Mus. II, 195 (1838)	W.A.	—	—	—	—	—	—	B.fl.III,24	M.fr.X,28.
V. Preissii; Schauer in Lehm. pl. Preiss. I, 101 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,24	M.fr.VIII,182.
V. acerosa; Lindley in Bot. Regist. XXV, App. VI (1839)...	W.A.	—	—	—	—	—	—	B.fl.III,25	
V. polytricha; Bentham, Fl. Austr. III, 25 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,25	

V. oxylepis; Turczaninow in Bullet. de l'Ac. de St. Petersb. 402 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,25	M.fr.VIII,182.
V. humilis; Bentham, Fl. Austr. III, 26 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,26	
V. penicillaris; F. v. M., fragm. I, 226 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.III,26	M.fr.I,226.
V. multiflora; Turczaninow in Bull. Mosc. XX, 159 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,26	M.fr.X,28.
V. Huegelii; Endlicher in Hueg. enum. 16 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.III,27	
V. insignis; Endlicher in Hueg. enum. 47 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.III,27	M.fr.X,28.
V. habrantha; Schauer in Lehm. pl. Preiss. I, 100 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,28	M.fr.I,164.
V. monadelpha; Turczaninow in Bull. Mosc. XX, 158 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,28	M.fr.I,227; X,28.
V. Lehmanni; Schauer in Lehm. pl. Preiss. I, 99 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,29	
V. Cunninghamii; Schauer in nov. act. acad. Caes. XIX, suppl. II, 55 (1841) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,29	M.fr.X,28.
V. Jamiesonii; F. v. M. in Wieg's S. Sc. Rec. III, 68 (1883) ...	W.A.	—	—	—	—	—	—	—	
V. picta; Endlicher in Ann. des Wien. Mus. II, 194 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.III,30	M.fr.X,28.
V. pennigera; Endlicher in Hueg. enum. 46 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.III,30	M.fr.X,28.
V. Drummondii; Schauer in nov. act. acad. Caes. XIX, suppl. II, 56 (1841) ...	W.A.	—	—	—	—	—	—	B.fl.III,31	
V. Hughani; F. v. M., fragm. XI, 10 (1878) ...	W.A.	—	—	—	—	—	—	—	M.fr.XI,10.
V. pholidophylla; F. v. M., fragm. I, 227 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.III,31	M.fr.I,227.
V. spicata; F. v. M., fragm. I, 226 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,226.
V. lepidophylla; F. v. M., fragm. I, 228 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,228.
V. Forrestii; F. v. M. in Wing's S. Sc. Rec. III, 67 (1883) ...	W.A.	—	—	—	—	—	—	—	
V. ovalifolia; Meissner in Journ. Linn. Soc. I, 40 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,32	M.fr.I,227; X,28.
V. chrysostachya; Meissner in Journ. Linn. Soc. I, 41 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.I,227; X,28.
V. oculata; Meissner in Journ. Linn. Soc. I, 41 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.I,227.
V. grandis; Drummond in Hook. Kew Misc. V, 119 (1853) ...	W.A.	—	—	—	—	—	—	B.fl.III,33	M.fr.V,14; X,28; XI,10.

PILEANTHUS; Labillardière, Nov. Holl. pl. specim. II, 11 (1806).

P. peduncularis; Endlicher in Ann. des Wien. Mus. II, 196 (1838) ...	W.A.	—	—	—	—	—	—	B.fl.III,34	M.fr.I,225.
P. Limacis; Labillardière, Nov. Holl. pl. specim. II, 11, t. 149 (1806) ...	W.A.	—	—	—	—	—	—	B.fl.III,34	
P. filifolius; Meissner in Journ. Linn. Soc. I, 45 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,35	

CALYCOTHRIX; Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806). (Calytrix, Calythrix.)

C. aurea; Lindley, Bot. Regist. XXV, App. 5, t. 3 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,41	
C. puberula; Meissner in Journ. Linn. Soc. I, 48 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. flavescens; Cunningham in Bot. Mag. 3323 (1834) ...	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. asperula; Schauer in Lehm. pl. Preiss. I, 106 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,42	
C. sapphirina; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,43	M.fr.I,224.
C. breviseta; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,43	M.fr.VIII,182.
C. simplex; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,43	
C. empetroides; Schauer in nov. act. acad. Caes. XIX, suppl. II, 102 (1841) ...	W.A.	—	—	—	—	—	—	B.fl.III,44	
C. variabilis; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,44	
C. muricata; F. v. M., fragm. I, 224 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.III,44	M.fr.I,224.
C. gracilis; Bentham, Fl. Austr. III, 45 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. brevifolia; Meissner in Journ. Linn. Soc. I, 46 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. brachyphylla; Turczaninow in Bull. Mosc. XX, 161 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,45	
C. Leschenaultii; Schauer in Lehm. pl. Preiss. I, 104 (1844) ...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. Oldfieldii; Bentham, Fl. Austr. III, 46 (1866) ...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. glutinosa; Lindley, Bot. Regist. XXV, App. 5 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III,46	
C. angulata; Lindley, Bot. Regist. XXV, App. 6 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.III.47	
C. depressa; Turczaninow in Bull. Soc. Mosc. XX, 162 (1847) ...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. tenuifolia; Meissner in Journ. Linn. Soc. I, 46 (1857) ...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. strigosa; Cunningham in Bot. Mag. 3323 (1834) ...	W.A.	—	—	—	—	—	—	B.fl.III,47	
C. plumulosa; F. v. M., fragm. X, 27 (1876) ...	W.A.	—	—	—	—	—	—	—	M.fr.X,27.

C. decandra; R. Brown in De Candolle, prodr. III, 208 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,48	M.fr.I,146.
C. Birdii; F. v. M., fragm. X, 26 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,26.
C. tenuiramea; Turczaninow in Bull. Mosc. XXII, 20 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,48	
C. Fraseri; Cunningham in Bot. Mag. 3323 (1834)	W.A.	—	—	—	—	—	—	B.fl.III,48	M.fr.I,13.
C. granulosa; Bentham, Fl. Austr. III, 49 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,49	
C. Creswelli; F. v. M., fragm. X, 27 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,27.
C. microphylla; Cunningham in Bot. Mag. 3323 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.III,49	
C. longiflora; F. v. M., fragm. I, 12 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,49	M.fr.I,12.
C. leptophylla; Bentham, Fl. Austr. III, 50 (1866)	—	—	—	—	—	Q.	—	B.fl.III,50	
C. megaphylla; F. v. M., fragm. I, 146 (1859)	—	—	—	—	—	—	N.A.	B.fl.III,50	M.fr.I,13,146.
C. tetragona; Labillardière, Nov. Holl. pl. specim. II, 8, t. 146 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,50	M.fr.IV,36; VIII,183.
C. Sullivani; F. v. M., fragm. IX, 1 (1875)	—	—	—	V.	—	—	—	—	M.fr.IX,1.
C. conferta; Cunningham in Bot. Mag. 3323 (1834)	—	—	—	—	—	—	N.A.	B.fl.III,51	
C. arborescens; F. v. M. in Transact. phil. Inst. Vict. III, 42 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,51	
C. brachychaeta; F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,52	
C. achaeta; F. v. M. in Transact. phil. Inst. Vict. III, 43 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,52	
C. laricina; R. Brown in Benth. Fl. Austr. III, 52 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,52	
LHOTZKYA; Schauer in Linnaea X, 309 (1835).									
L. glaberrima; F. v. M., fragm. I, 13 (1858)	—	S.A.	—	—	—	—	—	B.fl.III,53	M.fr.I,13.
L. genetylloides; F. v. M. in Transact. phil. Soc. Vict. I, 16 (1854)	—	S.A.	—	V.	—	—	—	B.fl.III,54	
L. Smeatoniana; F. v. M. in Austral. Journ. of Pharm. II, 5 (1887)	—	S.A.	—	—	—	—	—	—	
L. violacea; Lindley, Bot. Regist. XXV, App. VII (1839)...	W.A.	—	—	—	—	—	—	B.fl III,54	M.fr.XI,8.
L. ciliata; F. v. M. in Benth. Fl. Austr. III, 54 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,54	
L. brevifolia; Schauer in Lehm. pl. Preiss. I, 103 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,54	
L. purpurea; F. v. M., fragm. I, 224 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,55	M.fr.I,224.
L. ericoides; Schauer in Lindl. Introd. sec. ed. 439 (1835)...	W.A.	—	—	—	—	—	—	B.fl.III,55	M.fr.XI,9.
L. acutifolia; Lindley, Bot. Regist. XXV, App. VII (1839)	W.A.	—	—	—	—	—	—	B.fl.III,55	M.fr.XI,8.
L. Harvestiana; F. v. M., fragm. XI, 8 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,8.
THRYPTOMENE; Endlicher in den Ann. des Wien. Mus. II, 192 (1838). (Homalocalyx, Paryphantha, Astraea, Eremopyxis, Micromyrtus.)									
T. mucronulata; Turczaninow in Bull. Mosc. XX, 156 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,58	
T. Australis; Endlicher in den Ann. des Wien. Mus. II, 192 (1838)	W.A.	—	—	—	—	—	—	B.fl.III,58	
T. tenella; Bentham, Fl. Austr. III, 59 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,59	M.fr.VIII,182; X,24.
T. prolifera; Turczaninow in Bull. Mosc. XXXV, 324 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,59	
T. saxicola; Schauer in Lehm. pl. Preiss. I, 102 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,59	M.fr.IV,75.
T. Johnsonii; F. v. M., fragm. IV, 77 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,60	M.fr.IV,77.
T. racemulosa; Turczaninow in Bull. Mosc. XX, 156 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,60	
T. denticulata; Bentham in Fl. Austr. III, 60 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,60	M.fr.IV,75.
T. baeckeacea; F. v. M., fragm. IV, 65 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,60	M.fr.IV,65.
T. strongylophylla; F. v. M. in Benth. Fl. Austr. III, 61 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,61	
T. hyporhytis; Turczaninow in Bull. Mosc. XXXV, 324 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,61	
T. Maisonneuvii; F. v. M., fragm. IV, 64 (1864)	—	S.A.	—	—	—	—	N.A.	B.fl.III,61	M.fr.IV,64.
T. urceolaris; F. v. M., fragm. X, 25 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,25.
T. Mitchelliana; F. v. M., fragm. I, 11 (1858)	—	S.A.	—	V.	—	—	—	B.fl.III,61	M.fr.XI,62.
T. Miqueliana; F. v. M., fragm. I, 11 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,62	M.fr.I,11.
T. micrantha; J. Hooker in Hook. Kew. Misc. V, 299, t. 8 (1853)	—	—	T.	—	—	—	—	B.fl.III,62	
T. stenocalyx; F. v. M., fragm. X, 23 (1876)...	W.A.	—	—	—	—	—	—	—	M.fr.X,23.
T. ericaea; F. v. M., fragm. I, 12 (1858)	—	S.A.	—	V.	—	—	—	B.fl.III,62	M.fr.I,12.
T. oligandra; F. v. M., fragm. I, 11 (1858)	W.A.	—	—	—	—	Q.	N.A.	B.fl.III,63	M.fr.IV,169.
T. elobata; F. v. M., fragm. IV, 63 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,64	M.fr.IV,63.

T. racemosa ; F. v. M. in Benth. Fl. Austr. III, 64 (1866)	W.A.	—	—	—		—	—	—	B.fl.III,64	
T. trachycalyx ; F. v. M., fragm. X, 25 (1876)	W.A.	—	—	—		—	—	—	--	M.fr.X,25.
T. imbricata ; F. v. M., Cens. 53 (1882) from Benth. (1866)	W.A.	—	—	—		—	—	—	B.fl.III,64	
T. flaviflora ; F. v. M., fragm. VIII, 13 (1873)	W.A.	S.A.	—	—		—	—	—	—	M.fr.X,25.
T. Drummondii ; F. v. M., Cens. 53 (1882) from Benth. (1866)	W.A.	—	—	—		—	—	—	B.fl.III,64	M.fr.VIII,183.
T. Elliottii ; F. v. M., fragm. IX, 62 (1875)	—	S.A.	—	—		—	—	—	—	M.fr.IX,62.
T. auriculata ; F. v. M., fragm. X, 24 (1876)	—	S.A.	—	—		—	—	—	—	M.fr.X,24.
T. hymenonema ; F. v. M., fragm. X, 26 (1876)	W.A.	—	—	—		—	—	—	—	M.fr.X,26.
T. ciliata ; F. v. M. in Woolls, pl. of the neighb. of Sydney 23 (1880)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,65	M.fr.I,30&242; IV,63.
T. minutiflora ; F. v. M. in Woolls, pl. of the neighb. of Sydney 23 (1880) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,65	
T. leptocalyx ; F. v. M., Cens. 53 (1882) from Benth. (1866)	—	—	—	—	—	Q.	—	B.fl.III,65	M.fr.I,30.
T. homalocalyx ; F. v. M., fragm. IV, 63 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.III,56	M.fr.IV,63.
T. polyandra ; F. v. M., fragm. IV, 77 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.III,56	M.fr.IV,77.
WEHLIA ; F. v. M., fragm. X, 22 (1876).										
W. thryptomenoides ; F. v. M., fragm. X, 22 (1876)	W.A.	—	—	—		—	—	—	—	M.fr.X,22.
W. coarctata ; F. v. M., fragm. X, 23 (1876)...	W.A.	—	—	—		—	—	—	—	M.fr.X,23.
W. staminosa ; F. v. M. in Wing's S. Sc. Rec. III, 282 (1884)	W.A.	—	—	—		—	—	—	—	
BAECKEA ; Linné, spec. plant. 358 (1753). (Jungia, Imbricaria, Scholtzia, Schidiomyrtus, Rinzia, Euryomyrtus, Camphoromyrtus, Tetrapora, Harmogia, Oxymyrrhine, Babingtonia, Ericomyrtus, Piptandra, Anticoryne.)										
B. platystemonea ; Bentham, Fl. Austr. III, 74 (1866)	W.A.	—	—	—		—	—	—	B.fl.III,74	M.fr.VIII,183.
B. Fumana ; F. v. M., fragm. IV, 68 (1864)	W.A.	—	—	—		—	—	—	B.fl.III,74	M.fr.IV,68.
B. dimorphandra ; F. v. M. in Benth. Fl. Austr. III, 74 (1866)	W.A.	—	—	—		—	—	—	B.fl.III,74	
B. schollerifolia ; Lehmann, pl. Preiss. II, 369 (1847)	W.A.	—	—	—		—	—	—	B.fl.III,75	M.fr.IV,68; VIII,183.
B. oxycoccoides ; Bentham, Fl. Austr. III, 75 (1866)	W.A.	—	—	—		—	—	—	B.fl.III,75	M.fr.VIII,183.
B. Drummondii ; Bentham, Fl. Austr. III, 75 (1866)	W.A.	—	—	—		—	—	—	B.fl.III,75	M.fr.VIII,183.
B. diffusa ; Sieber in De Candolle, prodr. III, 230 (1828)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,76	M.fr.IV,67.
B. crassifolia ; Lindley in Mitch. Three Exped. II, 115 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,76	M.fr.IV,66.
B. cryptandroides ; F. v. M., fragm. X, 29 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,29.
B. tetragona ; F. v. M. in Benth. Fl. Austr. III, 77 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,77	
B. ericaea ; F. v. M., fragm. I, 31 (1858)	—	S.A.	—	V.	N.S.W.	—	--	B.fl.III,77	
B. polystemonea ; F. v. M., fragm. II, 124 (1861)	—	S.A.	—	—	—	—	—	B.fl.III,77	M.fr.III,24.
B. crenulata ; R. Brown in Flind. voy. 548 (1814)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,78	M.fr.IV,65.
B. brevifolia ; De Candolle, prodr. III, 230 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,78	
B. Gunniana ; Shauer in Walp. Rep. II, 920 (1843)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,78	M.fr.IV,66.
B. diosmifolia ; Rudge in Transact. Linn. Soc. VIII, 298, t. 13 (1807)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,79	M.fr.I,29.
B. leptocaulis ; J. Hooker, icon. pl. t. 298 (1840)	—	—	T.	—	—	—	—	B.fl.III,79	
B. arbuscula ; R. Brown in Benth. Fl. Austr. III, 79 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,79	
B. astarteoides ; Bentham, Fl. Austr. III, 80 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,80	
B. linifolia ; Rudge in Transact. Linn. Soc. VIII, 297, t. 12 (1807)	—	—	—	V.	N.S.W.	—	—	B.fl.III,80	M.fr.IV,71.
B. stenophylla ; F. v. M., fragm. I, 13 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,80	M.fr.I,13.
B. camphorata ; R. Brown in Bot. Mag. t. 2694 (1826)	—	—	—	V.	N.S.W.	—	—	B.fl.III,81	M.fr.IV,70; IV,177.
B. virgata ; Andrews, Bot. Reposit. t. 598 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.III,81	M.fr.I,69; II,31 ... P.
B. crenatifolia ; F. v. M., fragm. IV, 70 (1864)	—	—	—	V.	—	—	—	B.fl.III,82	M.fr.IV,70.
B. Cunninghamii ; Bentham, Fl. Austr. III, 82 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,82	
B. densifolia ; Smith in Transact. Linn. Soc. III, 260 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,82	M.fr.IV,71.
B. Behrii ; F. v. M., fragm. IV, 68 (1864)	W.A.	S.A.	—	V.	N.S.W.	—	--	B.fl.III,83	M.fr.IV,68.
B. uncinella ; Bentham, Fl. Austr. III, 84 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,84	
B. polyandra ; F. v. M., fragm. IV, 12 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,84	M.fr.II,30; IV,72.
B. corynophylla ; F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,84	M.fr.IV,72.

B. pachyphylla ; Bentham, Fl. Austr. III, 85 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,85	
B. crispiflora ; F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,85	M.fr.II,31.
B. Blackettii ; F. v. M., fragm. VIII, 181 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,181.
B. camphoresmae ; Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,86	
B. pulchella ; De Candolle, prodr. III, 230 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,86	M.fr.IV,67.
B. ochropetala ; F. v. M., fragm. X, 29 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,29.
B. pygmaea ; R. Brown in Benth. Fl. Austr. III, 86 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,86	
B. corymbulosa ; Bentham, Fl. Austr. III, 87 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,87	M.fr.VIII,183.
B. floribunda ; Bentham, Fl. Austr. III, 87 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,87	
B. pentandra ; F. v. M., fragm. II, 31 ; IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,87	M.fr.II,31; IV,72; VIII,183.
B. pentagonantha ; F. v. M., fragm. IV, 73 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,87	M.fr.IV,73.
B. robusta ; F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,88	M.fr.IV,72.
B. ovalifolia ; F. v. M., fragm. IV, 72 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,88	M.fr.II,32; IV,72; VIII,183.
B. subcuneata ; F. v. M., fragm. IV, 73 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,88	M.fr.IV,73.
B. grandiflora ; Bentham, Fl. Austr. III, 89 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,89	
B. uberiflora ; F. v. M., Cens. 53 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,67	M.fr.IV,74.
B. involucrata ; Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,68	M.fr.IV,74.
B. spathulata ; F. v. M., Cens. 54 (1882) from Benth. (1866)	W.A.	—	—	—	—	—	—	B.fl.III,68	
B. ciliata ; F. v. M., Cens. 54 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,68	M.fr.IV,76.
B. capitata ; F. v. M., Cens. 54 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,69	
B. umbellifera ; F. v. M., Cens. 54 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,69	M.fr.IV,75.
B. laxiflora ; F. v. M., Cens. 54 (1882) from Benth. (1866)	W.A.	—	—	—	—	—	—	B.fl.III,69	
B. leptantha ; F. v. M., Cens. 54 (1882) from Benth. (1866)	W.A.	—	—	—	—	—	—	B.fl.III,69	
B. parviflora ; F. v. M., Cens. 54 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,70	M.fr.IV,76.
B. oligandra ; F. v. M., Cens. 54 (1882)	W.A.	—	—	—	—	—	—	B.fl.III,70	
B. serpillifolia ; F. v. M., fragm. X, 30 (1876)	W.A.	—	—	—	—	—	—	B.fl.III,70	M.fr.VIII,183.
B. teretifolia ; F. v. M., Cens. 54 (1882) from Benth. (1866)	W.A.	—	—	—	—	—	—	B.fl.III,70	
ASTARTEA ; De Candolle in Dict. class. XI, 400 (1826).									
A. ambigua ; F. v. M., fragm. II, 32 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,89	M.fr.II.32.
A. fascicularis ; De Candolle, prodr. III, 210 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,90	M.fr.VIII,183.
A. intratropica ; F. v. M., fragm. I, 83 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,90	M.fr.I,83.
HYPOCALYMMA ; Endlicher in Hueg. enum. pl. Nov. Holl. Austr. occ. 51 (1837), implied.									
H. xanthopetalum ; F. v. M., fragm. II, 29 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,92	M.fr.II,29.
H. robustum ; Endlicher in Hueg. enum. 50 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,92	
H. longifolium ; F. v. M., fragm. II, 28 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,92	M.fr.II,28.
H. strictum ; Schauer in Lehm. pl. Preiss. I, 111 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,93	
H. tetrapterum ; Turczaninow in Bull. Mosc. XXXV, 325 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,93	
H. linifolium ; Turczaninow in Bull. Mosc. XXXV, 325 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,93	
H. angustifolium ; Endlicher in Hueg. enum. 50 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,94	M.fr.II,28.
H. ericifolium ; Bentham, Fl. Austr. III, 94 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,94	
H. cordifolium ; Lehmann, plant. Preiss. I, 112 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,94	
H. speciosum ; Turczaninow in Bull. de l'Acad. de St. Pet. 409 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,95	M.fr.VIII,183.
H. Phillipsii ; Harvey in nat. hist. Rev. V, 296, t. 22 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,95	
H. myrtifolium ; Turczaninow in Bull. de St. Pet. 410 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,95	M.fr.VIII,183.
BALAUSTION ; Hooker, icones plantarum 852 (1852). (Punicella, Cheynia.)									
B. pulcherrimum ; Hooker, icon. plant. 852 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,96	M.fr.VIII,183.
AGONIS ; De Candolle, prodr. III, 226 (1828), implied. (Billietia, Billetia, Billottia.)									
A. spathulata ; Schauer in Lehm. pl. Preiss. I, 117 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,97	M.fr.XI,120.
A. floribunda ; Turczaninow in Bull. Mosc. II, 20 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,97	

A. marginata ; De Candolle, prodr. III, 226 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.XI,120.				
A. lysicephala ; F. v. M. & Bailey in synops. Queensl. Flora sec. suppl. 24 (1888)	—	—	—	—	—	Q.	—	—					
A. linearifolia ; De Candolle, prodr. III, 227 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.VIII,183; XI,120.				
A. juniperina ; Schauer in Lehm. pl. Preiss. I, 118 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,98	M.fr.XI,120.				
A. parviceps ; Schauer in Lehm. pl. Preiss. I, 119 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,99	M.fr.XI,120.				
A. obtusissima ; F. v. M., fragm. XI, 119 (1881)	W.A.	—	—	—	—	—	—	—	M.fr.XI,119.				
A. flexuosa ; De Candolle, prodr. III, 226 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,99	M.fr.XI,120.				
A. undulata ; Bentham, Fl. Austr. III, 100 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,100					
A. hypericifolia ; Schauer in Lehm. pl. Preiss. I, 117 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,100	M.fr.XI,120.				
A. Scortechiniana ; F. v. M., fragm. XI, 118 (1881)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,118.				
A. grandiflora ; Bentham, Fl. Austr. III, 100 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,100	M.fr.XI,120.				
LEPTOSPERMUM ; R. et G. Forster, char. gen. 71, t. 36 (1776). (Fabricia, Homalospermum, Pericalymma.)													
L. Fabricia ; Bentham, Fl. Austr. III, 102 (1866)	—	—	—	—	—	Q.	—	B.fl.III,102					
L. laevigatum ; F. v. M., Annual Report 22 (1858)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,103	M.fr.IV,60.				
L. firmum ; Bentham, Fl. Austr. III, 104 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,104					
L. flavescens ; Smith in Transact. Linn. Soc. III, 262 (1797)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,104	M.fr.IV,61	...	As.		
L. scoparium ; R. & G. Forster, char. gen. 48 (1776)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,105	M.fr.IV,61	...	N.Z.		
L. arachnoideum ; Smith in Transact. Linn. Soc. III, 263 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.III,105					
L. spinescens ; Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,106					
L. lanigerum ; Smith in Transact. Linn. Soc. III, 263 (1797)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,106					
L. parvifolium ; Smith in Transact. Linn. Soc. III, 263 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.III,107					
L. stellatum ; Cavanilles, icon. pl. IV, 16, t. 330	—	—	—	—	N.S.W.	Q.	—	B.fl.III,107					
L. attenuatum ; Smith in Transact. Linn. Soc. III, 262 (1797)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,108	M.fr.II,28; IV,62.				
L. myrtifolium ; Sieber in De Candolle, prodr. III, 238 (1828)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,108					
L. rupestre ; J. Hooker in Hook. icon. pl. t. 308 (1841)	—	—	T.	—	—	—	—	B.fl.III,108					
L. myrsinoides ; Schlechtendal in Linnaea XX, 653 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,109	M.fr.VI,61.				
L. erubescens ; Schauer in Lehm. pl. Preiss. I, 121 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,109	M.fr.II,28; IV,61; VIII,183.				
L. abnorme ; F. v. M. in Benth. Fl. Austr. III, 109 (1866)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,109	M.fr.II,27.				
L. Roei ; Bentham, Fl. Austr. III, 110 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,110					
L. floridum ; Bentham, Fl. Austr. III, 110 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,110					
L. ellipticum ; Endlicher in Hueg. enum. 51 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,110					
L. crassipes ; Lehmann, ind. sem. hort. Hamb. (1842)	W.A.	—	—	—	—	—	—	B.fl.III,110					
KUNZEA ; Reichenbach, conspect. regn. veget. 175 (1828). (Salisia, Pentagonaster.)													
K. micrantha ; Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,112	M.fr.VIII,183.				
K. eriocalyx ; F. v. M., fragm. II, 28 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,112	M.fr.II,28.				
K. Muelleri ; Bentham, Fl. Austr. III, 113 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,113					
K. ericifolia ; Reichenbach, consp. 175 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,113					
K. Preissiana ; Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,113					
K. recurva ; Schauer in Lehm. pl. Preiss. I, 125 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,114	M.fr.VIII,183.				
K. micromera ; Schauer in Lehm. pl. Preiss. III, 223 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,114					
K. pauciflora ; Schauer in Lehm. pl. Preiss. I, 124 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,114	M.fr.II,28; VIII,183.				
K. calida ; F. v. M., fragm. VI, 23 (1867)	—	—	—	—	—	Q.	—	—	M.fr.VI,23.				
K. parvifolia ; Schauer in Lehm. pl. Preiss. I, 124 (1844)	—	—	—	—	N.S.W.	—	—	B.fl.III,115	M.fr.II,28; VIII,13.				
K. peduncularis ; F. v. M. in Transact. Vict. Inst. 144 (1855)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,115	M.fr.II,28; X,55.				
K. corifolia ; Reichenbach, consp. 175 (1828)...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,115					
K. capitata ; Reichenbach, consp. 175 (1828)...	—	—	—	—	N.S.W.	—	—	B.fl.III,116	M.fr.II,28.				
K. pomifera ; F. v. M. in Transact. Vict. Inst. 124 (1855)	—	S.A.	—	V.	—	—	—	B.fl.III,116	M.fr.IV,62.				
K. sericea ; Turczaninow in Bull. Mosc. XX, 162 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,117					
K. Baxteri ; Schauer in Lehm. pl. Preiss. I, 123 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,117	M.fr.III,153.				
K. opposita ; F. v. M., fragm. VI, 24 (1867)	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,24.				

CALLISTEMON; R. Brown in Flind. voy. II, 547 (1814).

C. speciosus; De Candolle, prodr. III, 224 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,119	M.fr.I,14.
C. phoeniceus; Lindley, Bot. Regist. XXV, App. 10 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,119	M.fr.IV,53.
C. lanceolatus; De Candolle, prodr. III, 224 (1828)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,120	M.fr.I,14; IV,53.
C. coccineus; F. v. M., fragm. I, 13 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,120	M.fr.IV,54.
C. salignus; De Candolle, prodr. III, 223 (1828)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,120	M.fr.I,14; IV,54.
C. pityoides; Miquel in Nederl. Kruidk. Arch. IV, 140 (1859)	—	—	—	V.	N.S.W.	Q.	—	—	
C. rigidus; R. Brown in Bot. Regist. t. 393 (1819)	—	—	—	—	N.S.W.	—	—	B.fl.III,121	M.fr.IV,54.
C. linearis; De Candolle, prodr. III, 223 (1828)	—	—	—	V.	N.S.W.	—	—	B.fl.III,122	M.fr.IV,54.
C. pinifolius; De Candolle, prodr. III, 223 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,122	M.fr.IV,53.
C. teretifolius; F. v. M. in Schlecht. Linnaea XXV, 387 (1852)	—	S.A.	—	—	—	—	—	B.fl.III,122	M.fr.IV,53.
C. brachyandrus; Lindley in Journ. Hort. Soc. IV, 112 (1849)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,122	M.fr.IV,52.

MELALEUCA; Linné, mantiss. plant. 14 (1767). (Gymnagathis, Asteromyrtus.)

M. macronychia; Turczaninow in Bullet. de St. Petersb. 420 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,129	M.fr.VIII,183.	
M. lateritia; Otto in allgem. Gart. Zeit. II, 257 (1834)	W.A.	—	—	—	—	—	—	B.fl.III,130		
M. calothamnoides; F. v. M., fragm. III, 114 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,130	M.fr.III,114.	
M. blaeriaefolia; Turczaninow in Bull. Mosc. XX, 165 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,130		
M. diosmifolia; Andrews, Bot. Reposit. t. 476 (1808)	W.A.	—	—	—	—	—	—	B.fl.III,130		
M. elliptica; Labillardière, Nov. Holl. pl. spec. II, 31, t. 173 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,131	M.fr.VIII,183.	
M. hypericifolia; Smith in Transact. Linn. Soc. III, 279 (1797)	—	—	—	V.	N.S.W.	—	—	B.fl.III,131		
M. fulgens; R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,131		
M. acuminata; F. v. M., fragm. I, 15 (1858)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,132	M.fr.I,15.	
M. leptoclada; Bentham, Fl. Austr. III, 132 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,132		
M. basicephala; Bentham, Fl. Austr. III, 133 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,133		
M. gibbosa; Labillardière, Nov. Holl. pl. spec. II, 30, t. 172 (1806)	—	S.A.	T.	V.	—	—	—	B.fl.III,133		
M. decussata; R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	—	S.A.	—	V.	—	—	—	B.fl.III,133		
M. Wilsonii; F. v. M., fragm. II, 124, t. 15 (1861)	—	S.A.	—	V.	—	—	—	B.fl.III,134	M.fr.II,124.	
M. thymifolia; Smith in Transact. Linn. Soc. III, 278 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,134	M.fr.X,55.	
M. violacea; Lindley, Bot. Regist. XXV, App. 8 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,135		
M. cardiophylla; F. v. M., fragm. I, 225 (1859)	W.A.	—	—	—	—	—	—	B.fl.III,135	M.fr.III,164.	
M. undulata; Bentham, Fl. Austr. III, 135 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,135		
M. depauperata; Turczaninow in Bullet. de St. Petersb. 424 (1852)	W.A.	—		—	—	—	—	B.fl.III,136	M.fr.III,120; VIII,183.	
M. lateriflora; Bentham, Fl. Austr. III, 136 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,136		
M. exarata; F. v. M., fragm. III, 114 (1862)...	W.A.	—	—	—	—	—	—	B.fl.III,136	M.fr.VIII,184.	
M. teretifolia; Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,137	M.fr.III,117.	
M. alsophila; Cunningham in Benth. Fl. Austr. III, 137 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,137		
M. acacioides; F. v. M., fragm. III. 116 (1862)	—	—	—	—	—	—	N.A.	B.fl.III,138	M.fr.III,116.	
M. Baxteri; Bentham, Fl. Austr. III, 138 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,138		
M. symphyocarpa; F. v. M. in Transact. phil. Inst. Vict. III, 44 (1858) ...	—	—	—	—	—	Q.	N.A.	B.fl.III,138		
M. angustifolia; Gaertner, de fruct. I, 172, t. 35 (1788)	—	—	—	—	—	Q.	—	B.fl.III,139		
M. pauciflora; Turczaninow in Bull. Mosc. XX, 166 (1847)...	—	—	—	—	N.S.W.	—	—	B.fl.III,139		
M. squarrosa; Donn, Hort. Cantabr. ed. sec. 101 (1800)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,139	M.fr.X,55.	
M. adnata; Turczaninow in Bullet. de St. Petersb. 425 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,140	M.fr.III,117; VIII,184.	
M. linariifolia; Smith in Transact. Linn. Soc. III, 278 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,140	M.fr.X,55.	
M. trichostachya; Lindley in Mitch. Trop. Austr. 277 (1848)	—	S.A.	—	—	—	Q.	N.A.	—		
M. radula; Lindley, Bot. Regist. XXV, App. 8 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,141	M.fr.X,55.	
M. pulchella; R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,141	M.fr.III,120.	
M. conferta; Bentham, Fl. Austr. III, 142 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,142		
M. Leucadendron; Linné, mantiss. 105 (1767)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,142	M.fr.IV,55 ... As., P.	

M. lasiandra; F. v. M., fragm. III, 115 (1862)	—	—	—	—	—	—	—	—	—	—	N.A.	B.&H. III, 143	M.&F. III, 115.	
M. genistifolia; Smith in Transact. Linn. Soc. III, 277 (1797)	—	—	—	—	—	—	—	—	N.S.W.	Q.	N.A.	B.&H. III, 143	M.&F. I, 15; X, 53.	
M. stypheloides; Smith in Transact. Linn. Soc. III, 276 (1797)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.&H. III, 144	M.&F. I, 15.	
M. Hueglii; Endlicher in Hueg. enum. 48 (1837)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 144		
M. sericifolia; F. v. M. in Austr. Journ. of Pharm. I, 273 (1886)	—	—	—	—	W.A.	—	—	—	—	—	—	—		
M. diosmifolia; F. v. M., fragm. III, 133 (1863)	—	—	—	—	—	—	—	—	—	—	N.A.	B.&H. III, 144	M.&F. III, 133.	
M. linophylla; F. v. M., fragm. III, 115 (1862)	—	—	—	—	—	—	—	—	—	—	N.A.	B.&H. III, 145	M.&F. III, 115.	
M. parviflora; Lindley, Bot. Regist. XXV, App. 8 (1839)	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.&H. III, 145	M.&F. III, 115.	
M. Dennei; F. v. M. in proceed. L. Soc. of N.S.W. sec. ser. I, 1106 (1886)	—	—	—	—	—	—	—	—	N.S.W.	—	—	—		
M. laxiflora; Turczaninow in Bullet. de St. Petersb. 421 (1852)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 145	M.&F. VIII, 184; X, 53.	
M. armillaris; Smith in Transact. Linn. Soc. III, 277 (1797)	—	—	—	—	—	—	—	V.	N.S.W.	Q.	—	B.&H. III, 146		
M. cylindrica; R. Brown in Benth. Fl. Austr. III, 146 (1866)	—	—	—	—	—	S.A.	—	—	—	—	—	—		
M. hamulosa; Turczaninow in Bull. Mosc. XX, 165 (1847)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 146	M.&F. III, 116.	
M. subdolenta; Turczaninow in Bullet. de St. Petersb. 422 (1852)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 146	M.&F. III, 119; VIII, 184.	
M. glaberrima; F. v. M., fragm. III, 119 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 147	M.&F. III, 119.	
M. rhaphiophylla; Schauer in Lehm. pl. Preiss. I, 143 (1844)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 147		
M. cymbifolia; Bentham, Fl. Austr. III, 148 (1866)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 148		
M. cuticularis; Labillardière, Nov. Holl. pl. spec. II, 30, t. 171 (1806)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 148		
M. sparviflora; Turczaninow in Bull. Mosc. XX, 167 (1847)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 148		
M. calycina; R. Brown in Ait. hort. Kew. sec. ed. IV, 416 (1812)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 148	M.&F. VIII, 184.	
M. cordata; Turczaninow in Bullet. de St. Petersb. 418 (1852)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 149	M.&F. VIII, 184.	
M. globifera; R. Brown in Ait. hort. Kew. sec. ed. IV, 411 (1812)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 149		
M. megacephala; F. v. M., fragm. III, 117 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 149	M.&F. III, 117.	
M. neurophila; F. v. M., fragm. III, 113 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 150	M.&F. III, 113.	
M. Oldfieldii; F. v. M., fragm. III, 118 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 150		
M. uncinata; R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	—	—	—	—	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.&H. III, 150	M.&F. III, 118.	
M. concreta; F. v. M., fragm. III, 118 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 151	M.&F. III, 118.	
M. filifolia; F. v. M., fragm. III, 119 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 151	M.&F. III, 119.	
M. glomerata; F. v. M., Rep. Babb. Exped. 8 (1858)	—	—	—	—	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.&H. III, 151		
M. pentagona; Labillardière, Nov. Holl. pl. spec. II, 27, t. 166 (1806)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 152	M.&F. VIII, 184.	
M. cilicna; Turczaninow in Bull. Soc. Mosc. XXXV, 326 (1862)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 152		
M. polycephala; Bentham, Fl. Austr. III, 152 (1866)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 152	M.&F. VIII, 184.	
M. spathulata; Schauer in Lehm. pl. Preiss. I, 134 (1844)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 153	M.&F. VIII, 184.	
M. eriantha; Bentham, Fl. Austr. III, 153 (1866)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 153		
M. subtrigona; Schauer in Lehm. pl. Preiss. I, 139 (1844)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 153		
M. seriata; Lindley, Bot. Regist. XXV, App. 8 (1839)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 154		
M. scabra; R. Brown in Ait. hort. Kew. sec. ed. IV, 414 (1812)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 154	M.&F. VIII, 184.	
M. urceolaris; F. v. M. in Benth. Fl. Austr. III, 154 (1866)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 154		
M. trichophylla; Lindley, Bot. Regist. XXV, App. 8 (1839)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 155	M.&F. III, 114.	
M. holosericea; Schauer in Lehm. pl. Preiss. I, 139 (1844)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 155		
M. squamea; Labillardière, Nov. Holl. pl. spec. II, 28, t. 168 (1806)	—	—	—	—	—	S.A.	T.	V.	N.S.W.	—	—	B.&H. III, 155		
M. densa; R. Brown in Ait. hort. Kew. sec. ed. IV, 415 (1812)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 156	M.&F. VIII, 184.	
M. thymoides; Labillardière, Nov. Holl. pl. spec. II, 27, t. 167 (1806)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 156		
M. striata; Labillardière, Nov. Holl. pl. spec. II, 26, t. 165 (1806)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 157		
M. polygaloides; Schauer in Lehm. pl. Preiss. I, 142 (1844)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 157		
M. incana; R. Brown in Bot. Regist. t. 410 (1819)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 157		
M. nodosa; Smith in Transact. Linn. Soc. III, 276 (1797)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.&H. III, 158		
M. pungens; Schauer in Lehm. pl. Preiss. I, 138 (1844)	—	—	—	—	W.A.	—	—	—	—	—	—	B.&H. III, 158	M.&F. VIII, 184.	
M. ericifolia; Smith in Transact. Linn. Soc. III, 276 (1797)	—	—	—	—	—	S.A.	T.	V.	N.S.W.	Q.	—	B.&H. III, 159		
M. pustulata; J. Hooker in Lond. Journ. VI, 476 (1844)	—	—	—	—	—	S.A.	T.	V.	N.S.W.	—	—	B.&H. III, 160		

M. viminea; Lindley, Bot. Regist. XXV, App. 8 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,159	
M. microphylla; Smith in Rees, Cyclop. XXIII (1812)	W.A.	—	—	—	—	—	—	B.fl.III,159	M.fr.III,116.
M. tenella; Bentham, Fl. Austr. III, 160 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,160	
M. leiocarpa; F. v. M., fragm. X, 55 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,55.
M. leiopyxis; F. v. M. in Bentham, Fl. Austr. III, 160 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,160	
M. acerosa; Schauer in Lehm. pl. Preiss. I, 137 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,160	M.fr.III,118.
M. pauperiflora; F. v. M., fragm. III, 116 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,161	M.fr.III,116.
M. aspalathoides; Schauer in Lehm. pl. Preiss. I, 140 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,161	M.fr.VIII,184.
M. cucullata; Turczaninow in Bullet. de St. Petersb. 424 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,161	M.fr.VIII,184.
M. minutifolia; F. v. M. in Transact. phil. Soc. Vict. III, 45 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.III,162	
M. foliolosa; Cunningham in Benth. Fl. Austr. III, 162 (1866)	—	—	—	—	—	Q.	—	B.fl.III,162	
M. micromera; Schauer in Lehm. pl. Preiss. I, 146 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,162	
M. thuyoides; Turczaninow in Bull. Soc. Mosc. XX, 167 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,162	M.fr.III,114.
M. tamariscina; Hooker in Mitch. Trop. Austr. 262 (1848)	—	—	—	—	—	Q.	—	B.fl.III,163	M.fr.III,115.
M. quadrifaria; F. v. M. in Wing's S. Sc. Rec. April (1886)	W.A.	S.A.	—	—	—	—	—	—	
CONOTHAMNUS; Lindley, Bot. Regist. XXV, App. 11 (1839). (Trichobasis.)									
C. trinervis; Lindley, Bot. Regist. XXV, App. IX (1839)	W.A.	—	—	—	—	—	—	B.fl.III,163	
C. divaricatus; Bentham, Fl. Austr. III, 164 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,164	M.fr.VIII,184.
BEAUFORTIA; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812). (Schizopleura.)									
B. sparsa; R. Brown in Ait. hort. Kew. sec. ed. IV, 419 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,165	
B. decussata; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,166	
B. squarrosa; Schauer in nov. act. acad. Caes. XXI, 11 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,166	
B. orbifolia; F. v. M., fragm. III, 110 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,166	M.fr.III,110.
B. anisandra; Schauer in nov. act. acad. Caes. XXI, 13, t. I (1844)	W.A.	—	—	—	—	—	—	B.fl.III,167	
B. macrostemon; Lindley, Bot. Regist. XXV, App. 10 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,167	
B. heterophylla; Turczaninow in Bullet. de St. Petersb. 427 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,167	M.fr.VIII,184.
B. Schaueri; Preiss in nov. act. acad. Caes. XXI, 18 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,168	
B. purpurea; Lindley, Bot. Regist. XXV, App. X, t. 3 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,168	
B. Dampieri; Cunningham in Bot. Mag. t. 3272 (1833)	W.A.	—	—	—	—	—	—	B.fl.III,168	M.fr.III,111.
B. elegans; Schauer in nov. act. acad. Caes. XXI, 20 (1844)	W.A.	—	—	—	—	—	N.A.	B.fl.III,169	
B. interstans; F. v. M., fragm. X, 30 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,30.
B. micrantha; Schauer in nov. act. acad. Caes. XXI, 22 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,169	M.fr.VIII,184.
REGELIA; Schauer in Linnaea, XVII, 243 (1843).									
R. grandiflora; Bentham, Fl. Austr. III, 170 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,170	M.fr.VIII,184; IX,46; XI,11.
R. ciliata; Schauer in nov. act. acad. Caes. XXI, 11 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,171	M.fr.I,225; XI,11.
R. inops; Schauer in Lehm. pl. Preiss. II, 224 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,171	M.fr.XI,11.
PHYMATOCARPUS; F. v. M., fragm. III, 120 (1862).									
P. porphyrocephalus; F. v. M., fragm. III, 121 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,172	M.fr.III,121.
P. Maxwelli; F. v. M., fragm. IX, 45 (1875)...	W.A.	—	—	—	—	—	—	—	M.fr.IX,45.
CALOTHAMNUS; Labillardière, Nov. Holl. pl. specim. II, 25 (1806). (Billotia.)									
C. pachystachyus; Bentham, Fl. Austr. III, 173 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,173	
C. longissimus; F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,174	M.fr.III,112.
C. blepharantherus; F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,174	M.fr.X,31&55.
C. sanguineus; Labillardière, Nov. Holl. pl. spec. II, 25, t. 164 (1806)	W.A.	—	—	—	—	—	—	B.fl.III,174	M.fr.X,31.
C. torulosus; Schauer in nov. act. acad. Caes. XXI, 25 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,175	M.fr.X,31.
C. gibbosus; Bentham, Fl. Austr. III, 175 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,175	
C. gracilis; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)...	W.A.	—	—	—	—	—	—	B.fl.III,175	

C. blepharospermus; F. v. M., fragm. III, 111 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,175	M.fr.III,111.
C. Gilesii; F. v. M., fragm. X, 31 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,31.
C. chrysantherus; F. v. M., fragm. III, 112 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,176	M.fr.III,112.
C. Oldfieldii; F. v. M., fragm. III, 113 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,176	M.fr.III,113.
C. planifolius; Lehmann, delect. sem. hort. bot. Hamb. 7 (1842)	W.A.	—	—	—	—	—	—	B.fl.III,176	M.fr.X,31.
C. lateralis; Lindley, Bot. Regist. XXV, App. 9 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,177	
C. affinis; Turczaninow in Bullet. de St. Petersb. 428 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,177	M.fr.III,111; VIII,184.
C. Preissii; Schauer in nov. act. acad. Caes. XXI, 31 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,177	
C. Schaueri; Lehmann, delect. sem. hort. bot. Hamb. 7 (1842)	W.A.	—	—	—	—	—	—	B.fl.III,178	
C. Lehmanni; Schauer in nov. act. acad. Caes. XXI, 31 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,178	
C. villosus; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,178	
C. pinifolius; F. v. M., fragm. III, 153 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,178	M.fr.III,153.
C. rupestris; Schauer in nov. act. acad. Caes. XXI, 26 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,179	
C. quadrifidus; R. Brown in Ait. hort. Kew. sec. ed. IV, 418 (1812)	W.A.	—	—	—	—	—	—	B.fl.III,179	M.fr.X,31.
C. asper; Turczaninow in Bull. Mosc. XXII, 25 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,180	M.fr.X,31.
C. homalophyllus; F. v. M., fragm. III, 111 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,180	M.fr.X,31.
LAMARCHEA; Gaudichaud in Freyc. Voy. Bot. 483, t. 110 (1826).									
L. hakeaefolia; Gaudichaud in Freyc. Voy. 484, t. 110 (1826)	W.A.	—	—	—	—	—	—	B.fl.III,123	
EREMAEA; Lindley, Bot. Regist. XXV, App. 11 (1839).									
E. fimbriata; Lindley, Bot. Regist. XXV, App. 11 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,181	M.fr.II,30; XI,11.
E. acutifolia; F. v. M., fragm. II, 30 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,181	M.fr.II,30.
E. violacea; F. v. M., fragm. XI, 10 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,10.
E. pilosa; Lindley, Bot. Regist. XXV, App. 11 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,182	M.fr.II,30; XI,11.
E. ebracteata; F. v. M., fragm. II, 29 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.
E. beaufortioides; Bentham, Fl. Austr. III, 182 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,182	M.fr.XI,11.
ANGOPHORA; Cavanilles, icon. IV, 21, t. 331 (1797).									
A. cordifolia; Cavanilles, icon. IV, 21, t. 338 (1797)	—	—	—	—	N.S.W.	—	—	B.fl.III,183	
A. subvelutina; F. v. M., fragm. I, 31 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,184	M.fr.I,31.
A. intermedia; De Candolle, prodr. III, 222 (1828)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,184	
A. lanceolata; Cavanilles, icon. IV, 22, t. 339 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,184	
EUCALYPTUS; L'Heritier, sertum Anglicum 18 (1788). (Eudesmia, Symphyomyrtus.)									
I. Renantherae.									
E. stellulata; Sieber in De Candolle, prodr. III, 217 (1828)	—	—	—	V.	N.S.W.	—	—	B.fl.III,200	M.fr.II,45; IX,172.
E. panciflora; Sieber in Sprengel, cur. poster. 195 (1827)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,201	M.fr.II,52.
E. regnans; F. v. M. in Report Acclim. Soc. Vict. 20 (1870)	—	—	T.	V.	N.S.W.	—	—	—	
E. amygdalina; Labillardière, Nov. Holl. pl. spec. II, 14, t. 154 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,202	M.fr.II,53; VII,42&44.
E. coccifera; J. Hooker in Lond. Journ. of Bot. VI, 477 (1847)	—	—	T.	—	—	—	—	B.fl.III,204	
E. obliqua; L'Heritier, sertum Anglicum 18, t. 20 (1788)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,204	M.fr.II,44&171; XI,33.
E. stricta; Sieber in Spreng. syst. veg. cur. post. 195 (1827)	—	—	—	—	N.S.W.	—	—	B.fl.III,217	M.fr.XI,44.
E. virgata; Sieber in Spreng. syst. veg. cur. post. 195 (1827)	—	—	—	—	N.S.W.	—	—	B.fl.III,202	M.fr.XI,38.
E. Planchoniana; F. v. M., fragm. XI, 43 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,43.
E. Baileyana; F. v. M., fragm. XI, 37 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,37.
E. sepulcralis; F. v. M., Eucalyptographia eighth dec. (1882)	W.A.	—	—	—	—	—	—	—	
E. buprestium; F. v. M., fragm. III, 57 (1862)	W.A.	—	—	—	—	—	—	B.fl.III,205	M.fr.III,57.
E. marginata; Donn, hort. Cantabr. sec. edit. 101 (1800)	W.A.	—	—	—	—	—	—	B.fl.III,209	M.fr.II,40&174; VII,43; XI,38.
E. santalifolia; F. v. M. in Transact. Vict. Inst. I, 35 (1854)	W.A.	S.A.	—	—	—	—	—	B.fl.III,206	
E. macrorrhyncha; F. v. M., First general Report, 12 (1853)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,207	M.fr.XI,45.
E. capitellata; Smith in White, Journ. Voy. N. S. Wales 216 (1790)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,206	M.fr.II,173; XI,38.

E. eugenioides; Sieber in Spreng. cur. poster. 195 (1827)	—	—	—	V.	N.S.W.	Q.	—	—	M.fr.II,52,64.
E. piperita; Smith in White, Journ. Voy. N. S. Wales 226 (1790)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,207	M.fr.II,173; VII,42,44; XI,38.
E. pilularis; Smith in Transact. Linn. Soc. III, 284 (1797)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,208	M.fr.II,61&172; VII,44; XI,38.
E. triantha; Link, enum. pl. hort. bot. Berol. II, 20 (1822)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VII,42.
E. haemastoma; Smith in Transact. Linn. Soc. III, 285 (1797)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,212	M.fr.II,51&175; VII,43; XI,38.
E. Sieberiana; F. v. M., Eucalyptographia sec. decade (1879)	—	S.A.	T.	V.	N.S W.	—	—	B.fl.III,202	M.fr.XI,37.
E. microcorys; F. v. M., fragm. II, 50 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,212	M.fr.II,50.
II. Porantherae.									
E. paniculata; Smith in Transact. Linn. Soc. III, 287 (1797)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,211	M.fr.II,52,174; VII,43.
E. Leucoxylon; F. v. M. in Transact. Vict. Inst. I, 33 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,209	M.fr.II,60&175.
E. melliodora; Cunningham in Walp. rep. bot. syst. II, 924 (1843)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,210	M.fr.VII,43.
E. polyanthema; Schauer in Walpers, rep. bot. syst. II, 924 (1843)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,213	M.fr.VII,44.
E. populifolia; Hooker, icon. plant. 279 (1852)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,214&243	M.fr.II,67.
E. ochrophloia; F. v. M., fragm. XI, 36 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.XI,36.
E. gracilis; F. v. M. in Transact. Vict. Inst. I, 35 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,222	M.fr.II,55,57; VII,43; VIII,184; XI,36.
E. uncinata; Turczaninow in Bull. de Mosc. XXII, 23 (1849)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,216	
E. odorata; Behr in Schlecht. Linnaea XX, 657 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,215	M.fr.II,66.
E. largiflorens; F. v. M. in Transact. Vict. Inst. I, 34 (1854)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,215	M.fr.VII,43.
E. Behriana; F. v. M. in Transact. Vict. Inst. I, 34 (1854)	—	S.A.	—	V.	—	—	—	B.fl.III,214	
E. hemiphloia; F. v. M., fragm. II, 62 (1860)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,216	M.fr.II,62.
E. pruinosa; Schauer in Walp. rep. bot. syst. II, 926 (1843)	—	—	—	—	—	Q.	N.A.	B.fl.III,213	M.fr.III,132.
E. melanophloia; F. v. M. in Journ. Linn. Soc. III, 93 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,220	M.fr.VII,43.
III. Parallelantherae.									
E. Cloeziana; F. v. M., fragm. XI, 44 (1878)...	—	—	—	—	—	Q.	—	—	M.fr.XI,44.
E. Howittiana; F. v. M. in Wing's S. Sc. Record, Aug. (1882)	—	—	—	—	—	Q.	—	—	
E. drepanophylla; F. v. M. in Benth. Fl. Austr. III, 221 (1866)	—	—	—	—	—	Q.	—	B.fl.III,221	
E. crebra; F. v. M. in Journ. Linn. Soc. III, 87 (1858)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,221	M.fr.XI,44.
E. Staigeriana; F. v. M. in Bailey's synops. Q. L. Flora 176 (1883)	—	—	—	—	—	Q.	—	—	
E. decipiens; Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,218	
E. concolor; Schauer in Lehm. pl. Preiss. I, 129 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,247	M.fr.II,42.
E. siderophloia; Bentham, Fl. Austr. III, 220 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,220	M.fr.II,61; VII,44.
E. microtheca; F. v. M. in Journ. Linn. Soc. III, 87 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,223	M.fr.II,40; VII,42; X,100; XI,14.
E. Raveretiana; F. v. M., fragm. X, 99 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,99.
E. brachyandra; F. v. M. in Journ. Linn. Soc. III, 97 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,223	M.fr.X,100.
E. pachyphylla; F. v. M. in Journ. Linn. Soc. III, 98 (1858)	—	S.A.	—	—	—	Q.	N.A.	B.fl.III,237	
E. pyriformis; Turczaninow in Bull. Mosc. XXII, 22 (1849)	W.A.	S.A.	—	—	—	—	—	B.fl.III,226	M.fr.II,32; X,5&84.
E. macrocarpa; Hooker, icones plant. t. 405-407 (1842)	W.A.	—	—	—	—	—	—	B.fl.III,224	M.fr.II,41; XI,38.
E. Preissiana; Schauer in Lehm. pl. Preiss. I, 131 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,232	M.fr.II,38; VII,42.
E. alpina; Lindley in Mitch. Three Exped. II, 175 (1838)	—	—	—	V.	—	—	—	B.fl.III.225	M.fr.II,68; VII,42.
E. Globulus; Labillardière, Voy. I, 153, t. 13 (1799)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,225	M.fr.II,68.
E. megacarpa; F. v. M., fragm. II, 70 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,232	M.fr.VII,43.
E. cosmophylla; F. v. M. in Transact. Vict. Inst. 32 (1854)	—	S.A.	—	—	—	—	—	B.fl.III,225	M.fr.VII,43.
E. cordata; Labillardière, Nov. Holl. pl. spec. II, 13, t. 152 (1806)	—	—	T.	—	—	—	—	B.fl.III,224	
E. urnigera; J. Hooker in Lond. Journ. of Bot. VI, 477 (1847)	—	—	T.	—	—	—	—	B.fl.III,227	M.fr.II,43.
E. longifolia; Link, enum. pl. hort. bot. Berol. II, 29 (1822)	—	—	—	V.	N.S.W.	—	—	B.fl.III,226	M.fr.II,50,147; VII,43.
E. erythronema; Turczaninow in Bull. de l'acad. de St. Petersb. 415 (1852) ...	W.A.	—	—	—	—	—	—	B.fl.III,227	M.fr.VIII,184.
E. caesia; Bentham, Fl. Austr. III, 227 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,227	M.fr.XI,40.
E. tetraptera; Turczaninow in Bull. de Mosc. XXII, 22 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,228	M.fr.II,34; IV,172; VIII,184
E. miniata; A. Cunningham in Walp. rep. bot. syst. II, 925 (1843)	—	—	—	—	—	Q.	N.A.	B.fl.III,228	M.fr.XI,38&42.
E. phoenicea; F. v. M. in Journ. Linn. Soc. III, 91 (1858)...	—	—	—	—	—	—	N.A.	B.fl.III,231	

E. robusta; Smith, Specim. Bot. New Holl. 40, t. 13 (1793)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,228	M.fr.II,43&171; VII,44.		
E. botryoides; Smith in Transact. Linn. Soc. III, 286 (1797)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,229	M.fr.II,48&175; VII,43.		
E. goniocalyx; F. v. M. in Nederl. Kruidk. Arch. IV, 134 (1859)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,229	M.fr.II,48; IV,52.		
E. pallidifolia; F. v. M., fragm. III, 131 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,236	M.fr.III,131.		
E. incrassata; Labillardière, Nov. Holl. pl. spec. II, 12, t. 150 (1806)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,231	M.fr.II,59.		
E. oleosa; F. v. M. in Nederl. Kruidk. Arch. IV, 127 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,248	M.fr.II,56; XI,14.		
E. goniantha; Turczaninow in Bullet. de Mosc. XX, 163 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,248			
E. falcata; Turczaninow in Bullet. de Mosc. XX, 163 (1847)	W.A.	—	—	—	—	—	—	B.fl.III,248			
E. salmonophloia; F. v. M., fragm. XI, 11 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,11.		
E. leptopoda; Bentham, Fl. Austr. III, 238 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,238	M.fr.XI,11&13.		
E. salubris; F. v. M., fragm. X, 54 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.XI,12.		
E. angustissima; F. v. M., fragm. IV, 25 (1863)	W.A.	S.A.	—	—	—	—	—	B.fl.III,238	M.fr.IV,25.		
E. Doratoxylon; F. v. M., fragm. II, 55 (1860)	W.A.	—	—	—	—	—	—	B.fl.III,249	M.fr.II,55.		
E. decurva; F. v. M., fragm. III, 130 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,249	M.fr.III,130.		
E. Cooperiana; F. v. M., fragm. XI, 83 (1880)	W.A.	—	—	—	—	—	—	—	M.fr.XI,83.		
E. corynocalyx; F. v. M., fragm. II, 43 (1860)	—	S.A.	—	V.	—	—	—	B.fl.III,218	M.fr.VII,43.		
E. gomphocephala; De Candolle, prodr. III, 220 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,231	M.fr.II,36; V,14.		
E. Oldfieldii; F. v. M., fragm. II, 37 (1860)	W.A.	S.A.	—	—	—	—	—	B.fl.III,237	M.fr.II,37.		
E. orbifolia; F. v. M., fragm. V, 50 (1865)	W.A.	—	—	—	—	—	—	B.fl.III,238	M.fr.V,50.		
E. diversicolor; F. v. M., fragm. III, 131 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,251	M.fr.III,131; V,14; VII,42.		
E. patellaris; F. v. M. in Journ. Linn. Soc. III, 84 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,244			
E. tessellaris; F. v. M. in Journ. Linn. Soc. III, 88 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,251			
E. clavigera; A. Cunningham in Walp. rep. bot. syst. II, 926 (1843)	—	—	—	—	—	Q.	N.A.	B.fl.III,250	...	...	P.
E. ferruginea; Schauer in Walp. repert. II, 926 (1843)	—	—	—	—	—	—	N.A.	B.fl.III,254			
E. grandifolia; R. Brown in Benth. Fl. Austr. III, 250 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,250			
E. aspera; F. v. M. in Journ. Linn. Soc. III, 95 (1858)	—	—	—	—	—	—	N.A.	B.fl.III,250			
E. patens; Bentham, Fl. Austr. III, 247 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,247			
E. Todtiana; F. v. M. in Wing's S. Sc. Record II, 171 (1882)	W.A.	—	—	—	—	—	—	—			
E. vernicosa; J. Hooker in Lond. Journ. of Bot. VI, 478 (1847)	—	—	T.	—	—	—	—	B.fl.III,232			
E. Gunnii; J. Hooker in Lond. Journ. Bot. III, 499 (1844)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,246	M.fr.II,62; VII,43; XI,38.		
E. resinifera; Smith in White's Journ. Voy. N.S. Wales 231 (1790)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,245	M.fr.II,172; VII,43.		
E. saligna; Smith in Transact. Linn. Soc. III, 285 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,245	M.fr.II,173; VII,44; II,52.		
E. punctata; De Candolle, prodr. III, 217 (1828)	—	—	—	—	N.S.W.	—	—	B.fl.III,242	M.fr.VII,43.		
E. oligantha; Schauer in Walp. rep. II, 926 (1843)	—	—	—	—	—	—	N.A.	B.fl.III,213			
E. alba; Reinwardt in Blume, Bijdr. 1101 (1826)	—	—	—	—	—	Q.	N.A.	B.fl.III,243	...	...	As., P.
E. pulverulenta; Sims, Bot. Mag. t. 2089 (1819)	—	—	—	V.	N.S.W.	—	—	B.fl.III,224	M.fr.II,70.		
E. Stuartiana; F. v. M. in Benth. Fl. Austr. III, 243 (1866)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,243	M.fr.II,62.		
E. viminalis; Labillardière, Nov. Holl. pl. spec. II, 12, t. 151 (1806)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,239	M.fr.II,64; VII,44.		
E. rostrata; Schlechtendal, Linnaea XX, 655 (1847)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,240	M.fr.II,40,65; VII,44.		
E. tereticornis; Smith, Spec. Bot. New Holl. 41 (1793)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,241			
E. rudis; Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,244	M.fr.XI,14.		
E. foecunda; Schauer in Lehm. pl. Preiss. I, 130 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,252			
E. redunca; Schauer in Lehm. pl. Preiss. I, 127 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,253	M.fr.XI,15.		
E. grossa; F. v. M. in Benth. Fl. Austr. III, 232 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,232	M.fr.VII,41.		
E. obcordata; Turczaninow in Bull. Acad. Petersb. 416 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,234	M.fr.VII,44; VIII,184.		
E. occidentalis; Endlicher in Hueg. enum. 49 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,235	M.fr.II,39; VII,43.		
E. cornuta; Labillardière, Voy. I, 403, t. 20 (1799)	W.A.	—	—	—	—	—	—	B.fl.III,234	M.fr.II,39; V,210; XI,38.		
E. gamophylla; F. v. M., fragm. XI, 40 (1878)	—	S.A.	—	—	—	—	N.A.	—	M.fr.XI,40.		
E. perfoliata; R. Brown in Benth. Fl. Austr. III, 253 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,253			
E. setosa; Schauer in Walpers, rep. bot. syst. II, 926 (1843)	—	—	—	—	—	Q.	N.A.	B.fl.III,254	M.fr.III,123; X,106.		

E. Torelliana; F. v. M., fragm. X, 106 (1877)											Q.	—	—	M.fr.XI,42.
E. peltata; Bentham, Fl. Austr. III, 254 (1866)											Q.	—	B.fl.III,254	
E. latifolia; F. v. M. in Journ. Linn. Soc. III, 94 (1858)											Q.	N.A.	B.fl.III,255	
E. ptychocarpa; F. v. M. in Journ. Linn. Soc. III, 90 (1858)											—	N.A.	B.fl.III,255	M.fr.VII,44.
E. Abergiana; F. v. M., fragm. XI, 41 (1878)											Q.	—	—	M.fr.XI,41.
E. calophylla; R. Brown in Journ. Geogr. Soc. 20 (1831)					W.A.						—	—	B.fl.III,255	M.fr.II,35&171.
E. ficifolia; F. v. M., fragm. II, 85 (1860)					W.A.						—	—	B.fl.III,256	M.fr.II,85&176; VI,25; XI,15.
E. corymbosa; Smith, Specim. Bot. New Holl. 43 (1793)								V.	N.S.W.		Q.	N.A.	B.fl.III,256	M.fr.II,46&174; VII,43; XI,42.
E. terminalis; F. v. M. in Journ. Linn. Soc. III, 89 (1858)					W.A.	S.A.			N.S.W.		Q.	N.A.	B.fl.III,257	M.fr.XI,42.
E. Foelscheana; F. v. M. in Melb. Chemist, No. 55, p. 56 (1882)											—	N.A.	—	
E. dichromophloia; F. v. M. in Journ. Linn. Soc. III, 89 (1858)											—	N.A.	B.fl.III,257	
E. trachyphloia; F. v. M. in Journ. Linn. Soc. III, 90 (1858)											Q.	—	B.fl.III,221	M.fr.XI,42.
E. eximia; Schauer in Walpers, rep. bot. syst. II, 925 (1843)									N.S.W.		—	—	B.fl.III,258	
E. maculata; Hooker, icones plant. t. 619 (1844)									N.S.W.		Q.	—	B.fl.III,258	M.fr.II,47,147; VII,43; XI,45.
E. Watsoniana; F. v. M., fragm. X, 98 (1877)											Q.	—	—	M.fr.X,98.
E. odontocarpa; F. v. M. in Journ. Linn. Soc. III, 98 (1858)					W.A.						—	—	B.fl.III,260	
E. tetrodonta; F. v. M. in Journ. Linn. Soc. III, 97 (1858)											Q.	N.A.	B.fl.III,260	
E. erythrocorys; F. v. M., fragm. II, 33 (1860)					W.A.						—	—	B.fl.III,258	M.fr.II,33.
E. tetragona; F. v. M., fragm. VI, 51 (1864)					W.A.						—	—	B.fl.III,259	M.fr.II,37.
E. eudesmioides; F. v. M., fragm. II, 35 (1860)					W.A.						—	—	B.fl.III,260	M.fr.II,35.
TRISTANIA; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812). (Lophostemon, Tristaniopsis.)														
T. neriifolia; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)									N.S.W.		—	—	B.fl.III,261	M.fr.IV,56.
T. suaveolens; Smith in Rees, Cycl. XXXVI (1817)									N.S.W.		Q.	N.A.	B.fl.III,261	M.fr.I,81; IV,57.
T. conferta; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)									N.S.W.		Q.	N.A.	B.fl.III,263	M.fr.I,82; IV,57.
T. lactiflora; F. v. M., fragm. I, 82 (1858)											—	N.A.	B.fl.III,263	M.fr.I,82.
T. exiliflora; F. v. M., fragm. V, 11 (1865)											Q.	—	B.fl.III,264	M.fr.IX,190.
T. laurina; R. Brown in Ait. hort. Kew. sec. ed. IV, 417 (1812)								V.	N.S.W.		Q.	—	B.fl.III,264	M.fr.I,81; IV,56.
T. psidioides; Cunningham in Bot. Regist. XXII, n. 1839 (1837)											—	N.A.	B.fl.III,264	
T. umbrosa; Cunningham in Bot. Regist. XXII, n. 1839 (1837)											—	N.A.	B.fl.III,265	
T. longivalvis; F. v. M. in Wing's S. Sc. Record, new ser. vol. II, April (1886)											Q.	N.A.	—	
METROSIDEROS; Banks in Gaertner, de fruct. I, 170, t. 34 (1788). (Syncarpia, Kamptzia, Lysicarpus, Xanthostemon, Nania, Fremya, Cloezia, Spermalepis.)														
M. glomulifera; Smith in Transact. Linn. Soc. III, 269 (1797)									N.S.W.		Q.	—	B.fl.III,265	M.fr.I,79.
M. leptopetala; F. v. M., Docum. Intercol. Exhib. 30 (1867)									N.S.W.		Q.	—	B.fl.III,266	M.fr.I,79.
M. ternifolia; F. v. M., Docum. Intercol. Exhib. 30 (1867)											Q.	—	B.fl.III,267	M.fr.I,82.
M. eucalyptoides; F. v. M., fragm. I, 243 (1858)											—	N.A.	B.fl.III,267	M.fr.I,81.
M. polymorpha; Gaudichaud, Freyc. Voy. Bot. 99, t. 108 (1826)									N.S.W.		—	—	—	M.fr.VIII,14 ... ... P.
M. nervulosa; C. Moore & F. v. M., fragm. VIII, 15 (1873)									N.S.W.		—	—	—	M.fr.I,80.
M. chrysantha; F. v. M., fragm. IV, 159 (1864)											Q.	—	B.fl.III,268	M.fr.IV,159.
M. paradoxa; F. v. M., fragm. I, 243 (1859)											—	N.A.	B.fl.III,269	M.fr.I,80.
M. tetrapetala; F. v. M., fragm. VII, 41 (1870)											Q.	N.A.	—	M.fr.VII,41.
M. pachysperma; F. v. M. & Bailey in Bailey's occas. papers on Q. Flora I, 4 (1886)											Q.	—	—	
BACKHOUSIA; Hooker and Harvey in Bot. Mag. t. 4133 (1845).														
B. myrtifolia; Hooker & Harvey in Bot. Mag. t. 4133 (1845)								V.	N.S.W.		Q.	—	B.fl.III,269	M.fr.I,78; IX,43.
B. angustifolia; F. v. M., fragm. I, 79 (1858)											Q.	—	B.fl.III,270	M.fr.IX,43.
B. sciadophora; F. v. M., fragm. II, 26 (1860)									N.S.W.		Q.	—	B.fl.III,270	M.fr.II,171.
B. citriodora; F. v. M., fragm. I, 78 (1858)											Q.	—	B.fl.III,270	M.fr.I,78.
B. Bancroftii; F. v. M. & Bailey in catal. of Queensland Woods 46 (1886)											Q.	—	—	
OSBORNIA; F. v. M., fragm. III, 30 (1862).														
O. octodonta; F. v. M., fragm. III, 31 (1862)											Q.	N.A.	B.fl.III,271	M.fr.IX,143 ... ... As., P.

RHODOMYRTUS; De Candolle, prodr. III, 240 (1828).

R. psidioides; Bentham, Fl. Austr. III, 272 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,272	M.fr.II,86; III,165; IX,142.		
R. trineura; F. v. M. in Benth. Fl. Austr. III, 272 (1866)	—	—	—	—	—	Q.	—	B.fl.III,272	M.fr.IV,117; IX,142.		
R. cymiflora; F. v. M. in Benth. Fl. Austr. III, 273 (1866)	—	—	—	—	—	Q.	—	B.fl.III,273	M.fr.V,12; IX,149.		
R. macrocarpa; Bentham, Fl. Austr. III, 273 (1866)	—	—	—	—	—	Q.	—	B.fl.III,273	M.fr.IX,142.		

MYRTUS; Tournefort, inst. 640, t. 409 (1700), from Plinius.

M. rhytisperma; F. v. M., fragm. I, 77 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,274	M.fr.IX,147.		
M. tenuifolia; Smith in Transact. Linn. Soc. III, 280 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,274	M.fr.IX,147.		
M. gonoclada; F. v. M. in Benth. Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,275	M.fr.IX,144.		
M. lasioclada; F. v. M., fragm. IX, 148 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,148.		
M. Hillii; Bentham, Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,275	M.fr.IX,149.		
M. Becklerii; F. v. M., fragm. II, 85 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,272	M.fr.IX,147.		
M. Bidwillii; Bentham, Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,275			
M. racemulosa; Bentham, Fl. Austr. III, 276 (1866)	—	—	—	—	—	Q.	—	B.fl.III,276	M.fr.IX,147.		
M. acmenioides; F. v. M., Fragm. I, 77 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,276	M.fr.IX,147.		
M. fragrantissima; F. v. M. in Benth. Fl. Austr. III, 277 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,277			
M. Shepherdi; F. v. M., fragm. IX, 148 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,148.		

RHODAMNIA; Jack in Hook. Comp. to the Bot. Mag. I, 153 (1835).

R. sessiliflora; Bentham, Fl. Austr. III, 275 (1866)	—	—	—	—	—	Q.	—	B.fl.III,277	M.fr.IX,142.		
R. trinervia; Blume, Mus. Bot. Lugd. I, 79 (1849)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,278	M.fr.I,76; IX,142.		
R. argentea; Bentham, Fl. Austr. III, 276 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,278	M.fr.IX,142.		
R. Blairiana; F. v. M., fragm. IX, 141 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,141.		

FENZLIA; Endlicher in Atacta 19, t. 17 & 18 (1833).

F. obtusa; Endlicher, Atacta 19, t. 17 (1833)	—	—	—	—	—	Q.	—	B.fl.III,279	M.fr.IX,142.		
F. retusa; Endlicher, Atacta 20, t. 18 (1833)	—	—	—	—	—	Q.	N.A.	B.fl.III,279			

DECASPERMUM; R. & G. Forster, char. gen. 73, t. 37 (1776). (Nelitris.)

D. paniculatum; Baillon, hist. des plant. 355 (1871)	—	—	—	—	—	Q.	—	B.fl.III,279	M.fr.IV,56; IX,149	...	As.

EUGENIA; Micheli, nov. pl. gen. 226, t. 108 (1729). (Plinia, Plumier 1703, Acmene, Jambosa, Syzygium, Cleistocalyx partly.)

E. rariflora; Bentham in Hook. Lond. Journ. II, 221 (1843)	—	—	—	—	—	Q.	—	B.fl.III,282	M.fr.III,130; V,15; IX,143.		P.
E. Smithii; Poiret, Encycl. méthod. Suppl. III, 126 (1813)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.III,282	M.fr.IV,59; IX,145.		
E. hemilampra; F. v. M., fragm. IX, 145 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,145.		
E. Ventenatii; Bentham, Fl. Austr. III, 283 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,283	M.fr.IV,58; IX,145.		
E. leptantha; Wight, Illustr. II, 15 (1850)	—	—	—	—	—	Q.	—	B.fl.III,283	M.fr.IX,143	...	As.
E. Moorei; F. v. M., fragm. V, 33 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,283	M.fr.IX,143.		
E. corynantha; F. v. M., fragm. IX, 144 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,144.		
E. cormiflora; F. v. M., fragm. V, 32 (1865)	—	—	—	—	—	Q.	—	B.fl.III,284	M.fr.V,32.		
E. Hodgkinsoniae; F. v. M., fragm. IX, 145 (1875)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,145.		
E. Tierneyana; F. v. M., fragm. V, 14 (1865)	—	—	—	—	—	Q.	—	B.fl.III,284	M.fr.IX,143.		
E. grandis; Wight, Illustr. II, 17 (1850)	—	—	—	—	—	Q.	—	B.fl.III,285	M.fr.V,13; IX,143	...	As.
E. suborbicularis; Bentham, Fl. Austr. III, 285 (1866)	—	—	—	—	—	Q.	—	B.fl.III,285	M.fr.IX,143.		
E. Wilsonii; F. v. M., fragm. V, 12 (1865)	—	—	—	—	—	Q.	—	B.fl.III,285	M.fr.IX,143.		
E. eucalyptoides; F. v. M., fragm. IV, 55 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.III,285	M.fr.I,226; IX,143.		
E. Australis; Wendland in Link, enum. pl. hort. bot. II, 28 (1822)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,286	M.fr.I,225; V,33; IX,143.		
E. angophoroides; F. v. M., fragm. V, 33 (1865)	—	—	—	—	—	Q.	—	B.fl.III,286	M.fr.IX,145.		
E. Armstrongii; Bentham, Fl. Austr. III, 286 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,286			
E. oleosa; F. v. M., fragm. V, 15 (1865)	—	—	—	—	—	Q.	—	B.fl.III,287	M.fr.V,33; IX,146.		
E. Dallachiana; F. v. M. in Benth. Fl. Austr. III, 287 (1866)	—	—	—	—	—	Q.	—	B.fl.III,287	M.fr.IX,145.		
E. cryptophlebia; F. v. M., fragm. IX, 144 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,144.		
E. Holtzeana; F. v. M. in Austr. Journ. of Pharm. I, 199 (1886)	—	—	—	—	—	—	N.A.	—			

ACICALYPTUS ; A. Gray, New Gen. Un. St. Exped. 9 (1853). (Calyptranthes partly, Cleistocalycis subgenus.)

A. Fullagari ; F. v. M., fragm. VIII, 15 (1873)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,281.			

BARRINGTONIA ; R. & G. Forster, char. gen. 75, t. 38 (1776). (Stravadium.)

B. speciosa ; R. & G. Forster, char. gen. 76, t. 38 (1776)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,288	M.fr.IX,118	...	...	As., P.
B. racemosa ; Gaudichaud in Freyc. Voy. Bot. 483, t. 107 (1826)	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,118	...	...	As., P.	
B. acutangula ; Gaertner, de fruct. II, 97, t. 101 (1791)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,288	M.fr.IX,119	...	...	As.

CAREYA ; Roxburgh, pl. Corom. III, 13, t. 217 (1816).

C. Australis ; F. v. M., fragm. V, 183 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,289	M.fr.V,183.		

SONNERATIA ; Linné, fil. suppl. plant. 38 (1781).

S. acida ; Linné, fil. suppl. plant. 38 (1781)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,301	M.fr.III,109	...	...	As., Afr.

MELASTOMACEAE.

A. L. de Jussieu, Gen. plant. 328 (1789).—Genera 5, Species 7.

MEMECYLON ; Linné, Fl. Zeyl. app. 9 (1747).

M. umbellatum ; N. L. Burmann, Fl. Ind. 87 (1768)	...	...	...	..	—	—	—	—	—	Q.	N.A.	B.fl.III,293	...	...	...	As.

OSBECKIA ; Linné, spec. plant. I, 345 (1753).

O. Chinensis ; Linné, spec. plant. 345 (1753)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,291	M.fr.IV,160	...	...	As.
O Australiana ; Naudin in Ann. sc. Nat. trois. sér. XIV, 59 (1855)	...	...	...	—	—	—	—	—	—	—	N.A.	B.fl.III,291	M.fr.IV,161.				
O. perangusta ; F. v. M., fragm. V, 181 (1866)	...	...	...	..	...	—	—	—	—	—	—	N.A.	—	M.fr.V,181.			

OTANTHERA ; Blume in Flora, Regensb. IV, 488 (1831).

O. bracteata ; Korthals in Verh. Nat. Gesch. der Ned. Bes. 235, t. 51 (1841)	...	—	—	—	—	—	Q.	—	B.fl.III,292	...	...	..	P.				

MELASTOMA ; J. Burmann, thesaur. Zeyl. 72 (1737).

M. Malabathricum ; Linné, spec. plant. 390 (1753)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,292	M.fr.IV,161	...	..	As., P.

MEDINILLA ; Gaudichaud in Freycin. voy. Bot. 484, t. 106 (1826).

Balls-Headleyi ; F. v. M. in Vict. Naturalist new ser. II, 125 (1887)	...	—	—	—	—	—	Q.	—	—								

RHAMNACEAE.

A. L. de Jussieu, Gen. plant. 376 (1789), from B. de Jussieu (1759).—Genera 10, Species 90.

VENTILAGO ; Gaertner, de fruct. I, 223, t. 49 (1788).

V. viminalis ; Hooker in Mitch. Trop. Austr. 369 (1848)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.I,411	M.fr.III,63; IX,140.				
V. ecorollata ; F. v. M. in Wing's S. Sc. Rec. III, 176 (1883)	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,141.					

ZIZYPHUS ; Tournefort, inst. 627, t. 403 (1700), from Plinius. (Ziziphus.)

Z. Oenoplia ; Miller, Gardeners' Dictionary n. 3 (1731)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,412					
Z. Jujuba ; Lamarck, Encycl. méth. III, 318 (1789)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.I,412	..	...	..	...	As., Afr.
Z. quadrilocularis ; F. v. M., fragm. III, 57 (1862)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.I,412	M.fr.III,57.				

DALLACHYA ; F. v. M., fragm. IX, 140 (1875). (Rhamnus partly.)

D. Vitiensis ; F. v. M., fragm. IX, 140 (1875)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.XI,134	...	...	P.

COLUBRINA ; L. C. Richard in Ann. des sc. nat. X, 368 (1827).

C. Asiatica ; Brogniart in Ann. des sc. nat. X, 369 (1827)	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.I,413	M.fr.III,63; IX,139	...	As., Afr., P		

ALPHITONIA ; Reisseck in Endl. gen. plant. 1098 (1840).

A. excelsa ; Reisseck in Endl. gen. plant. 1098 (1840)	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.I,414	M.fr.III,63; IX,140	...	As., P.		

EMMENOSPERMA; F. v. M., fragm. III, 63 (1862).

E. alphitonioides; F. v. M., fragm. III, 63 (1862)	--	—	—	—	N.S.W.	Q.	—	B.fl.I,415	M.fr.IX,140.
E. Cunninghamii; Bentham, Fl. Austr. I, 415 (1863)	—	—	—	—	—	—	N.A.	B.fl.I,415	

POMADERRIS; Labillardière, Nov. Holl. pl. specim. I, 61, t. 86 (1804). (Pomatoderris.)

P. lanigera; Sims, Bot. Mag. t. 1823 (1816)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,416	
P. elliptica; Labillardière, Nov. Holl. pl. spec. I, 61, t. 86 (1804)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,417	M.fr.III,69; IX,139.
P. phillyroides; Sieber in De Candolle, prodr. II, 33 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,418	
P. vaccinifolia; Reisseck & F. v. M. in Linnaea XXIX, 266 (1857)	—	—	—	V.	N.S.W.	—	—	B.fl.I,418	M.fr.III,71.
P. myrtilloides; Fenzl in Hueg. enum. 22 (1837)	W.A.	S.A.	—	—	—	—	—	B.fl.I,419	M.fr.IX,138.
P. grandis; F. v. M., fragm. III, 68 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,417	M.fr.IX,139.
P. Calvertiana; F. v. M., fragm. IX, 138 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,138.
P. Iedifolia; Cunningham in Field, N.S.W. 351 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.I,419	M.fr.III,69.
P. apetala; Labillardière, Nov. Holl. pl. spec. I, 62, t. 87 (1804)	--	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,419	M.fr.III,69&73; IX,139.
P. cinerea; Bentham, Fl. Austr. I, 420 (1863)	—	—	—	V.	N.S.W.	—	—	B.fl.I,420	
P. prunifolia; Cunningham in Hueg. enum. 22 (1837)	—	—	—	V.	N.S.W.	Q	—	B.fl.I,420	M.fr.III,75; IX,139.
P. ligustrina; Sieber in De Candolle, prodr. II, 34 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.I,420	M.fr.III,71; IX,139.
P. betulina; Cunningham in Bot. Mag. t. 3212 (1833)	—	—	—	V.	N.S.W.	—	—	B.fl.I,421	M.fr.III,76; IX,139.
P. Forrestiana; F. v. M., fragm. IX, 139 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,139.
P. obcordata; Fenzl in Hueg. enum. 23 (1837)	W.A.	S.A.	—	V.	—	—	—	B.fl.I,421	M.fr.III,73.
P. racemosa; Hooker, Journ. of Bot. I, 256 (1834)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,421	M.fr.III,75&168; IX,139.
P. subrepanda; Reisseck & F. v. M. in Linnaea XXIX, 267 (1857)	—	—	—	V.	—	—	—	B.fl.I,422	M.fr.III,74.
P. clachophylla; F. v. M., fragm. II, 131 (1861)	—	—	—	V.	—	—	—	B.fl.I,422	M.fr.III,72,166; IX,139 ... N.Z.
P. phylicifolia; Loddiges, Bot. Cabinet t. 120 (1818)	—	—	T.	V.	N.S.W.	—	—	B.fl.I,422	M.fr.III,73.
P. intangenda; F. v. M., fragm. X, 52 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,53.

CRYPTANDRA; Smith in Transact. Linn. Soc. IV, 217 (1798). (Trymalium, Spyridium, Stenanthemum, Wichurea, Stenodiscus.)

C. albicans; F. v. M., Cens. 60 (1882) from Reisseck (1847)	W.A.	—	—	—	—	—	—	B.fl.I,423	
C. Billardieri; F. v. M., Cens. 60 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,423	
C. ledifolia; F. v. M., Cens. 60 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,424	
C. Daltoni; F. v. M., Cens. 60 (1882)	—	—	—	V.	—	—	—	—	M.fr.IX,135,198.
C. angustifolia; F. v. M., Cens. 60 (1882) from Reisseck (1847)	W.A.	—	—	—	—	—	—	B.fl.I,424	M.fr.IX,135.
C. Wichureae; F. v. M., Cens. 60 (1882) from Nees (1847)	W.A.	—	—	—	—	—	—	B.fl.I,425	M.fr.IX,135.
C. ericifolia; Smith in Rees, Cycloped. (1808)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,438	M.fr.III,66;IX,137.
C. propinqua; Cunningham in Hueg. enum. 23 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,442	M.fr.III,67;IX,137.
C. hispidula; Reisseck & F. v. M. in Linnaea XXIX, 294 (1857)	—	S.A.	—	—	—	—	—	B.fl.I,439	
C. spyridioides; F. v. M., fragm. III, 68 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,439	M.fr.IX,138.
C. scoparia; Reisseck in Lehm. pl. Preiss. II, 283 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,439	
C. spinescens; Sieber in De Candolle, prodr. II, 38 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.I,439	
C. amara; Smith in Transact. Linn. Soc. X, 245, t. 18 (1811)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.I,440	M.fr.III,66&67; IX,137.
C. lanosiflora; F. v. M., fragm. III, 65 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,440	M.fr.III,65.
C. tomentosa; Lindley in Mitch. Three Exped. II, 178 (1838)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.I,441	M.fr.III,67; IX,138.
C. nutans; Steudel in Lehm. pl. Preiss. I, 186 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,441	M.fr.IX,138.
C. glabriflora; Bentham, Fl. Austr. I, 441 (1863)	W.A.	—	—	—	—	—	—	B.fl.I,441	
C. alpina; J. Hooker, Fl. Tasm. I, 75, t. 12 (1860)	—	—	T.	—	—	—	—	B.fl.I,441	
C. Leucopogon; Meissner in Lehm. pl. Preiss. II, 287 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,442	
C. parvifolia; Turczaninow in Bull. de Mosc. I, 459 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,442	
C. buxifolia; Fenzl in Hueg. enum. 23 (1837)	—	—	—	—	N.S.W.	—	—	B.fl.I,442	M.fr.IX,138.
C. pungens; Steudel in Lehm. pl. Preiss. I, 187 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,443	M.fr.IX,138.
C. mutila; Nees in Lehm. pl. Preiss. II, 289 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,443	
C. longistaminea; F. v. M., fragm. III, 64 (1862)	—	—	—	—	N.S.W.	—	—	B.fl.I,443	M.fr.IX,138.

C. arbutiflora; Fenzl in Hueg. enum. 26 (1837)	W.A.	—	—	—	—	—	—	B.fl.I,444	M.fr.III,64; IX,138.
C. miliaris; Reisseck in Lehm. pl. Preiss. II, 288 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,444	
C. nudiflora; F. v. M., fragm. III, 64 (1862)	W.A.	—	—	—	—	—	—	B.fl.I,444	M.fr.III,64.
C. ulicina; Hooker, Journ. of Bot. I, 257 (1834)	—	—	T.	—	—	—	—	B.fl.I,434	M.fr.IX,137.
C. pomaderroides; Reisseck in Endl. nov. stirp. dec. 29 (1839)	W.A.	—	—	—	—	—	—	B.fl.I,435	M.fr.IX,137.
C. pumila; F. v. M., Cens. 61 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.IX,137.
C. leucophracta; Schlechtendal, Linnaea XX, 640 (1847)	—	S.A.	T.	V.	—	—	—	B.fl.I,435	M.fr.III,77; IX,136.
C. coronata; Reisseck in Lehm. pl. Preiss. II, 288 (1847)	W.A.	—	—	—	—	—	—	B.fl.I,436	M.fr.IX,137.
C. Scortechinii; F. v. M. in Melb. Chem. VI, 71 (1884)	—	—	—	V.	N.S.W.	Q.	—	—	
C. humilis; F. v. M., Cens. 61 (1882) from Bentham (1863)	W.A.	—	—	—	—	—	—	B.fl.I,436	
C. Waterhousii; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I,436	M.fr.III,83.
C. tridentata; Steudel in Lehm. pl. Preiss. I, 186 (1844)	W.A.	—	—	—	—	—	—	B.fl.I,427	M.fr.IX,137.
C. divaricata; F. v. M., Cens. 61 (1882) from Bentham (1863)	W.A.	—	—	—	—	—	—	B.fl.I,427	M.fr.III,68.
C. obcordata; J. Hooker, Fl. Tasm. I, 71 (1860)	—	—	T.	V.	—	—	—	B.fl.I,427	M.fr.III,80.
C. Lawrencii; J. Hooker, Fl. Tasm. I, 72 (1860)	—	—	T.	—	—	—	—	B.fl.I,430	M.fr.IX,136.
C. Hookeri; F. v. M., Cens. 61 (1882)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,428	M.fr.III,79; IX,136.
C. spadicea; F. v. M., Cens. 61 (1882) from Fenzl (1837)	W.A.	—	—	—	—	—	—	B.fl.I,428	M.fr.IX,137.
C. globulosa; F. v. M., Cens. 61 (1882) from G. Don (1830)	W.A.	—	—	—	—	—	—	B.fl.I,429	M.fr.IX,137.
C. obovata; J. Hooker, Fl. Tasm. I, 74 (1860)	—	S.A.	T.	—	—	—	—	B.fl.I,429	M.fr.IX,136.
C. spathulata; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I,430	M.fr.IX,136.
C. cordata; Turczaninow in Bull. de Mosc. XXXI, 459 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,430	M.fr.IV,25; IX,137.
C. rotundifolia; F. v. M., Cens. 61 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.IV,25.
C. phlebophylla; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I,430	
C. coactilifolia; F. v. M., Cens. 61 (1882) from Reiss. (1856)	—	S.A.	—	—	—	—	—	B.fl.I,431	M.fr.IX,136.
C. complicata; F. v. M., Cens. 61 (1882)	W.A.	—	—	—	—	—	—	B.fl.I,431	M.fr.III,78.
C. westringifolia; F. v. M., Cens. 61 (1882) from Reiss. (1847)	W.A.	—	—	—	—	—	—	B.fl.I,431	
C. villosa; Turczaninow in Bull. de Mosc. XXXI, 458 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,432	
C. pauciflora; Turczaninow in Bull. de Mosc. XXXI, 458 (1858)	W.A.	—	—	—	—	—	—	B.fl.I,432	
C. halmaturina; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I,432	
C. bifida; F. v. M., Cens. 61 (1882)	—	S.A.	—	—	—	—	—	B.fl.I,432	M.fr.IX,136.
C. subochreata; F. v. M., Cens. 61 (1882)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.I,432	M.fr.III,82; IX,136.
C. vexillifera; Hooker, Journ. Bot. I, 257 (1834)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.I,433	M.fr.III,81; IX,136.
C. Wayii; F. v. M. and Tate in Transact. R. Soc. S.A. V, 89 (1882)	—	S.A.	—	—	—	—	—	—	
COLLETIA; Commerçon in A. L. de Jussieu, gen. 380 (1789). (Discaria, Tetrapasma.)									
C. pubescens; Brogniart in Annal. des sc. nat. X, 366 (1827)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.I,445	M.fr.III,83; IV,175; IX,140.
GOUANIA; N. J. Jacquin, stirp. Amer. hist. 263, t. 179 (1763).									
G. Hillii; F. v. M., fragm. IX, 163 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,163.
G. Australiana; F. v. M., fragm. IV, 144 (1864)	—	—	—	—	—	Q.	—	—	M.fr.IV,144.

VINIFERAE.

J. de St. Hilaire, Exp. fam. II, 48, t. 79 (1805).—Genus 1, Species 17.

VITIS; Tournefort, inst. 613, t. 384 (1700), e Latinis. (Cissus.)									
V. Baudiniana; F. v. M., fragm. IV, 136 (1864)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,447	M.fr.V,212; VI,178.
V. oblonga; Bentham, Fl. Austr. I, 447 (1863)	—	—	—	—	—	Q.	—	B.fl.I,447	M.fr.IV,136; VI,178; IX,125.
V. repens; Wight & Arnott, prodr. fl. penins. Ind. I, 186 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.I,447	M.fr.II,73; V,210; VI,178; IX,126. As.
V. adnata; Wallich in Wight & Arnott, prodr. 126 (1834)	—	—	—	—	—	—	N.A.	B.fl.I,448	M.fr.VI,178. ... ... As., P
V. nitens; F. v. M., fragm. II, 73 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,448	M.fr.VI,178.
V. brachypoda; F. v. M., fragm. IX, 125 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,125.

V. saponaria ; Seemann, Bonplandia, 254 (1859)	—	—	—	—	—	—	Q.	—	B.fl.I,448	M.fr.VI,178.		
V. acris ; F. v. M., fragm. II, 75 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,449	M.fr.VI,178.		
V. trifolia ; Linné, spec. plant. 203 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.I,449	M.fr.II,75; VI,177; IX,126	...	As., P.
V. clematidea ; F. v. M., fragm. II, 74 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,449	M.fr.V,210; VI,177; IX,126.		
V. Japonica ; Thunberg, Fl. Jap. 104 (1775) ...	—	—	—	—	—	Q.	—	—	... ...	...	As., P.
V. acetosa ; F. v. M., plants of Vict. I, 44 (1860)	—	—	—	—	—	Q.	N.A.	B.fl.I,449	M.fr.VI,177; IX,126.		
V. hypoglauca ; F. v. M., plants of Vict. I, 95 (1860)	—	—	—	V.	N.S.W.	Q.	—	B.fl.I,450	M.fr.VI,177.		
V. sterculifolia ; F. v. M. in Benth. Fl. Austr. I, 450 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,450	M.fr.IX,126.		
V. penninervis ; F. v. M., fragm. VI, 177 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,177.		
V. opaca ; F. v. M. in Benth. Fl. Austr. I, 450 (1863)	—	—	—	—	N.S.W.	Q.	—	B.fl.I,450	M.fr.VI,177; IX,125.		
V. augustissima ; F. v. M., fragm. I, 141 (1859)	W.A.	—	—	—	—	—	—	B.fl.I,450	M.fr.I,244; II,180;IX,126.		

LEEACEAE.

F. v. M. from De Candolle, prodr. I, 635 (1824).—Genus 1, Species 2.

LEEA ; Linné, mantiss. 17 (1767).

L. staphylea ; Roxburgh, hort. Beng. 18 (1814)	—	—	—	—	—	Q.	—	B.fl.I,451	M.fr.VI,178	...	As.
F. Brunouiana ; Clarke in Britten's journ. of Bot. X, 105 (1881)	—	—	—	—	—	Q.	N.A.	—	M.fr.VI,178.		

ARALIACEAE.

Ventenat, Tabl. III, (1799). (Umbelliferarum subordo.)—Genera 10, Species 22.

ASTROTRICHA ; De Candolle, Mémoire snr la fam. des Ombellifères 25 (1829).

A. pterocarpa ; Bentham, Fl. Austr. III, 379 (1866)...	—	—	—	—	—	Q.	—	B.fl.III,379	M.fr.VII,148.		
A. floccosa ; De Candolle, Mém. des Ombell. 30, t. 5 (1829)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,380			
A. longifolia ; Bentham in Hueg. enum. 55 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,380	M.fr.VII,148.		
A. ledifolia ; De Candolle, Mém. des Ombell. 30, t. 6 (1829)	—	—	—	V.	N.S.W.	—	—	B.fl.III,380	M.fr.VII,148.		
A. Hamptoni ; F. v. M., fragm. VI, 125 (1868)	W.A.	—	—	—	—	—	N.A.	—	M.fr.VI,125&250.		

DELARBREA ; Vieillard in Bullet. de la Soc. Linn. Normand. IX, 342 (1865). (Porospermum.)

D. Michieana ; F. v. M., fragm. VII, 95 (1870)	—	—	—	—	—	Q.	—	—	M.fr.VII,95.		

PANAX ; Linné, spec. plant. edit. sec. 1513 (1763). (Nothopanax.)

P. Gunnii ; J. Hooker in Lond. Journ. of Bot. VI, 466 (1847)	—	—	T.	—	—	—	—	B.fl.III,381			
P. Murrayi ; F. v. M., fragm. II, 106 (1861) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,381	M.fr.VII,96.		
P. mollis ; Bentham, Fl. Austr. III, 382 (1866)	—	—	—	—	—	Q.	—	B.fl.III,382	M.fr.VII,96.		
P. Macgillivraei ; Bentham, Fl. Austr. III, 382 (1866)	—	—	—	—	—	Q.	—	B.fl.III,382			
P. sambucifolius ; Sieber in De Candolle, prodr. III, 255 (1825)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,382	M.fr.II,107; VII,95.		
P. cephalobotrys ; F. v. M., fragm. II, 83 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,382	M.fr.II,83,107,176; VII,95.		
P. elegans ; C. Moore and F. v. M. in Trans. phil. Inst. Vict. II, 68 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,383	M.fr.II,107.		
P. Cissodendron ; C. Moore and F. v. M., fragm. VII, 96 (1870) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,280; IX,96.		

PENTAPANAX ; Seemann, Journal of Botany II, 294 (1864).

P. Willmottii ; F. v. M. in Vict. Naturalist, new ser. II, 125 (1887)	—	—	—	—	—	Q.	—	—			

MOTHERWELLIA ; F. v. M., fragm. VII, 107 (1871).

M. haplosciadea ; F. v. M., fragm. VII, 107 (1871) ...	—	—	—	—	—	Q.	—	—	M.fr.VII,107.		

CISSODENDRON ; Seemann, Journ. of Bot. II, 303 (1864). (Hedera partly, Irvingia, Kissodendron.)

C. Australianum ; Seemann, Journ. of Bot. II, 303 (1864) ...	—	—	—	—	—	Q.	—	B.fl.III,384	M.fr.IV,120; V,18.		

HEPTAPLEURUM ; Gaertner, de fruct. II, 472, t. 178 (1791). (Paratropia.)

H. venulosum ; Seemann, Journ. of Bot. III, 80 (1865)	—	—	—	—	—	Q.	—	B.fl.III,384	M.fr.II,108...	...	As.

BRASSAIA; Endlicher, nov. stirp. decad. I, 89 (1839).

B. actinophylla; Endlicher, nov. stirp. dec. I, 89 (1839)	—	—	—	—	—	—	—	—	—	Q.	—	B.fl.III,385	M.fr.II,108; IV,121.		

MERYTA; R. & G. Forster, char. gen. 119, t. 60 (1776). (Bothryodendron.)

M. latifolia; Seemann in Bonplandia, 295 (1862)	—	—	—	—	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.		
M. angustifolia; Seemann in Bonplandia, 296 (1862)	—	—	—	—	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.		

MACKINLAYA; F. v. M., fragm. IV, 120 (1864).

M. macrosciadea; F. v. M., fragm. IV, 120 (1864)	—	—	—	—	—	—	—	—	—	Q.	—	B.fl.III,383	M.fr.II,108&176.		

UMBELLIFERAE.

MORISON, hist. pl. II, lib. 3, sect. 9 (1680).—Genera 16, Species 106.

HYDROCOTYLE; Tournefort, inst. 328, t. 173 (1700).

H. vulgaris; Linné, spec. plant. 234 (1753)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,339	M.fr.IV,180	...	E., As., Afr., Am.				
H. hirta; R. Brown in annal. des sc. phys. VI, 64 (1820)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,339	M.fr.IV,181.						
H. Candollei; F. v. M., key to the syst. of Vict. pl. I, 267 (1888)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,340	M.fr.IV,179&180.						
H. rhombifolia; F. v. M., fragm. VII, 147 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,147.						
H. pedicellosa; F. v. M., fragm. IV, 182 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,341	M.fr.VII,147.						
H. tripartita; R. Brown in annal. des sc. phys. VI, 69, t. 61 (1820)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,341	M.fr.IV,181; VII,147.						
H. comocarpa; F. v. M. in Vict. Nat. III, 126 (1887)	—	S.A.	—	—	—	—	—	—							
H. crassiuscula; Tate in Transact. R. S. of S.A. VI, 140	—	S.A.	—	—	—	—	—	—							
H. plebeja; R. Brown in annal. des sc. phys. VI, 46, t. 60 (1820)	W.A.	—	—	—	—	—	—	B.fl.III,341							
H. pterocarpa; F. v. M. in Transact. Vict. Inst. 126 (1855)	—	S.A.	T.	V.	—	—	—	B.fl.III,342	...	..	..	N.Z.			
H. geranifolia; F. v. M. in Transact. Vict. Inst. 126 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.III,342	M.fr.IV,181.						
H. medicaginoides; Turczaninow in Bull. Mosc. XXII, 27 (1849)	W.A.	S.A.	—	V.	—	—	—	B.fl.III,342	M.fr.IV,178.						
H. muriculata; Turczaninow in Bull. Mosc. 28 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,342							
H. callicarpa; Bunge in Lehm. pl. Preiss. I, 283 (1844)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,343	M.fr.IV,181.						
H. scutellifera; Bentham, Fl. Austr. III, 343 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,343							
H. hispidula; Bunge in Lehm. pl. Preiss. I, 283 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,343							
H. trachycarpa; F. v. M. in Schlecht. Linnaea XXV, 394 (1852)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,343	M.fr.IV,181.						
H. pilifera; Turczaninow in Bull. Mosc. XXII, 26 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,344							
H. capillaris; F. v. M., First Gen. Report 13 (1853)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,344	M.fr.IV,178.						
H. rugulosa; Turczaninow in Bull. Soc. Mosc. XXII, 27 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,344							
H. grammatocarpa; F. v. M., fragm. II, 128 (1861)	—	—	—	—	—	—	N.A.	B.fl.III,344	M.fr.II,128.						
H. diantha; De Candolle, prodr. IV, 63 (1830)	W.A.	S.A.	—	—	—	—	—	B.fl.III,345	M.fr.IV,179.						
H. lemnoides; Bentham, Fl. Austr. III, 345 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,345							
H. alata; R. Brown in annal. des sc. phys. VI, 75, t. 61 (1820)	W.A.	—	—	—	—	—	—	B.fl.III,345							
H. tetragonocarpa; Bunge in Lehm. pl. Preiss. I, 284 (1844)	W.A.	—	—	—	—	—	—	B.fl.III,345							
H. blepharocarpa; F. v. M. in Wing's S. Sc. Rec. III, 175 (1883)	W.A.	—	—	—	—	—	—	—							
H. glochidiata; Bentham, Fl. Austr. III, 346 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,346							
H. homalocarpa; F. v. M., fragm. II, 129 (1861)	W.A.	—	—	—	—	—	—	B.fl.III,346	M.fr.IV,181; VII,147.						
H. Asiatica; Linné, spec. plant. 234 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,346	M.fr.VII,147	...	As., Afr., Am., P., N.Z.				

DIDISCUS; De Candolle in Bot. Mag. t. 2857 (1828). (Dimetopia, Pritzelia, Huegelia, Cesatia, Hemicarpus.)

D. pusillus; F. v. M., fragm. IX, 47 (1875)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,348	M.fr.I,231.						
D. cyanopetalus; F. v. M., fragm. IX, 46 (1875)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,348	M.fr.I,231.						
D. eriocarpus; F. v. M., fragm. IX, 46 (1875)	W.A.	S.A.	—	—	—	—	—	B.fl.III,348	M.fr.I,231; IX,46.						
D. villosus; F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857)	—	—	—	—	—	—	N.A.	B.fl.III,349							
D. coeruleus; De Candolle in Hook. Bot. Mag. 2875 (1828)	W.A.	—	—	—	—	—	—	B.fl.III,349	M.fr.IX,47.						
D. pilosus; Bentham in Hueg. enum. 54 (1837)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,349							

D. glaucifolius; F. v. M. in Schlecht. Linnaea XXV, 395 (1852)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,350	M.fr.IX,47.
D. glandulosus; F. v. M. in Proc. Roy. Soc. Tasm. III, 238 (1857)	—	—	—	—	—	Q.	N.A.	B.fl.III,350	
D. incisus; Hooker, Bot. Magaz. 2875 (1828)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,350	
D. procumbens; F. v. M. in Proc. Roy. Soc. Tasm. III, 237 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,350	M.fr.IX,47.
D. hemicarpus; F. v. M. in Transact. Bot. Soc. Edinb. VII, 491 (1857) ...	—	—	—	—	—	—	N.A.	B.fl.III,351	M.fr.IX,47.
D. humilis; J. Hooker in icon. pl. t. 304 (1841)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,351	
TRACHYMENE; Rudge in Transact. Linn. Soc. X, 300 (1810). (Siebera, Fischera, Platysace, Platycarpidium.)									
T. compressa; Rudge in Transact. Linn. Soc. X, 300 (1810) implied	W.A.	—	—	—	—	—	—	B.fl.III,352	
T. juncea; Bunge in Lehm. pl. Preiss. I, 286 (1844)...	W.A.	—	—	—	—	—	—	B.fl.III,353	
T. haplosciadea; F. v. M., Cens. 62 (1882) from Bentham (1866)	W.A.	—	—	—	—	—	—	B.fl.III,353	
T. cirrosa; F. v. M., Cens. 62 (1882) from Bunge (1844)	W.A.	—	—	—	—	—	—	B.fl.III,354	M.fr.I,231.
T. heterophylla; F. v. M., First Gen. Rep. 13 (1853)	—	S.A.	—	V.	—	—	—	B.fl.III,354	
T. tenuissima; F. v. M., Cens. 62 (1866) from Bentham (1866)	W.A.	—	—	—	—	—	—	B.fl.III,354	
T. dissecta; F. v. M., Cens. 62 (1882) from Bentham (1866)	W.A.	—	—	—	—	—	—	B.fl.III,354	
T. commutata; Turczaninow in Bull. Mosc. XXII, 30 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,355	
T. effusa; Turczaninow in Bull. Mosc. XXII, 31 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,355	M.fr.I,231.
T. deflexa; Turczaninow in Bull. Mosc. XXII, 31 (1849)	W.A.	—	—	—	—	—	—	B.fl.III,355	
T. valida; F. v. M., Cens. 62 (1882)	—	—	—	—	—	Q.	—	B.fl.III,355	M.fr.I,232.
T. ericoides; Sieber in Spreng. syst. veg. cur. post. 116 (1827)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,356	
T. linearis; Sprengel, spec. umbellif. 7 (1818)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,356	
T. Billardieri; F. v. M., Cens. 62 (1882) from Bentham (1866)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,356	
T. Stephensonii; Turczaninow in Bull. de Mosc. XX, 170 (1847)	—	—	—	—	N.S.W.	—	—	B.fl.III,357	
XANTHOSIA; Rudge in Transact. Linn. Soc. X, 361 (1810). (Leucolaena, Schoenolaena, Pentapeltis.)									
X. juncea; Bentham, Fl. Austr. III, 359 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. tenuior; Bentham, Fl. Austr. III, 359 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. peltigera; Steudel, nomencl. bot. 791 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. hederifolia; Bentham, Fl. Austr. III, 359 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,359	
X. candida; Steudel, nomencl. bot. 790 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,259	
X. tridentata; De Candolle, prodr. IV, 75 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.III,260	
X. singuliflora; F. v. M., fragm. IV, 184 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,260	M.fr IV
X. pilosa; Rudge in Transact. Linn. Soc. X, 301, t. 22 (1811)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,260	
X. ciliata; Hooker, icon. pl. t. 726 (1848)	W.A.	—	—	—	—	—	—	B.fl.III,261	
X. pusilla; Bunge in Lehm. pl. Preiss. I, 291 (1844)	W.A.	S.A.	T.	V.	—	—	—	B.fl.III,261	
X. fruticulosa; Bentham, Fl. Austr. III, 361 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III 261	
X. Huegelii; Steudel, nomencl. bot. 791 (1841)	W.A.	—	—	—	—	—	—	B.fl.III,262	
X. dissecta; J. Hooker in icon. plant. t. 302 (1841)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,362	
X. peduncularis; Bentham, Fl. Austr. III, 362 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,362	
X. vestita; Bentham, Fl. Austr. III, 363 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,363	
X. Atkinsoniana; F. v. M., fragm. II, 127 (1861)	W.A.	—	—	V.	N.S.W.	—	—	B.fl.III,363	M.fr.II,127.
X. rotundifolia; De Candolle, prodr. IV, 75 (1830)	W.A.	—	—	—	—	—	—	B.fl.III,633	M.fr.II,128.
AZORELLA; Lamarck, Encycl. Méthod. I, 344, t. 189 (1783). (Fragosa, Pozoa, Microsciadium, Schizeilema, Oschatzia, Centella partly, Dichopetalum.)									
A. Muelleri; Bentham, Fl. Austr. III, 364 (1866)	—	—	—	V.	—	—	—	B.fl.III,364	
A. cuneifolia; F. v. M. in Transact. phil. Inst. Vict. I, 103 (1855)	—	—	—	V.	—	—	—	B.fl.III,365	
A. saxifraga; Bentham, Fl. Austr. III, 365 (1866)	—	—	T.	—	—	—	—	B.fl.III,365	
A. dichopetala; Bentham, Fl. Austr. III, 365 (1866)	—	—	T.	V.	—	—	—	B.fl.III,365	
HUANACA; Cavanilles, icon. VI, 18, t. 528 (1801). (Diplaspis, Pozoopsis.)									
H. hydrocotylea; Bentham & J. Hooker, gen. pl. I, 877 (1867)	—	—	T.	V.	—	—	—	B.fl.III,366	
H. cordifolia; Bentham & J. Hooker, gen. pl. I, 877 (1867)	—	—	T.	—	—	—	—	B.fl.III,366	

ACTINOTUS ; Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804). (Hemiphues, Eriocalia, Holotome.)

A. Helianthi ; Labillardière, Nov. Holl. pl. spec. I, 67, t. 92 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,367				
A. Schwarzii ; F. v. M. in proceed. R. Soc. of S.A. IX, 7 & 200 (1886)	—	S.A.	—	—	—	—	—	—				
A. leucocephalus ; Bentham in Hueg. enum. 56 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,368	M.fr.II,130.			
A. minor ; De Candolle, prodr. IV, 83 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.III,368	M.fr.VI,23.			
A. Gibbonsii ; F. v. M., fragm. VI, 23 (1867)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VI,23.			
A. omnifertilis ; Bentham, Fl. Austr. III, 368 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,368	M.fr.II,129.			
A. rhomboideus ; Bentham, Fl. Austr. III, 368 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,368				
A. bellidioides ; Bentham, Fl. Austr. III, 369 (1866)	—	—	T.	—	—	—	—	B.fl.III,369				
A. glomeratus ; Bentham, Fl. Austr. III, 369 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,369	M.fr.VI,23.			

ERYNGIUM ; Tournefort, inst. 327, t. 173 (1700) from Theophrastos and Dioscorides.

E. rostratum ; Cavanilles, icon. pl. VI, 35, t. 552 (1801)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,370				Am.
E. vesiculosum ; Labillardière, Nov. Holl. pl. spec. I, 73, t. 98 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III.370	M.fr.VII,148			Am., N.Z.
E. plantagineum ; F. v. M. in Pap. Roy. Soc. Tasm. III, 235 (1857)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,371	M.fr.VII,148.			
E. expansum ; F. v. M. in Pap. Roy. Soc. Tasm. III, 236 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,371	M.fr.VII,148.			

APIUM ; Tournefort, inst. 305, t. 160 (1700).

A. prostratum ; Labillardière, relat. du voy. I, 141 (1799)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,372	M.fr.VII,148			As., Afr., N.Z., P.
A. leptophyllum ; F. v. M. in Benth. Fl. Austr. III, 372 (1866)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,372	M.fr.IV,184; VII,148.			Am., P.

SIUM ; Tournefort, inst. 308, t. 162 (1700) from C. Bauhin (1623). (Helosciadium).

S. latifolium ; C. Bauhin, pinax 154 (1623)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,336				E., As.
S. erectum ; Hudson, fl. Angl. p. 103 (1762)	—	—	—	—	N.S.W.	—	—	B.fl.III,336				E., As., Am.

SESELI ; Rivinus in Rupp. Fl. Jen. 267 (1718).

S. Harveyanum ; F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.III,373				
S. algens ; F. v. M. in Transact. phil. Soc. Vict. I, 104 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.III,373				

CRANTZIA ; Nuttall, Gen. of North Amer. plants I, 177 (1818). (Non Scopoli, 1777, Crantziola, F. v. M.)

C. lineata ; Nuttall, Gen. of North Amer. pl. I, 178 (1818)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,374				Am., N.Z.

OENANTHE ; Tournefort, inst. 312, t. 166 (1700) from l'Obel (1576).

O. stolonifera ; Wallich, numerical list 585 (1825)	—	—	—	—	—	Q.	—	—	M.fr.V,182			As.

ACIPHYLLA ; R. & G. Forster, char. gen. 135, t. 68 (1776). (Anisotome, Calosciadium, Gingidium partly, Ligustici subgenus.)

A. simplicifolia ; F. v. M. in Benth. Fl. Austr. III, 375 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,375				
A. glacialis ; F. v. M. in Benth. Fl. Austr. III, 375 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,375				
A. procumbens ; F. v. M. in Benth. Fl. Austr. III, 375 (1866)	—	—	T.	—	—	—	—	B.fl.III,375				

DAUCUS ; Tournefort, inst. 307, t. 161 (1700) from l'Ecluse (1576).

D. brachiatus ; Sieber in De Candolle, prodr. IV, 214 (1830)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,376				Am., N.Z.

OREOMYRRHIS ; Endlicher, gen. plantar. 787 (1839). (Caldasia.)

O. andicola ; Endlicher, gen. plantar. 787 (1839)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,377	M.fr.VIII,185			Am., N.Z.
O. pulvinifica ; F. v. M., fragm. VIII, 185 (1874)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,185.			

SYNPETALEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney, 27 (1880).

ELAEAGNEAE.

R. Brown, prodr. I, 350 (1810).—Genus 1, Species 1.

ELAEAGNUS; Tournefort, coroll. 53, t. 459 (1703) from Camerarius (1586).

E. latifolia; Linné, spec. plant. 121 (1753) ... ... — — — — — Q. — B.fl.VI,39 M.fr.IV,146; IX,190 .. As.

OLACINEAE.

Mirbel in Bull. de la Soc. philomat. 377 (1813).—Genera 9, Species 15.

XIMENIA; Plumier, nova plant. Amer. genera 6, t. 21 (1703).

X. Americana; Linné, spec. plant. 1193 (1753) ... ... — — — — — Q. N.A. B.fl.I,391 M.fr.VI,3; IX,150 ... As., Afr., Am., P.

OLAX; Linné, amoen. acad. 387 (1747). (Spermaxyrum, Lopadocalyx.)

O. phyllanthi; R. Brown, prodr. 358 (1810) ... ... W.A. — — — — — — B.fl.I,392 M.fr.IX,151.
O. retusa; F. v. M. in Benth. Fl. Austr. I, 392 (1863) ... ... — — — — N.S.W. Q. — B.fl.I,392 M.fr.IX,151.
O. stricta; R. Brown, prodr. 358 (1810) ... ... — — — V. N.S.W. Q. — B.fl.I,392 M.fr.IX,151.
O. Benthamiana; Miquel in Lehm. pl. Preiss. I, 228 (1844) ... ... W.A. S.A. — — — — N.A. B.fl.I,393 M.fr.IX,151.
O. aphylla; R. Brown, prodr. 358 (1810) .. .. — — — — — Q. N.A. B.fl.I,393 M.fr.IX,151.

CANSJERA; A. L. de Jussieu, gen. pl. 448 (1789).

C. leptostachya; Bentham in Hook. Lond. Journ. of Bot. II, 231 (1843) ... — — — — — Q. N.A. B.fl.I,394 M.fr.VI,3 ... ... P.

OPILIA; Roxburgh, pl. Corom. II, 31, t. 158 (1799).

O. amentacea; Roxburgh, pl. Corom. II, 31, t. 158 (1799) .. .. — — — — — Q. N.A. B.fl.I,394 M.fr.VI,4;IX,150 ... As., Afr.

PHLEBOCALYMMA; Griffith in Benth. & J. Hook. gen. pl. I, 353 (1862). (Phlebocalymna.)

P. lobospora; F. v. M., fragm. IX, 151 (1875) ... ... — — — — — Q. — — M.fr.IX,151.

PENNANTIA; R. & G. Forster, char. gen. 133, t. 67 (1776).

P. Cunninghamii; Miers in Ann. of nat. hist. sec. ser. IX, 491 (1852) ... ... — — — — N.S.W. Q. — B.fl.I,395 M.fr.VI,4; IX,150.
P. Endlicheri; Reisseck in Schlecht. Linnaea XVI, 341, t. 13 (1843) ... ... — — — — N.S.W. — — — M.fr.IX,150.

APODYTES; E. Meyer in Hook. Journ. of Bot. III, 155 (1841).

A. brachystylis; F. v. M., fragm. IX, 149 (1875) ... ... — — — — — Q. — — M.fr.IX,149.

VILLARESIA; Ruiz & Pavon, flor. Peruv. et Chil. prodr. 35 (1794).

V. Moorei; F. v. in Benth. Fl. Austr. I, 396 (1863)... ... ... — — — — N.S.W. — — B.fl.I,395
V. Smythii; F. v. M., fragm. V, 156 (1866) ... ... — — — — — Q. — — M.fr.IX,150.

GOMPHANDRA; Wallich, Catal. n. 3718 & 7204 (1832).

G. Australiana; F. v. M., fragm. VI, 3 (1877) ... ... — — — — — Q. — — M.fr.VI,3&253; IX,150.

BALANOPHOREAE.

L. Cl. Richard in Mém. du Mus. VIII, 404 & 429 (1822).—Genus 1, Species 1.

BALANOPHORA; R. & G. Forster, char. gen. 99, t. 50 (1776).

B. fungosa; R. & G. Forster, char. gen. 99, t. 50 (1776) ... ... — — — — — Q. — B.fl.VI,232 ... . P.

SANTALACEAE.

R. Brown, prodr. I, 350 (1810).—Genera 7, Specief 43.

EXOCARPOS; Labillardière, Relat. du voy. à la rech. de La Pérouse I, 155 (1789). (Exocarpus.)

E. latifolia; R. Brown, prodr. 356 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,228	M.fr.VIII,10	...	...	As., P.
E. odorata; A. de Candolle, prodr. XIV, 689 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,228	M.fr.VIII,10.			
E. cupressiformis; Labillardière, voy. I, 155, t. 14 (1798)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,229				
E. spartea; R. Brown, prodr. 356 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,229	M.fr.VIII,10; IX,20.			
E. aphylla; R. Brown, prodr. 357 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,230	M.fr.VIII,10; IX,169; XI,20.			
E. homaloclada; C. Moore & F. v. M., fragm. VIII. 9 (1871)	—	—	—	—	N.S.W.	—	—	B.fl.VI,230	M.fr.VIII,9.			
E. phyllanthoides; Endlicher, prodr. pl. ins. Norf. 46 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.			
E. stricta; R. Brown, prodr. 357 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,230	M.fr.IX,169.			
E. humifusa; R. Brown, prodr. 356 (1810)	—	—	T.	—	—	—	—	B.fl.VI,231				
E. nana; J. Hooker in Lond. Journ. of Bot. VI, 281 (1847)	—	—	T.	V.	—	—	—	B.fl.VI,230				

ANTHOBOLUS; R. Brown, prodr. 357 (1810).

A. filifolius; R. Brown, prodr. 357 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,226	M,fr.I,213; IX,3.			
A. triqueter; R. Brown, prodr. 357 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,226				
A. foveolatus; F. v. M., fragm. I, 212 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,226	M.fr.II,181; IX,3; XI,20.			
A. leptomerioides; F. v. M., fragm. I, 21 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,227	M.fr.IX,3.			
A. exocarpoides; F. v. M., fragm. IX, 3 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,3.			

OMPHACOMERIA; Endlicher, gen. pl. 326 (1838).

O. acerba; A. de Candolle, prodr. XIV, 681 (1856)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,224					

LEPTOMERIA; R. Brown, prodr. 353 (1810).

L. spinosa; A. de Candolle, prodr. XIV, 678 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,220	M.fr.VIII,10.				
L. Preissiana; A. de Candolle, prodr. XIV, 678 (1856)	W.A.	—	—	—	—	—	—	B.fl.VI,221	M.fr.VIII,10.				
L. pauciflora; R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,221	M.fr.VIII,10.				
L. scrobiculata; R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,221	M.fr.VIII,10.				
L. acida; R. Brown, prodr. 353 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,222	M.fr.VIII,10.				
L. Billardieri; R. Brown, prodr. 354 (1810)	—	—	T.	—	N.S.W.	Q.	—	B.fl.VI,222	M.fr.VIII,10.				
L. aphylla; R. Brown, prodr. 354 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,222	M.fr.VIII,10.				
L. glomerata; F. v. M. in J. Hook. Fl. Tasm. II, 370 (1860)	—	—	T.	—	—	—	—	B.fl.VI,223	M.fr.VIII,10.				
L. squarrulosa; R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,223	M.fr.VIII,11.				
L. Cunninghamii; Miquel in Lehm. pl. Preiss. I, 611 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,223	M.fr.VIII,10.				
L. empetriformis; Miquel in Lehm. pl. Preiss. I, 610 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,224					
L. axillaris; R. Brown, prodr. 354 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,224					
L. laxa; Miquel in Lehm. pl. Preiss. I, 612 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,224					
L. obovata; Miquel in Lehm. pl. Preiss. I, 613 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,224					

CHORETRUM; R. Brown, prodr. 354 (1810).

C. glomeratum; R. Brown, prodr. 354 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,218	M.fr.I,21.				
C. chrysanthum; F. v. M. in Transact. phil. Soc. Vict. I, 23 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,218	M.fr.I,21; VIII,11; XI,138.				
C. spicatum; F. v. M., fragm. I, 21 (1858)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,218	M.fr.I,21; VIII,11.				
C. lateriflorum; R. Brown, prodr. 354 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI.219	M.fr.I,21.				
C. Candollei; F. v. M. in Bentham's Fl. Austr. VI, 219 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,219					

SANTALUM; Linné, gen. ed. sec. 165 (1742) from C. Bauhin (1623). (Fusanus, Eucarya.)

S. lanceolatum; R. Brown, prodr. 356 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,214	M.fr.I,85;VIII,11;XI,3.				
S. ovatum; R. Brown, prodr. 355 (1810)	—	—	—	—	—	—	N.A.	B.fl.VI,214					
S. obtusifolium; R. Brown, prodr. 356 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,215	M.fr.VIII,11.				

S. acuminatum ; A. de Candolle, prodr. XIV, 684 (1856)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,216	M.fr.I,85; IX,169; XI,20.							
S. persicarium ; F. v. M. in Transact. Vict. Inst. 41 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,216	M.fr.I,86; IX,3; XI,20.							
S. cygnorum ; Miquel in Lehm. pl. Preiss. I, 615 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,217	M.fr.VIII,11.							
S. crassifolium ; A. de Candolle, prodr. XIV, 685 (1856)	—	—	—	—	N.S.W.	—	—	B.fl.VI,217								
THESIUM ; Linné, gen. pl. 60 (1737).																
T. Australe ; R. Brown, prodr. 353 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,212	M.fr.IX,3	As.						

LORANTHACEAE.

A. L. de Jussieu in Ann. du Mus. XII, 292 (1808).—Genera 5, Species 27.

VISCUM ; Tournefort, inst. 609, t. 380 (1700), from Camerarius (1586), from W. Turner (1538).																
V. orientale ; Willdenow, spec. plant. IV, 737 (1805)	—	—	—	—	—	Q.	—	B.fl.III,396	... ...	As., P.						
V. angulatum ; Heyne in De Candolle, prodr. IV, 283 (1830)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,396	... ...	As.						
V. articulatum ; N. L. Burmann, Fl. Ind. 311 (1768)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,396	... ...	As., P.						
NOTOTHIXOS ; Oliver in Journ. Linn. Soc. VII, 104 (1865).																
N. incanus ; Oliver in Journ. Linn. Soc. VII, 104 (1865)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,397	M.fr.II,109,176; IV,173.							
LORANTHUS ; Linné, syst. nat. ed. sec. 22 (1740).																
L. celastroides ; Sieber in Roem. and Schult. syst. veg. VII, 163 (1829)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,389	M.fr.VI,252.							
L. Bidwillii ; Bentham, Fl. Austr. III, 390 (1866)	—	—	—	—	—	Q.	—	B.fl.III,390								
L. myrtifolius ; Cunningham in Benth. Fl. Austr. III, 390 (1866)...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,390								
L. longiflorus ; Desrousseaux in Lam. Encycl. méth. IV, 598 (1796)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,390	... ...	As.						
L. angustifolius ; R. Brown in Benth. Fl. Austr. III, 390 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,390								
L. dictyophlebus ; F. v. M., Rep. Burdek. Exped. 14 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,391								
L. maytenifolius ; A. Gray, Bot. Wilk. Expl. Exped. I, 739, t. 99 (1854) ...	—	—	—	—	N.S.W.	—	—	B.fl.III,393								
L. alyxifolius ; F. v. M. in Bentham Fl. Austr. III, 391 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,391								
L. odontocalyx ; F. v. M. in Bentham Fl. Austr. III, 391 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,391								
L. linearifolius ; Hooker in Mitchell, Trop. Austr. 102 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,391								
L. Murrayi ; Tate in Transact. R. Soc. of S.A. VI, 109 (1883)	—	S.A.	—	—	—	—	—	—								
L. Exocarpi ; Behr in Schlecht. Linnaea XX, 624 (1847)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,392								
L. acacioides ; Cunningham in Bentham, Fl. Austr. III, 392 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,392								
L. signatus ; F. v. M. in Bentham, Fl. Austr. III, 392 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,392	... ...	As.						
L. sanguineus ; F. v. M., fragm. I, 177 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.III,393	M.fr.IV,171.							
L. bifurcatus ; Bentham, Fl. Austr. III, 393 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,393								
L. linophyllus ; Fenzl in Huegel, enum. 56 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,393								
L. gibberulosus ; Tate in Transact. R. S. of S.A. VIII, 71 Oct. (1885)	—	S.A.	—	—	—	—	—	—								
L. pendulus ; Sieber in Sprengel, cur. poster. 139 (1827)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,394								
L. Quandang ; Lindley in Mitch. Three Exped. II, 69 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,395								
L. grandibracteus ; F. v. M., Rep. Burdek. Exped. 14 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,395								
ATKINSONIA ; F. v. M., fragm. V, 34 (1865). (Gaiadendron partly.)																
A. ligustrina ; F. v. M., fragm. V, 34 (1865)	—	—	—	—	N.S.W.	—	—	B.fl.III,388	M.fr.II,130; VI,252.							
NUYTSIA ; R. Brown in Journ. of the geogr. Soc. I, 17 (1831).																
N. floribunda ; R. Brown in Journ. geogr. Soc. I, 17 (1831)	W.A.	—	—	—	—	—	—	B.fl.III,387	M.fr.VI,252.							

PROTEACEAE.

A. L. de Jussieu, gen. plant. 78 (1789).—Genera 33, Species 597.

PETROPHILA ; R. Brown in Transact. Linn. Soc. X, 67 (1809).																
P. teretifolia ; R. Brown in Transact. Linn. Soc. X, 68 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,321	M.fr.VI,245.							
P. longifolia ; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,322	M.fr.VI,255.							

P. media; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,322	M.fr.VI,255.
P. acicularis; R. Brown in Transact. Linn. Soc. X, 69 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,323	M.fr.VI,245.
P. megalostegia; F. v. M., fragm. X, 61 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,61.
P. linearis; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,323	M.fr.VI,243.
P. anceps; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	—	M.fr.VI,243; X,47.
P. heterophylla; Lindley, Bot. Regist. XXV, App. 35 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,324	M.fr.VI,244.
P. biloba; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,324	M.fr.VI,255.
P. propinqua; R. Brown, prot. nov. 7 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,325	
P. squamata; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,325	M.fr.VI,244.
P. colorata; Meissner in Lehm. pl. Preiss. II, 246 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,326	
P. striata; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,326	M.fr.VI,244.
P. divaricata; R. Brown, prot. nov. 7 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,326	M.fr.VI,344.
P. Serruriae; R. Brown, prot. nov. 6 (1830)	W.A.	—		—	—	—	—	B.fl.V,327	M.fr.VI,243.
P. inconspicua; Meissner in Hook. Kew Misc. VII, 68 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,327	M.fr.VI,243.
P. trifida; R. Brown in Transact. Linn. Soc. X, 70 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,328	M.fr.VI,244.
P. carduacea; Meissner in Hook. Kew Misc. IV, 182 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,328	M.fr.VI,244.
P. Shuttleworthiana; Meissner in Lehm. pl. Preiss. II, 246 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,329	M.fr.VI,245.
P. macrostachya; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,329	M.fr.VI,245.
P. diversifolia; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,329	M.fr.VI,244.
P. biternata; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,330	M.fr.VI,245.
P. plumosa; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,330	M.fr.VI,244.
P. ericifolia; R. Brown, prot. nov. 5 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,331	M.fr.VI,243.
P. chrysantha; Meissner in Hook. Kew Misc. VII, 68 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,331	M.fr.VI,243.
P. pedunculata; R. Brown in Transact. Linn. Soc. X, 70 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,332	M.fr.VI,242.
P. pulchella; R. Brown in Transact. Linn. Soc. X, 69 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,332	M.fr.VI,242.
P. sessilis; Sieber in Roem. and Schult. mant. 262 (1822)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,332	
P. fastigiata; R. Brown in Transact. Linn. Soc. X, 70 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,333	M.fr.VI,245.
P. seminuda; Lindley, Bot. Regist. XXV, App. 34 (1839)...	W.A.	—	—	—	—	—	—	B.fl.V,333	M.fr.VI,243.
P. circinata; Kippist in Hook. Kew Misc. VII, 67 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,333	M.fr.VI,244.
P. Drummondii; Meissner in Lehm. pl. Preiss. I, 496 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,334	M.fr.VI,243.
P. crispata; R. Brown, prot. nov. 6 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,334	
P. rigida; R. Brown in Transact. Linn. Soc. X, 69 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,334	M.fr.VI,245.
P. multisecta; F. v. M., fragm. VI, 242 (1868)	—	S.A.	—	—	—	—	—	B.fl.V.335	M.fr.VI,242.
P. conifera; Meissner in Hook. Kew Misc. VII, 67 (1855)...	W.A.	—	—	—	—	—	—	B.fl.V,335	M.fr.VI,245.
P. semifurcata; F. v. M. in Bentham Fl. Austr. V, 335 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,335	M.fr.X,47.
ISOPOGON; R. Brown in Transact. Linn. Soc. IX, 70 (1809).									
I. latifolius; R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,338	M.fr.VI,237.
I. cuneatus; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,339	M.fr.VI,238.
I. linearis; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,339	M.fr.VI,236.
I. polycephalus; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—		—	B.fl.V,339	M.fr.VI,236.
I. attenuatus; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,340	M.fr.VI,237.
I. sphaerocephalus; Lindley, Bot. Regist. XXV, App. 34 (1839)...	W.A.	—	—	—	—	—	—	B.fl.V,340	M.fr.VI,236.
I. uncinatus; R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,341	M.fr.VI,237.
I. buxifolius; R. Brown in Transact. Linn. Soc. X, 74 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,341	M.fr.VI,237.
I. axillaris; R. Brown in Transact. Linn. Soc. X, 74 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,351	M.fr.VI,237.
I. tridens; F. v. M., fragm. VI, 239 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,342	M.fr.VI,239.
I. Baxteri; R. Brown, prot. nov. 9 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,342	M.fr.VI,240.
I. roseus; Lindley, Bot. Regist. Misc. n. 37 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,343	M.fr.VI,240.
I. adenanthoides; Meissner in Hook. Kew Misc. VII, 69 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,343	M.fr.VI,241.
I. trilobus; R. Brown in Transact. Linn. Soc. X, 72 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,343	M.fr.VI,239.

I. tripartitus ; R. Brown, prot. nov. 8 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,344	
I. longifolius ; R. Brown in Transact. Linn. Soc. X, 73 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,344	M.fr.VI,237.
I. Drummondii ; Bentham, Fl. Austr. V, 344 (1870)...	W.A.	—	—	—	—	—	—	B.fl.V,344	M.fr.VI,241.
I. heterophyllus ; Meissner in Lehm. pl. Preiss. I, 504 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. villosus ; Meissner in De Candolle, prodr. XIV, 277 (1856)	W.A.	—	—	—	—	—	—	B.fl.V,345	M.fr.VI,241.
I. teretifolius ; R. Brown in Transact. Linn. Soc. X, 71 (1809)	W.A.	—	—	—	—	—	—	B.fl.V.345	M.fr.VI,241.
I. anethifolius ; R. Brown in J. Knight, proteac. 94 (1809)	W.A.	—	—	—	N.S.W.	—	—	B.fl.V,346	M.fr.VI,238.
I. petiolaris ; Cunningham in R. Brown, prot. nov. 8 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,346	
I. anemonifolius ; R. Brown in J. Knight, prot. 93 (1809)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,347	M.fr.VI,238; X,90.
I. ceratophyllus ; R. Brown in Transact. Linn. Soc. X, 72 (1809)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,347	M.fr.VI,238.
I. asper ; R. Brown, prot. nov. 8 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,348	M.fr.VI,240.
I. crithmifolius ; F. v. M., fragm. VI, 239 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,348	M.fr.VI,239.
I. formosus ; R. Brown in Transact. Linn. Soc. X, 72 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,349	M.fr.VI,240.
I. divergens ; R. Brown, prot. nov. 7 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,349	M.fr.VI,241.
I. scabriusculus ; Meissner in Hook. Kew Misc. IV, 182 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,349	M.fr.VI,240.
ADENANTHOS ; Labillardière, Nov. Holl. pl. specim. I, 28, t. 36 (1804).									
A. Detmoldi ; F. v. M., fragm. VIII, 149 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,149.
A. barbigerus ; Lindley, Bot. Regist. XXV, App. 36 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,351	M.fr.VI,205; VIII,149.
A. obovatus ; Labillardière, Nov. Holl. pl. spec. I, 29, t. 37 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,352	
A. cuneatus ; Labillardière, Nov. Holl. pl. spec. I, 28, t. 36 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,352	
A. Cunninghami ; Meissner in Lehm. pl. Preiss. I, 513 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,352	
A. pungens ; Meissner in Lehm. pl. Preiss. I, 515 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,353	
A. venosus ; Meissner in Hook. Kew Misc. IV, 183 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,353	
A. Dobsoni ; F. v. M., fragm. VI, 204 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,353	M.fr.VI,204.
A. Forrestii ; F. v. M. in Wing's S. Sc. Rec. II, 230 (1882)	W.A.	—	—	—	—	—	—	—	
A. linearis ; Meissner in Hook. Kew Misc. IV, 183 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,354	
A. sericeus ; Labillardière, Nov. Holl. pl. specim. I, 29, t. 38 (1804)	W.A.	S.A.	—	—	—	—	—	B.fl.V,354	
A. Meissneri ; Lehmann, pl. Preiss. I, 515 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,354	
A. filifolius ; Bentham, Fl. Austr. V, 355 (1873)	W.A.	—	—	—	—	—	—	B.fl.V,355	
A. terminalis ; R. Brown in Transact. Linn. Soc. X, 152 (1809)	—	S.A.	—	V.	—	—	—	B.fl.V,355	M.fr.I,158.
A. flavidiflorus ; F. v. M., fragm. I, 157 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,355	M.fr.I,157.
A. apiculatus ; R. Brown, prot. nov. 9 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,356	
SIMSIA ; R. Brown in Transact. Linn. Soc. X, 152 (1809). (Stirlingia.)									
S. simplex ; F. v. M., Cens. of Austr. pl. 66 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.V,357	M.fr.VI,248.
S. abrotanoides ; F. v. M., Cens. of Austr. pl. 66 (1882) from Meissner (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,357	
S. teretifolia ; F. v. M., Cens. of Austr. pl. 66 (1882) from Meissner (1845) ...	W.A.	—	—	—	—	—	—	B.fl.V,357	
S. tenuifolia ; R. Brown in Transact. Linn. Soc. X, 152 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,358	
S. latifolia ; R. Brown, prot. nov. 9 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,358	M.fr.VI,248.
SYNAPHEA ; R. Brown in Transact. Linn. Soc. X, 155 (1809).									
S. polymorpha ; R. Brown in Transact. Linn. Soc. X, 156 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,360	
S. dilatata ; R. Brown in Transact. Linn. Soc. X, 156 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,360	
S. favosa ; R. Brown in Transact. Linn. Soc. X, 156 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,360	
S. Preissii ; Meissner in Lehm. pl. Preiss. I, 529 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,361	
S. acutiloba ; Meissner in Lehm. pl. Preiss. I, 528 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,361	
S. petiolaris ; R. Brown in Transact. Linn. Soc. X, 156 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,361	
S. decorticans ; Lindley, Bot. Regist. XXV, App. 32 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,362	
S. pinuata ; Lindley, Bot. Regist. XXV, App. 32 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,362	
CONOSPERMUM ; Smith in Transact. Linn. Soc. IV, 213 (1798).									
C. capitatum ; R. Brown in Transact. Linn. Soc. X, 155 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,365	
C. petiolare ; R. Brown, prot. nov. 11 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,365	

C. teretifolium; R. Brown in Transact. Linn. Soc. X, 155 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,365	M.fr.VI,224.
C. flexuosum; R. Brown, prot. nov. 11 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,365	M.fr.VI,224.
C. acerosum; Lindley, Bot. Regist. XXV, App. 30 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,366	
C. amoenum; Meissner in Lehm. pl. Preiss. I, 522 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,366	M.fr.I,157; VI,224.
C. nervosum; Meissner in Hook. Kew Misc. VII, 71 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,367	M.fr.VI,224.
C. diffusum; Bentham, Fl. Austr. V, 367 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,367	
C. glumaceum; Lindley, Bot. Regist. XXV, App. 30 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,367	
C. ephedroides; Kippist in Hook. Kew Misc. VII, 70 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,368	
C. Toddii; F. v. M., fragm. X, 20 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,20,90.
C. polycephalum; Meissner in Lehm. pl. Preiss. II, 249 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,368	
C. coeruleum; R. Brown in Transact. Linn. Soc. X, 154 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,368	M.fr.VI,224.
C. debile; Kippist in Hook. Kew Misc. VII, 70 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. scaposum; Bentham, Fl. Austr. V, 369 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. Huegelii; R. Brown in Endlicher, nov. stirp. dec. 58 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. densiflorum; Lindley, Bot. Regist. XXV, App. 32 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,369	
C. Brownii; Meissner in Lehm. pl. Preiss. II, 249 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,370	
C. longifolium; Smith, Exot. Bot. II, 45, t. 82 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.V,370	
C. tenuifolium; R. Brown in Transact. Linn. Soc. X, 154 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,370	M.fr.VI,223.
C. Mitchellii; Meissner in De Candolle, prodr. XIV, 320 (1856)	—	S.A.	—	V.	—	—	—	B.fl.V,371	
C. sphacelatum; Hooker in Mitchell, Trop. Austr. 342 (1848)	—	—	—	—	—	Q.	N.A.	B.fl.V,371	M.fr.VI,223.
C. patens; Schlechtendal, Linnaea XX, 587 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,371	M.fr.VI,223.
C. taxifolium; Smith in Rees's Cycl. IX (1808)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,372	M.fr.VI,224.
C. ericifolium; Smith in Rees's Cycl. IX (1808)	—	—	—	—	N.S.W.	—	—	B.fl.V,372	
C. ellipticum; Smith in Rees's Cycl. IX (1808)	—	—	—	—	N.S.W.	—	—	B.fl.V,372	
C. distichum; R. Brown in Transact. Linn. Soc. X, 155 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,373	M.fr.VI,224.
C. florihundum; Bentham, Fl. Austr. V, 373 (1870)...	W.A.	—	—	—	—	—	—	B.fl.V,373	
C. incurvum; Lindley, Bot. Regist. XXV, App. 30 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,373	
C. brachyphyllum; Lindley, Bot. Regist. XXV, App. 31 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,374	
C. stoechadis; Endlicher in Ann. Wien. Mus. II, 208 (1838)	W.A.	—	—	—	—	—	—	B.fl.V,374	M.fr.XI,20.
C. triplinervium; R. Brown, prot. nov. 11 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,375	
C. bracteosum; Meissner in Lehm. pl. Preiss. I, 518 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,375	M.fr.VI,224.
C. crassinervium; Meissner in Hook. Kew Misc. IV, 184 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,375	
FRANKLANDIA; R. Brown in Transact. Linn. Soc. X, 157 (1809).									
F. fucifolia; R. Brown in Transact. Linn. Soc. X, 157 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,376	M.fr.VI,223; VII,133.
F. triaristata; Bentham, Fl. Austr. V, 377 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,377	
SYMPHYONEMA; R. Brown in Transact. Linn. Soc. X, 157 (1809). (Symphionema.)									
S. montanum; R. Brown in Transact. Linn. Soc. X, 158 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,377	M.fr.VI,223.
S. paludosum; R. Brown in Transact. Linn. Soc. X, 158 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,378	M.fr.VI,223.
BELLENDENA; R. Brown in Transact. Linn. Soc. X, 156 (1809).									
B. montana; R. Brown in Transact. Linn. Soc. X, 166 (1809)	—	—	T.	—	—	—	—	B.fl.V,378	
AGASTACHYS; R. Brown in Transact. Linn. Soc. X, 158 (1809).									
A. odorata; R. Brown in Transact. Linn. Soc. X, 158 (1809)	—	—	T.	—	—	—	—	B.fl.V,379	M.fr.X,90.
CENARRHENES; Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804).									
C. nitida; Labillardière, Nov. Holl. pl. spec. I, 36, t. 50 (1804)	—	—	T.	—	—	—	—	B.fl.V,380	
PERSOONIA; Smith in Transact. Linn. Soc. IV, 215 (1798). (Linkia 1797, Pentadactylon.)									
P. hakeiformis; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,383	M.fr.VI,223.
P. teretifolia; R. Brown in Transact. Linn. Soc. X, 160 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,383	M.fr.VI,223.
P. saccata; R. Brown, prot. nov. 12 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,384	M.fr.VI,223.

P. Saundersiana; Kippist in Hook. Kew Misc. VII, 72 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,384	
P. diadena; F. v. M., fragm. X, 46 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,46.
P. comata; Meissner in Hooker's Kew Misc. VII, 71 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,385	
P. brachystylis; F. v. M., fragm. VI, 221 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,385	
P. falcata; R. Brown in Transact. Linn. Soc. X, 162 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,385	M.fr.VI,222.
P. trinervis; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,386	
P. tortifolia; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,386	
P. angustiflora; Bentham, Fl. Austr. V, 386 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,386	
P. rudis; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,387	
P. microcarpa; R. Brown in Transact. Linn. Soc. X, 160 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,387	M.fr.VI,222.
P. sulcata; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,387	
P. acicularius; F. v. M., fragm. VI, 220 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,388	M.fr.VI,220.
P. scabrella; Meissner in Hooker's Kew Misc. VII, 72 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,388	
P. dillwynioides; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,388	M.fr.VI,222.
P. striata; R. Brown, prot. nov. 13 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,388	M.fr.VI,223.
P. quinquenervis; Hooker, icon. plant. t. 425 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,389	
P. rufiflora; Meissner in Hooker's Kew Misc. VII, 72 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,389	
P. scabra; R. Brown in Transact. Linn. Soc. X, 162 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,389	
P. graminea; R. Brown in Transact. Linn. Soc. X, 164 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,390	M.fr.VI,222.
P. longifolia; R. Brown in Transact. Linn. Soc. X, 164 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,390	M.fr.VI,222.
P. articulata; R. Brown in Transact. Linn. Soc. X, 164 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,390	
P. elliptica; R. Brown in Transact. Linn. Soc. X, 164 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,391	M.fr.VI,222.
P. ferruginea; Smith, Exot. Bot. II, 47, t. 83 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.V,391	
P. media; R. Brown, prot. nov. 16 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,391	
P. cornifolia; Cunningham in R. Brown, prot. nov. 16 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,392	
P. marginata; R. Brown, prot. nov. 16 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,492	
P. sericea; Cunningham in R. Brown, prot. nov. 14 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,393	
P. Mitchellii; Meissner in Hook. Kew Misc. VII, 73 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,393	
P. fastigiata; R. Brown, prot. nov. 13 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,393	
P. hirsuta; Persoon, synops. plant. I, 118 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.V,394	M.fr.VI,222.
P. Chamaepitys; Cunningham in Field, N.S.W. 329 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.V,394	M.fr.VI,223.
P. arborea; F. v. M., gen. rep. 18 (1860)	—	—	—	V.	—	—	—	B.fl.V,394	M.fr.V,37; VI,221.
P. salicina; Persoon, synops. plant. I, 118 (1805)	—	—	—	V.	N.S.W.	—	—	B.fl.V,395	
P. prostrata; R. Brown in Transact. Linn. Soc. X, 163 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,395	
P. lanceolata; Andrews, Bot. Reposit. t. 74 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,395	M.fr.VI,222.
P. confertiflora; Bentham, Fl. Austr. V, 396 (1870)	—	—	—	V.	—	—	—	B.fl.V,396	
P. lucida; R. Brown in Transact. Linn. Soc. X, 161 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,396	
P. linearis; Andrews, Bot. Reposit. t. 77 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,397	M.fr.VI,222.
P. pinifolia; R. Brown in Transact. Linn. Soc. X, 160 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,397	M.fr.VI,222; X,47.
P. Caleyi; R. Brown, prot. nov. 13 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,398	
P. ledifolia; Cunningham in De Candolle, prodr. XIV, 339 (1856)	—	—	—	—	N.S.W.	—	—	B.fl.V,398	
P. revoluta; Sieber in Roem. & Schult. mant. 272 (1822)	—	—	—	V.	N.S.W.	—	—	B.fl.V,398	M.fr.VI,221.
P. Gunnii; J. Hooker in Lond. Journ. Bot. VI, 283 (1843)	—	—	T.	—	—	—	—	B.fl.V,399	
P. mollis; R. Brown in Transact. Linn. Soc. X, 161 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,399	M.fr.VI,221.
P. rigida; R. Brown, prot. nov. 14 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.V,399	M.fr.VI,221; VIII,149; X,47.
P. curvifolia; R. Brown, prot. nov. 13 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,400	
P. oblongata; Cunningham in R. Brown, prot. nov. 14 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,400	
P. Cunninghamii; R. Brown, prot. nov. 15, (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,400	
P. myrtilloides; Sieber in Roem. & Schult. mant. 272 (1822)	—	—	—	V.	N.S.W.	—	—	B.fl.V,401	M.fr.VI,221.
P. oxycoccoides; Sieber in Roem. & Schult. mant. 270 (1822)	—	—	—	V.	N.S.W.	—	—	B.fl.V,401	M.fr.VI,221.

P. nutans ; R. Brown in Transact. Linn. Soc. X, 162 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,401	M.fr.VI,222.
P. angulata ; R. Brown, prot. nov. 14 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,402	
P. virgata ; R. Brown in Transact. Linn. Soc. X, 161 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,402	
P. Chamaepeuce ; Lhotsky in De Candolle, prodr. XIV, 336 (1856)	—	—	—	V.	N.S.W.	—	—	B.fl.V,402	M.fr.VIII,149; X,47.
P. juniperina ; Labillardière, Nov. Holl. pl. spec. I, 33, t. 45 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,403	
P. tenuifolia ; R. Brown, prot. nov. 12 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,403	
P. acerosa ; Sieber in Roem. & Schult. mant. 269 (1822)	—	—	—	—	N.S.W.	—	—	B.fl.V,403	M.fr.VI,222.

MACADAMIA ; F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857). (Panopsis partly, Andriapetalum partly.)

M. ternifolia ; F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,406	M.fr.II,91; III,38; VI,191,224; VII,59.

HELICIA ; Loureiro, Fl. Cochinch. I, 83 (1790).

H. Youngiana ; C. Moore & F. v. M., fragm. IV, 84 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,406	M.fr.V,186.
H. praealta ; F. v. M., fragm. III, 37 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,404	M.fr.IV,174;VI,224.
H. Australasica ; F. v. M. in Hook. Kew Misc. IX, 27 (1857)	—	—	—	—	—	—	N.A.	B.fl.V,405	M.fr.II,91.
H. glabriflora ; F. v. M., fragm. II, 91 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,405	M.fr.III,38; V,38,186.
H. ferruginea ; F. v. M., fragm. III, 37 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,405	M.fr.IV,174; V,186; VI,191,224.

ROUPALA ; Aublet, Hist. des pl. de la Guian. I, 83, t. 32 (1775). (Rhopala, Bleasdalea, Adenostephanus partly, Kermadecia.)

R. Bleasdalei ; F. v. M. in Journ. Roy. Soc. N.S.W. XV, 212 (1881) implied	—	—	—	—	—	Q.	—	B.fl.V,417	M.fr.V,90.

HICKSBEACHIA ; F. v. M. in Wing's South. Science Record III, 32 (1883).

H. pinnatifolia ; F. v. M. in S. Sc. Rec. III, 33 (1883)	—	—	—	—	N.S.W.	Q.	—	—	

XYLOMELUM ; Smith in Transact. Linn. Soc. IV, 214 (1798).

X. pyriforme ; Smith in J. Knight, prot. 105 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,408	M.fr.VI,220.
X. occidentale ; R. Brown, prot. nov. 31 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,408	
X. salicinum ; Cunningham in R. Brown, prot. 31 (1830)	—	—	—	—	—	Q.	—	B.fl.V,408	M.fr.VI,220.
X. Scottianum ; F. v. M., fragm. V, 174 (1866)	—	—	—	—	—	Q.	—	—	M.fr.IV,107; V,215.
X. angustifolium ; Kippist in Hook. Kew Misc. IV, 209 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,409	M.fr.VI,220.

LAMBERTIA ; Smith in Transact. Linn. Soc. IV, 214 (1798).

L. uniflora ; R. Brown in Transact. Linn. Soc. X, 188 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,413	M.fr.VI,248.
L. rariflora ; Meissner in Lehm. pl. Preiss. II, 263 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,414	
L. inermis ; R. Brown in Transact. Linn. Soc. X, 188 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,414	M.fr.VI,248.
L. ericifolia ; R. Brown, prot. nov. 30 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,414	M.fr.VI,248.
L. multiflora ; Lindley, Bot. Regist. XXV, App. 32 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,415	
L. formosa ; Smith in Transact. Linn. Soc. IV, 214, t. 20 (1798)	—	—	—	—	N.S.W.	—	—	B.fl.V,415	M.fr.VII,133.
L. echinata ; R. Brown in Transact. Linn. Soc. X, 189 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,416	M.fr.VI,248.
L. ilicifolia ; Hooker, icon. plant. t. 553 (1843)	W.A.	—	—	—	—	—	—	B.fl.V,416	

ORITES ; R. Brown in Transact. Linn. Soc. X, 189 (1809). (Oritina.)

O. excelsa ; R. Brown, prot. nov. 32 (1832)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,411	M.fr.V,153; VI,223.
O. diversifolia ; R. Brown in Transact. Linn. Soc. X, 190 (1809)	—	—	T.	—	—	—	—	B.fl.V,411	M.fr.V,153.
O. Milligani ; Meissner in Hook. Kew Misc. IV, 209 (1852)	—	—	T.	—	—	—	—	B.fl.V,411	
O. lancifolia ; F. v. M. in Transact. phil. Soc. Vict. I, 108 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.V,412	M.fr.V,153; VI,223; VII,133.
O. revoluta ; R. Brown in Transact. Linn. Soc. X, 19 (1809)	—	—	T.	—	—	—	—	B.fl.V,412	
O. acicularis ; R. Brown, prot. nov. 32 (1830)	—	—	T.	—	—	—	—	B.fl.V,412	

STRANGEA ; Meissner in Hooker's Kew Misc. VII, 66 (1855). (Molloya.)

S. linearis ; Meissner in Hook. Kew Misc. VII, 66 (1855)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,453	M.fr.VI,223; VII,132.
S. cynanchocarpa ; F. v. M., fragm. VII, 132 (1871)	W.A.	—	—	—	—	—	—	B.fl.V,453	M.fr.VII,132.

GREVILLIA ; R. Brown in Transact. Linn. Soc. X, 168 (1809). (Anadenia, Lyssanthe, Stylurus, Manglesia.)

G. Pinaster ; Meissner in Hook. Kew Misc. VII, 76 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,427	M.fr.VI,212.
G. obtusifolia ; Meissner in Hook. Kew Misc. IV, 187 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,427	
G. sparsiflora ; F. v. M., fragm. VI, 206 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,206.
G. macrostylis ; F. v. M., fragm. I, 137 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,211.
G. tripartita ; Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,213.
G. platypoda ; F. v. M., fragm. VI, 205 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,428	M.fr.VI,205.
G. patentiloba ; F. v. M., fragm. I, 137 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,429	M.fr.I,137.
G. pectinata ; R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,429	M.fr.VI,212.
G. plurijuga ; F. v. M., fragm. IV, 84 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,430	M.fr.VI,212.
G. nudiflora ; Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,430	M.fr.I,35; VI,212.
G. stenomera ; F. v. M., fragm. IV, 85 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,430	M.fr.IV,85.
G. Thelemanniana ; Endlicher, nov. stirp. dec. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,431	M.fr.VI,212.
G. concinna ; R. Brown in Transact. Linn. Soc. X, 172 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,431	M.fr.VI,212.
G. Hookeriana ; Meissner in Lehm. pl. Preiss. I, 546 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,432	M.fr.VI,212.
G. Baxteri ; R. Brown, prot. nov. 22 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,432	
G. eriobotrya ; F. v. M., fragm. X, 44 (1876)	W.A.	—	—	—	—	—	—	—	B.fl.X,44.
G. pterosperma ; F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,432	M.fr.VI,210; X,45.
G. stenobotrya ; F. v. M., fragm. IX, 3 (1875)	W.A.	S.A.	—	—	—	Q.	N.A.	—	M.fr.IX,3.
G. eriostachya ; Lindley, Bot. Regist. XXV, App. 36 (1839)	—	—	—	—	—	—	N.A.	B.fl.V,433	M.fr.VI,208&213; X,45.
G. thyrsoides ; Meissner in Hooker's Kew Misc. VII, 77 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,433	
G. Chrysodendron ; R. Brown in Transact. Linn. Soc. X, 176 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,434	M.fr.VI,210.
G. Banksii ; R. Brown in Transact. Linn. Soc. X, 176 (1809)	—	—	—	—	—	Q.	—	B.fl.V,434	M.fr.VI,210; VIII,150; X,46.
G. Caleyi ; R. Brown, prot. nov. 22 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,435	
G. asplenifolia ; R. Brown in J. Knight, prot. 120 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.V,435	M.fr.VI,212.
G. cirsiifolia ; Meissner in Lehm. pl. Preiss. II, 253 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,436	
G. laurifolia ; Sieber in Roem. & Schult. mant. 279 (1822)	—	—	—	—	N.S.W.	—	—	B.fl.V,436	
G. Barklyana ; F. v. M., gen. report 14 & 18 (1861)	—	—	—	V.	—	—	—	B.fl.V,436	M.fr.VII,133; VIII,150; X,46.
G. repens ; F. v. M. in Schlecht. Linnaea XXVI, 355 (1853)	—	—	—	V.	—	—	—	B.fl.V,437	
G. Aquifolium ; Lindley in Mitchell, Three Exped. II, 178 (1838)	—	S.A.	—	V.	—	—	—	B.fl.V,437	
G. ilicifolia ; R. Brown, prot. nov. 21 (1830)	—	S.A.	—	V.	—	Q.	—	B.fl.V,437	M.fr.VI,212.
G. Scortechinii ; F. v. M. in Bailey's synop. of Queensland Flora 435 (1883)	—	—	—	—	N.S.W.	Q.	—	—	
G. Gandichaudii ; R. Brown in Freyc. Voy. Bot. 443, t. 46 (1826)	—	—	—	—	N.S.W.	—	—	B.fl.V,438	
G. acanthifolia ; Cunningham in Field, N.S.W. 328 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.V,438	M.fr.VI,212; VIII,150.
G. bipinnatifida ; R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,439	M.fr.VI,212.
G. armigera ; Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,439	
G. asparagoides ; Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,439	M.fr.IX,123.
G. Treueriana ; F. v. M., fragm. IX, 123 (1875)	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,123.
G. floribunda ; R. Brown, prot. nov. 19 (1830)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,440	
G. cinerea ; R. Brown in Transact. Linn. Soc. X, 173 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,440	
G. alpina ; Lindley in Mitchell, Three Exped. II, 179 (1838)	—	—	—	V.	—	—	—	B.fl.V,441	
G. montana ; R. Brown in Transact. Linn. Soc. X, 172 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,441	
G. obtusiflora ; R. Brown, prot. nov. 19 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,442	
G. arenaria ; R. Brown in Transact. Linn. Soc. X, 172 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,442	
G. mucronulata ; R. Brown in Transact. Linn. Soc. X, 173 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,443	
G. Baueri ; R. Brown in Transact. Linn. Soc. X, 173 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,443	
G. lanigera ; Cunningham in R. Brown, prot. nov. 20 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.V,444	
G. rosmarinifolia ; Cunningham in Field, N. S. Wales, 328 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.V,445	
G. Goodii ; R. Brown in Transact. Linn. Soc. X, 174 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,446	M.fr.VI,211.
G. venusta ; R. Brown in Transact. Linn. Soc. X, 175 (1809)	—	—	—	—	—	Q.	—	B.fl.V,446	

G. longistyla; Hooker in Mitchell, Trop. Austr. 343 (1848)	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,446	M.fr.VI,211.		
G. juncifolia; Hooker in Mitchell, Trop. Austr. 341 (1848)	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,447	M.fr.VI,209; VIII,150.		
G. Wilsoni; Cunningham in Wilson's Voy. 273 (1835)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,447	M.fr.VI,212; X,45.		
G. erectiloba; F. v. M., fragm. X, 44 (1876)	...	...	...	W.A.	—	—	—	—	—	—	—	M.fr.X,44.		
G. lavandulacea; Schlechtendal, Linnaea XX, 586 (1847)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,448			
G. insignis; Kippist in De Candolle, prodr. XIV, 379 (1856)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,448	M.fr.VIII,150; X,45.		
G. Brownii; Meissner in Lehm. pl. Preiss. I, 537 (1845)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,449	M.fr.VI,213.		
G. fasciculata; R. Brown, prot. nov. 20 (1830)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,449			
G. aspera; R. Brown in Transact. Linn. Soc. X, 172 (1809)	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.V,450	M.fr.VII,211.		
G. brachystylis; Meissner in Lehm. pl. Preiss. I, 538 (1845)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,450			
G. saccata; Bentham, Fl. Austr. V, 450 (1870)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,450			
G. Drummondii; Meissner in Lehm. pl. Preiss. I, 536 (1845)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,451			
G. deflexa, F. v. M. in Melb. Chemist 57, p. 72 (1883)	...	...	...	W.A.	—	—	—	—	—	—	—			
G. disjuncta; F. v. M., fragm. VI, 206 (1868)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,451	M.fr.VI,206.		
G. haplantha; F. v. M. in Bentham, Fl. Austr. V, 451 (1870)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,452			
G. pinifolia; Meissner in Hook. Kew Misc. 186 (1852)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,452			
G. Kennedyana; F. v. M. in proceed. R.S. Vict. XXIV, p. 172 (1887)	...	...	...	—	—	—	—	N.S.W.	Q.	—	—			
G. acuaria; F. v. M. in Bentham, Fl. Austr. V., 452 (1870)	..	...	...	W.A.	—	—	—	—	—	—	B.fl.V,452			
G. singuliflora; F. v. M., fragm. VI, 92 (1868)	...	...	...	—	—	—	—	—	Q.	—	B.fl.V,452	M.fr.VI,92.		
G. pauciflora; R. Brown in Transact. Linn. Soc. X, 171 (1809)	...	...	...	W.A.	S.A.	—	—	—	—	—	B.fl.V,453	M.fr.I,135; VI,206,208.		
G. quercifolia; R. Brown, prot. nov. 23 (1830)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,454	M.fr.VI,213.		
G. angulata; R. Brown, prot. nov. 24 (1830)	...	...	...	—	S.A.	—	—	—	—	N.A.	B.fl.V,455			
G. Wickhami; Meissner in Hook. Kew Misc. 187 (1852)	...	...	...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.V,455			
G. agrifolia; Cunningham in R. Brown, prot. nov. 24 (1830)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,455			
G. Cunninghamii; R. Brown, prot. nov. 23 (1830)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,456			
G. pungens; R. Brown in Transact. Linn. Soc. X, 175 (1809)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,456			
G. Huegelii; Meissner in Lehm. pl. Preiss. I, 543 (1845)	...	...	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,457	M.fr.VI,212; X,46.		
G. dimidiata; F. v. M., fragm. III, 146 (1863)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,457	M.fr.III,146.		
G. heliosperma; R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,457	M.fr.VI,210.		
G. refracta; R. Brown in Transact. Linn. Soc. X, 176 (1809)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.V,458			
G. Dryandri; R. Brown in Transact. Linn. Soc. X, 175 (1809)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.V,458	M.fr.VI,209.		
G. polystachya; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.V,459	M.fr.VI,210.		
G. robusta; Cunningham in R. Brown, prot. nov. 24 (1830)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,459	M.fr.VI,210; VII,133.		
G. annulifera; F. v. M., fragm. IV., 85 (1864)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,460	M.fr.IV,85; VI,212; VII,132; XI,20.		
G. leucopteris, Meissner in Hook. Kew Misc. VII, 76 (1855)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,460	M.fr.III,145; IV,85,176; XI,20.		
G. Leucadendron; Cunningham in R. Brown, prot. nov. 25 (1830)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,461	M.fr.I,136-7.		
G. pyramidalis; Cunningham in R. Brown, prot. nov. 25 (1830)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,462			
G. striata; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,462			
G. mimosoides; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	—	—	—	—	—	—	N.A.	B.fl.V,462			
G. Hilliana; F. v. M. in Transact. phil. Inst. Vict. II, 72 (1857)	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.V,463			
G. pinnatifida; Bailey, papers on Q. plants I, 6 (1886)	...	...	...	—	—	—	—	—	Q.	—	—			
G. gibbosa; R. Brown in Transact. Linn. Soc. X, 177 (1809)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.V,463	M.fr.VI,210	...	... P.
G. buxifolia; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,464			
G. phylicoides; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,464	M.fr.VI,213.		
G. sphacelata; R. Brown in Transact. Linn. Soc. X, 174 (1809)	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.V,464			
G. occidentalis; R. Brown in Transact. Linn. Soc. X, 173 (1809)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,465	M.fr.I,136.		
G. accrosa; F. v. M., fragm. I, 136 (1859)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,465	M.fr.I,136.		
G. umbellulata; Meissner in Lehm. pl. Preiss. II, 252 (1847)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.V,465			
G. oxystigma; Meissner in Lehm. pl. Preiss. I, 540 (1845)	...	...	...	W.A.	—	—	—	—	—	—	B.fl V,466			
G. Candolleana; Meissner in Lehm. pl. Preiss. I, 541 (1845)	...	...	...	W.A.	—	—	—	—	—	—	B.fl V,466	M.fr.VI,213.		

G. scabra; Meissner in Lehm. pl. Preiss. I, 541 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,466	
G. Miquelliana; F. v. M. in Transact. Vict. Inst. 132 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.V,467	
G. Victoriae; F. v. M. in Transact. phil. Soc. Vict. I, 107 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.V,468	M.fr.VIII,142,150.
G. punicea; R. Brown in Transact. Linn. Soc. X, 169 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,468	M.fr.XI,123.
G. oleoides; Sieber in Roem. & Schult. mant. 277 (1822)	—	—	—	V.	N.S.W.	—	—	B.fl.V,468	
G. trinervis; R. Brown, prot. nov. 18 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,469	M.fr.VI,211.
G. juniperina; R. Brown in Transact. Linn. Soc. X, 171 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,469	
G. sericea; R. Brown in Transact. Linn. Soc. X, 170 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,470	
G. capitellata; Meissner in Hook. Kew Misc. IV, 187 (1852)	—	—	—	—	N.S.W.	—	—	B.fl.V,470	
G. leiophylla; F. v. M. in Bentham, Fl. Austr. V, 471 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,471	
G. linearis; R. Brown in Transact. Linn. Soc. X, 170 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,471	
G. confertifolia; F. v. M. in Transact. phil. Soc. Vict. I, 22 (1854)	—	—	—	V.	—	—	—	B.fl.V,471	M.fr.VI,211.
G. parviflora; R. Brown in Transact. Linn. Soc. X, 171 (1809)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,472	
G. halmaturina; Tate in Transact. R.S. of S.A. VI, 141 (1883)	—	S.A.	—	—	—	—	—	—	
G. Australis; R. Brown in Transact. Linn. Soc. X, 171 (1809)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,472	M.fr.VII,133.
G. commutata; F. v. M., fragm. VI, 207 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,473	M.fr.VII,207.
G. pinnatisecta; F. v. M. in Bentham, Fl. Austr. V, 473 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,473	
G. argyrophylla; Meissner in Hook. Kew Misc. VII, 75 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,474	M.fr.VI,411.
G. brachystachys; Meissner in Lehm. pl. Preiss. II, 254 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,474	M.fr.VI,211.
G. Endlicheriana; Meissner in Lehm. pl. Preiss. I, 546 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,474	M.fr.VI,211.
G. manglesoides; Meissner in Lehm. pl. Preiss. I, 547 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,475	
G. diversifolia; Meissner in Lehm. pl. Preiss. I, 547 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,475	
G. filifolia; Meissner in Lehm. pl. Preiss. I, 547 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,475	
G. hakeoides; Meissner in Lehm. pl. Preiss. II, 252 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,476	
G. teretifolia; Meissner in Lehm. pl. Preiss. II, 255 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,476	
G. eryngioides; Bentham, Fl. Austr. V, 476 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,476	M.fr.X,46.
G. bracteosa; Meissner in Lehm. pl. Preiss. II, 254 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,477	
G. crithmifolia; R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,477	
G. trachytheca; F. v. M., fragm. VI, 207 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,477	M.fr.VI,207.
G. triternata; R. Brown, prot. nov. 21 (1830)	—	—	—	V.	N.S.W.	—	—	B.fl.V,478	
G. ramosissima; Meissner in De Candolle, prodr. XIV, 388	—	—	—	V.	N.S.W.	—	—	B.fl.V,478	M.fr.VIII,150; X,46.
G. monticola; Meissner in Lehm. pl. Preiss. II, 259 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,478	M.fr.VI,209
G. Muelleri; Bentham, Fl. Austr. V, 479 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,479	
G. trifida; Meissner in Lehm. pl. Preiss. I, 553 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,479	M.fr.VI,209.
G. Synapheae; R. Brown, prot. nov. 23 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,480	
G. flexuosa; Meissner in Lehm. pl. Preiss. I, 553 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,480	
G. leptobotrya; Meissner in Lehm. pl. Preiss. II, 256 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,480	M.fr.VI,209.
G. brevicuspis; Meissner in Lehm. pl. Preiss. II, 256 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,481	
G. intricata; Meissner in Hook. Kew Misc. VII, 74 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,481	
G. didymobotrya; Meissner in Hook. Kew Misc. IV, 186 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,481	M.fr.I,136; VI,208; X,45.
G. polybotrya; Meissner in Hook. Kew Misc. IV, 185 (1852)	W.A.	—	—	—	—	—	N.A.	B.fl.V,482	M.fr.IV,129; VI,208.
G. nematophylla; F. v. M., fragm. I, 136 (1859)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,482	M.fr.I,136; VI,208.
G. anethifolia; R. Brown, prot. nov. 21 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,482	M.fr.VI,209.
G. paradoxa; F. v. M., fragm. VI, 246 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,483	M.fr.VI,246.
G. petrophiloides; Meissner in Lehm. pl. Preiss. II, 257 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,483	M.fr.VI,209; X,46.
G. tenuiflora; Meissner in Lehm. pl. Preiss. I, 554 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,483	
G. pulchella; Meissner in Lehm. pl. Preiss. I, 553 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,484	M.fr.VI,209.
G. rudis; Meissner in Hook. Kew Misc. VII, 73 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,484	
G. apiciloba; F. v. M., fragm. X, 45 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,45.
G. Shuttleworthiana; Meissner in Lehm. pl. Preiss. II, 258 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,484	

G. integrifolia; Meissner in De Candolle, prodr. XIV, 385 (1856)	W.A.	—	—	—	—	—	—	B.fl.V,485		
G. stenocarpa; F. v. M. in Bentham, Fl. Austr. V, 485 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,485		
G. acrobotrya; Meissner in Hooker's Kew Misc. VII, 74 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,486	M.fr.VI,213.	
G. glabrata; Meissner in Lehm. pl. Preiss. I, 549 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,486		
G. ornithopoda; Meissner in Lehm. pl. Preiss. II, 256 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,486		
G. paniculata; Meissner in Lehm. pl. Preiss. I, 550 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,487		
G. biternata; Meissner in Lehm. pl. Preiss. I, 549 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,487	M.fr.VI,213.	
G. triloba; Meissner in Hooker's Kew Misc. VII, 74 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,487	M.fr.VI,213.	
G. amplexans; F. v. M. in Bentham, Fl. Austr. V, 488 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,488		
G. vestita; Meissner in Lehm. pl. Preiss. I, 548 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,488	M.fr.VI,213.	
G. tridentifera; Meissner in Lehm. pl. Preiss. 1, 547 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,488		
G. erinacea; Meissner in Hooker's Kew Misc. VII, 74 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,489	M.fr.X,45.	
HAKEA; Schrader, sert. Hannov. I, fasc. 3, 27, t. 17 (1797). (Conchium.)										
H. chordophylla; F. v. M. in Hooker's Kew Misc. IX, 23 (1857)	—	—	—	—	—	Q.	N.A.	B.fl.V,495	M.fr.V,190.	
H. Cunninghamii; R. Brown, prot. nov. 26 (1830)	W.A.	—	—	—	—	—	N.A.	B.fl.V,495	M.fr.VI,190.	
H. lorea; R. Brown, prot. nov. 25 (1830)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,495	M.fr.VI,189; VII,133.	
H. Fraseri; R. Brown, prot. nov. 26 (1830)	—	—	—	—	N.S.W.	—	—	B.fl.V,496		
H. macrocarpa; Cunningham in R. Brown, prot. nov. 30 (1830)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.V,496	M.fr.VI,190,210.	
H. arborescens; R. Brown in Transact. Linn. Soc. X, 187 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.V,497	M.fr.VI,190.	
H. Edneiana; Tate in Transact. R. Soc. of S.A. VII, 70 (1884)	—	S.A.	—	—	—	—	—	—		
H. Persiehana; F. v. M. in Austr. Journ. of Pharm. I, 430 (1886)	—	—	—	—	—	Q.	—	—		
H. stenophylla; Cunningham in De Candolle, prodr. XIV, 417 (1856)	—	—	—	—	—	—	N.A.	B.fl.V,497		
H. trineura; F. v. M., fragm. III, 146 (1863)	—	—	—	—	—	Q.	—	B.fl.V,497	M.fr.VI,216.	
H. cyclocarpa; Lindley, Bot. Regist. XXV, App. 36 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,498		
H. crassifolia; Meissner in Lehm. pl. Preiss. I, 570 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,498	M.fr.VI,216.	
H. Brooksiana; F. v. M. in Austr. Journ. of Pharm. I, 430 (1886)	W.A.	—	—	—	—	—	—	—		
H. pandanocarpa; R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,499	M.fr.VI,216.	
H. Roei; Bentham, Fl. Austr. V, 499 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,499		
H. adnata; R. Brown, prot. nov. 26 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,499		
H. obliqua; R. Brown in Transact. Linn. Soc. X, 180 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,500	M.fr.VI,218.	
H. Hookeriana; Meissner in Hook. Kew Misc. IV, 208 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,500	M.fr.VI,216.	
H. incrassata; R. Brown, prot. nov. 29 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,500		
H. flabellifolia; Meissner in Hook. Kew Misc. VII, 116 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,501		
H. Brownii; Meissner in Lehm. pl. Preiss. I, 569 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,501		
H. Baxteri; R. Brown, prot. nov. 28 (1830)	W.A.	S.A.	—	—	—	—	—	B.fl.V,501		
H. ceratophylla; R. Brown in Transact. Linn. Soc. X, 184 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,501	M.fr.IV,49; VI,217.	
H. lasiantha; R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,502	M.fr.VI,215.	
H. eriantha; R. Brown, prot. nov. 29 (1830)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,502	M.fr.VI,217.	
H. megalosperma; Meissner in Hook. Kew Misc. VII, 117 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,503		
H. clavata; Labillardière, Nov. Holl. pl. spec. I, 31, t. 41 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,503	M.fr.VI,216.	
H. orthorrhyncha; F. v. M., fragm. V, 214 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,503	M.fr.VI,214.	
H. Candolleana; Meissner in Lehm. pl. Preiss. II, 262 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,504	M.fr.VI,215.	
H. trifurcata; R. Brown in Transact. Linn. Soc. X, 183 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,504	M.fr.VI,220.	
H. erinacea; Meissner in Lehm. pl. Preiss. I, 559 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,505	M.fr.VI,220.	
H. platysperma; Hooker, icon. pl. t. 433 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,505		
H. brachyptera; Meissner in Hook. Kew Misc. IV, 208 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,505	M.fr.VI,217.	
H. Kippistiana; Meissner in Hooker's Kew Misc. VII, 115 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,506		
H. Preissii; Meissner in Lehm. pl. Preiss. I, 557 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,506	M.fr.VI,219.	
H. pugioniformis; Cavanilles, Annal. hist. nat. I, 213, t. 11 (1800)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,506	M.fr.VI,217.	
H. vittata; R. Brown in Transact. Linn. Soc. X, 182 (1809)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,507		

H. rostrata; F. v. M. in Schlecht. Linnaea XXVI, 259 (1853)	—	S.A.	—	V.	—	—	—	B.fl.V,508	M.fr.VI,218; VII,133.
H. rugosa; R. Brown in Transact. Linn. Soc. X, 179 (1809)	—	S.A.	—	V.	—	—	—	B.fl.V,508	M.fr.VI,218.
H. Epiglottis; Labillardière, Nov. Holl. pl. spec. I, 30, t. 40 (1804)	—	—	T.	—	—	—	—	B.fl.V,508	
H. amplexicaulis; R. Brown in Transact. Linn. Soc. X, 184 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,509	M.fr.VI,217; VII,133.
H. glabella; R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,509	
H. auriculata; Meissner in Hook. Kew Misc. VII, 116 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,510	M.fr.VI,217.
H. cristata; R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,510	
H. linearis; R. Brown in Transact. Linn. Soc. X, 183 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,511	
H. stenocarpoides; F. v. M. in Bentham Fl. V, 511 (1870)...	W.A.	—	—	—	—	—	—	B.fl.V,511	
H. pedunculata; F. v. M. in Melb. Chem. July (1883)	—	—	—	—	—	Q.	—	—	
H. ruscifolia; Labillardière, Nov. Holl. pl. spec. I, 30, t. 39 (1804)	W.A.	—	—	—	—	—	—	B.fl.V,511	
H. saligna; R. Brown in J. Knight, prot. 108 (1809)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,512	M.fr.VI,217.
H. verrucosa; F. v. M., fragm. V, 25 (1865)	W.A.	—	—	—	—	—	—	B.fl.V,512	M.fr.V,25; VI,218.
H. rhombales; F. v. M., fragm. X, 90 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.X,90.
H. purpurea; Hooker in Mitch. Trop. Austr. 348 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,513	
H. pubescens; Schrader. sert. Hannover. I, 25 (1795), ...	—	—	—	—	N.S.W.	—	—	B.fl.V,513	M.fr.VI,218.
H. propinqua; Cunningham in Field's N.S.Wales, 327 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.V,513	M.fr.V,26.
H. Macraeana; F. v. M. in Austr. Journ. of Phar. I, 430 (1886)	—	—	—	—	N.S.W.	—	—	—	
H. nodosa; R. Brown in Transact. Linn. Soc. X, 179 (1809)	—	S.A.	T.	V.	—	—	—	B.fl.V,514	M.fr.V,26; VI,217.
H. sericea; Schrader sert. Hannover. I, 27 (1795)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,514	M.fr.VI,218.
H. leucoptera; R. Brown in Transact. Linn. Soc. X, 180 (1809)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,515	M.fr.VI,219; VII,133.
H. cycloptera; R. Brown in Transact. Linn. Soc. X, 182 (1809)	—	S.A.	—	—	—	—	—	B.fl.V,515	M.fr.VI,217.
H. microcarpa; R. Brown in Transact. Linn. Soc. X, 182 (1809)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,516	M.fr.VI,217.
H. recurva; Meissner in Hook. Kew Misc. IV, 207 (1852)...	W.A.	—	—	—	—	—	—	B.fl.V,516	M.fr.VI,218.
H. circumalata; Meissner in Hook. Kew Misc. VII, 114 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,516	M.fr.VI,219.
H. commutata; F. v. M., fragm. V, 26 (1865)	W.A.	—	—	—	—	—	—	B.fl.V,517	M.fr.VI,219.
H. strumosa; Meissner in Hook. Kew Misc. IV, 208 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,517	M.fr.VI,219.
H. multilineata; Meissner in Lehm. pl. Preiss. II, 261 (1847)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.V,518	M.fr.V,25; VI,215.
H. laurina; R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,518	M.fr.IV,130,216.
H. obtusa; Meissner in Hook. Kew Misc. IV, 209 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,519	
H. cinerea; R. Brown in Transact. Linn. Soc. X, 186 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,519	
H. corymbosa; R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,519	M.fr.VI,216.
H. undulata; R. Brown in Transact. Linn. Soc. X, 185 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,520	M.fr.VI,216.
H. petiolaris; Meissner in Lehm. pl. Preiss. I, 577 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,520	
H. neurophylla; Meissner in Hook. Kew Misc. VII, 117 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,521	
H. loranthifolia; Meissner in Lehm. pl. Preiss. I, 574 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,521	
H. cucullata; R. Brown, prot. nov. 30 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,521	M.fr.VI,216.
H. ferruginea; Sweet, Fl. Austral. t. 45 (1828)	W.A.	—	—	—	—	—	—	B.fl.V,522	M.fr.VI,216.
H. smilacifolia; Meissner in Lehm. pl. Preiss. I, 567 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,522	M.fr.VI,217.
H. elliptica; R. Brown in Transact. Linn. Soc. X, 187 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,523	M.fr.VI,216.
H. ambigua; Meissner in Lehm. pl. Preiss. II, 260 (1847)...	W.A.	—	—	—	—	—	—	B.fl.V,523	M.fr.VI,216.
H. plurinervia; F. v. M. in Benth. Fl. Austr. V, 523 (1870)	—	—	—	—	—	Q.	—	B.fl.V,523	
H. dactyloides; Cavanilles, Anal. hist. nat. I, 215, t. 12 (1800)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,524	M.fr.VI,215.
H. ulicina; R. Brown, prot. nov. 29 (1830)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,524	
H. falcata; R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,524	
H. pycnoneura; Meissner in Hook. Kew Misc. VII, 117 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,525	M.fr.VI,215.
H. stenocarpa; R. Brown, prot. nov. 29 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,525	
H. marginata; R. Brown in Transact. Linn. Soc. X, 185 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,356	M.fr.VI,216.
H. myrtoides; Meissner in Lehm. pl. Preiss. I, 577 (1845)...	W.A.	—	—	—	—	—	—	B.fl.V,526	
H. costata; Meissner in Lehm. pl. Preiss. I, 575 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,526	

H. oleifolia; R. Brown in Transact. Linn. Soc. X, 185 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,527	M.fr.VI,215.
H. florida; R. Brown in Transact. Linn. Soc. X, 183 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,527	
H. varia; R. Brown in Transact. Linn. Soc X, 183 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,527	M.fr.VI,219.
H. sulcata; R. Brown in Transact. Linn. Soc. X, 180 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,528	M.fr.VI,219.
H. Meissneriana; Kippist in Hook. Kew Misc. VII, 114 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,529	M.fr.VI,219.
H. subsulcata; Meissner in Lehm. pl. Preiss. I, 555 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,529	M.fr.VI,219.
H. Lehmanniana; Meissner in Lehm. pl. Preiss. I, 557 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,529	M.fr.VI,218.
H. nitida; R. Brown in Transact. Linn. Soc. X, 184 (1809)	W.A. S.A.	—	—	—	—	—	—	B.fl.V,530	M.fr.V,72; VI,217.
H. Oldfieldii; Bentham, Fl. Austr. V, 530 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,530	
H. suaveolens; R. Brown in Transact. Linn. Soc. X, 182 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,531	M.fr.VI,220.
H. lissocarpha; R. Brown, prot. nov. 27 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,531	M.fr.VI,219.
H. bipinnatifida; R. Brown, prot. nov. 28 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,532	
CARNARVONIA; F. v. M., fragm. VI, 81 (1867).									
C. aralifolia; F. v. M., fragm. VI, t. 55 (1867)	—	—	—	—	—	Q.	—	B.fl.V,410	M.fr.VI,81,247,254; VII,59.
BUCKINGHAMIA; F. v. M., fragm. VI, 248 (1868).									
B. celsissima; F. v. M., fragm. VI, 248 (1868)	—	—	—	—	—	Q.	—	B.fl.V,532	M.fr.VI,248.
DARLINGIA; F. v. M., fragm. V, 152 (1866).									
D. spectatissima; F. v. M., fragm. V, 152 (1866)	—	—	—	—	—	Q.	—	B.fl.V,533	M.fr.V,24; X,90.
HOLLANDAEA; F. v. M. in Melb. Chem. and Drugg., new. ser. II, 173 (1887).									
H. Sayeri; F. v. M. in Melb. Chem. and Drugg., new ser. II, 173 (1887) ...	—	—	—	—	—	Q.	—	—	
CARDWELLIA; F. v. M., fragm. V, 23 (1865).									
C. sublimis; F. v. M., fragm. V, 24 (1865)	—	—	—	—	—	Q.	—	B.fl.V,538	M.fr.V,38,73,152.
STENOCARPUS; R. Brown in Transact. Linn. Soc. X, 201 (1809). (Cybele, 1809, Agnostus.)									
S. sinuatus; Endlicher, gen. pl. suppl. IV, 88 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,539	M.fr.V,154; VI,224.
S. salignus; R. Brown in Transact. Linn. Soc. X, 202 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,539	M.fr.I,134,244; VI,224; VIII,133.
S. Cunninghamii; R. Brown, prot. nov. 34 (1830)	—	—	—	—	—	—	N.A.	B.fl.V,540	
LOMATIA; R. Brown in Transact. Linn. Soc. X, 199 (1809). (Tricondylus, 1809.)									
L. fraxinifolia; F. v. M. in Benth. Fl. Austr. V, 536 (1870)	—	—	—	—	—	Q.	—	B.fl.V,536	
L. ilicifolia; R. Brown in Transact Linn. Soc. X, 200 (1809)	—	—	—	V.	N.S.W.	—	—	B.fl.V,536	M.fr.V,153; VII,133.
L. Fraseri; R. Brown, prot. nov. 34 (1830)	—	—	—	V.	N.S.W.	Q.	—	—	
L. longifolia; R. Brown in Transact. Linn. Soc. X, 200 (1809)	—	—	—	V.	N.S.W.	—	—	B.fl.V,537	M.fr.V,153.
L. silaifolia; R. Brown in Transact. Linn. Soc. X, 199 (1809)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,537	M.fr.V,153; VI,191.
L. tinctoria; R. Brown in Transact. Linn. Soc. X, 199 (1809)	—	—	T.	—	—	—	—	B.fl.V,537	M.fr.VI,224.
L. polymorpha; R. Brown in Transact. Linn. Soc. X, 200 (1809)	—	—	T.	—	—	—	—	B.fl.V,538	
EMBOTHRIUM; R. & G. Forster, charact. gen. 15, t. 8 (1776). (Oreocallis.)									
E. Wickhami; Hill & F. v. M., fragm. VIII, 164 (1874)	—	—	—	—	—	Q.	—	—	M.fr.IX,194.
TELOPEA; R. Brown in Transact. Linn. Soc. X, 197 (1809). (Hylogyne, 1809).									
T. speciosissima; R. Brown in Transact. Linn. Soc. X, 198 (1809)	—	—	—	—	N.S.W.	—	—	B.fl.V,534	M.fr.X,90.
T. oreades; F. v. M., Annual Report 18 (1861)	—	—	—	V.	N.S.W.	—	—	B.fl.V,534	M.fr.II,170; VII,133.
T. truncata; R. Brown in Transact. Linn. Soc. X, 198 (1809)	—	—	T.	—	—	—	—	B.fl.V,535	M.fr.II,171.
BANKSIA; Linné, fil. suppl. 15 et 126 (1781).									
B. pulchella; R. Brown in Transact. Linn. Soc. X, 202 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,544	M.fr.VII,54.
B. Meissneri; Lehmann, plant. Preiss. I, 582 (1845)...	W.A.	—	—	—	—	—	—	B.fl.V,545	

B. nutans; R. Brown in Transact. Linn. Soc. X, 203 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,545	M.fr.IV,108; VII,54.
B. sphaerocarpa; R. Brown in Transact. Linn. Soc. X, 203 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
B. tricuspis; Meissner in Hook. Kew Misc. VII, 118 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
B. occidentalis; R. Brown in Transact. Linn. Soc. X, 204 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,546	M.fr.VII,54.
B. litoralis; R. Brown in Transact. Linn. Soc. X, 204 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,547	M.fr.VII,54.
B. ericifolia; Linné fil., suppl. 127 (1781)	—	—	—	—	N.S.W.	—	—	B.fl.V,547	M.fr.VII,54.
B. spinulosa; Smith, Specim. Bot. New Holl. 13, t. 4 (1793)	—	—	—	—	N.S.W.	—	—	B.fl.V,547	
B. collina; R. Brown in Transact. Linn. Soc. X, 204 (1809)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,548	M.fr.VII,54.
B. verticillata; R. Brown in Transact. Linn. Soc. X, 207 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,548	
B. dryandroides; Baxter in Sweet, hort. Brit., 350 (1827)	W.A.	—	—	—	—	—	—	B.fl.V,549	M.fr.VII,58.
B. Brownii; Baxter in R. Brown prot. nov. 37 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,549	M.fr.VII,58.
B. attenuata; R. Brown in Transact. Linn. Soc. X, 209 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,549	M.fr.VII,55.
B. media; R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,550	M.fr.IV,109.
B. Solandri; R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,550	M.fr.VII,58.
B. Goodii; R. Brown, prot. nov. 36 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,550	M.fr.VII,58.
B. petiolaris; F. v. M., fragm. IV, 109 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,551	M.fr.IV,109.
B. repens; Labillardière, Voy. I, 411, t. 23 (1798)	W.A.	—	—	—	—	—	—	B.fl.V,551	M.fr.IV,108,177; VII,58.
B. prostrata; R. Brown, prot. nov. 36 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,551	M.fr.VII,57.
B. grandis; Willdenow, spec. plant. I, 535 (1797)	W.A.	—	—	—	—	—	—	B.fl.V,552	M.fr.VII,57.
B. quercifolia; R. Brown in Transact. Linn. Soc. X, 210 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,552	M.fr.VII,57.
B. Baueri; R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,552	M.fr.IV,107; VII,56.
B. marginata; Cavanilles, Anal. hist. nat. I, 227, t. 13 (1800)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,553	M.fr.VII,55,133; X,90.
B. integrifolia; Linné fil., suppl. 127 (1781)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,554	M.fr.VII,55.
B. dentata; Linné fil., suppl. 127 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.V,555	M.fr.VII,57.
B. Robur; Cavanilles in Ansl. hist. nat. I, 226 (1800)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,555	M.fr.VII,56.
B. serrata; Linné fil., suppl. 126 (1781)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,556	M.fr.VII,56.
B. serratifolia; Salisbury, prodr. 51 (1796)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,556	M.fr.VII,56.
B. ornata; F. v. M. in Schlecht. Linnaea XXVI, 352 (1853)	—	S.A.	—	V.	—	—	—	B.fl.V,557	M.fr.VII,56.
B. coccinea; R. Brown in Transact. Linn. Soc. X, 207 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,557	M.fr.VII,56.
B. Sceptrum; Meissner in Hook. Kew Misc. VII, 120 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,557	M.fr.VII,56.
B. Menziesii; R. Brown, prot. nov. 36 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,558	
B. laevigata; Meissner in Hook. Kew Misc. IV, 210 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,558	M.fr.VII,56.
B. Hookeriana; Meissner in Hook. Kew Misc. VII, 119 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,558	
B. prionotes; Lindley, Bot. Regist. XXV, App. 34 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,558	M.fr.VII,56.
B. Victoriae; Meissner in Hook. Kew Misc. VII, 119 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
B. speciosa; R. Brown in Transact. Linn. Soc. X, 210 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
B. Baxteri; R. Brown, prot. nov. 36 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,559	M.fr.VII,58.
B. praemorsa; Andrews, Bot. repos. t. 258 (1802)	W.A.	—	—	—	—	—	—	B.fl.V,560	M.fr.VII,56.
B. Lehmanniana; Meissner in Hook. Kew Misc. 210 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,560	M.fr.VII,57.
B. Caleyi; R. Brown, prot. nov. 35 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,560	M.fr.VII,57.
B. Lindleyana; Meissner in Hook. Kew Misc. VII, 120 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,561	M.fr.VII,55.
B. elegans; Meissner in Hook. Kew Misc. VII, 119 (1855)...	W.A.	—	—	—	—	—	—	B.fl.V,561	
B. Candolleana; Meissner in Hook. Kew Misc. VII, 118 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,561	M.fr.VII,58.
B. ilicifolia; R. Brown in Transact. Linn. Soc. X, 211 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,561	M.fr.VII,58.

DRYANDRA; R. Brown in Transact. Linn. Soc. X, 211 (1809). (Josephia, Hemiclidia.)

D. quercifolia; Meissner in Hook. Kew Misc. IV, 210 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
D. praemorsa; Meissner in Lehm. pl. Preiss. II, 265 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
D. cuneata; R. Brown in Transact. Linn. Soc. X, 212 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,566	M.fr.VII,50.
D. falcata; R. Brown in Transact. Linn. Soc. X, 213 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,567	
D. armata; R. Brown in Transact. Linn. Soc. X, 212 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,567	M.fr.VII,50.

D. longifolia; R. Brown in Transact. Linn. Soc. X, 215 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,568					
D. Fraseri; R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,568	M.fr.VII,52.				
D. floribunda; R. Brown in Transact. Linn. Soc. X, 212 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VI,92; VII,50.				
D. carduacea; Lindley, Bot. Regist. XXV, App. 33 (1839)...	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VII,51.				
D. carlinoides; Meissner in Lehm. pl. Preiss. II, 267 (1847)	W.A.	—	—	—	—	—	—	B.fl.V,569	M.fr.VII,53.				
D. polycephala; Bentham, Fl. Austr. V, 570 (1870)...	W.A.	—	—	—	—	—	—	B.fl.V,570					
D. Kippistiana; Meissner in Hook. Kew Misc. VII, 122 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,570	M.fr.VII,52.				
D. squarrosa; R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,52.				
D. Serra; R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,51.				
D. concinna; R. Brown, prot. nov. 38 (1830)...	W.A.	—	—	—	—	—	—	B.fl.V,571	M.fr.VII,52.				
D. foliolata; R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,572	M.fr.VII,51.				
D. stupposa; Lindley, Bot. Regist. XXV, App. 33 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,572	M.fr.VII,51.				
D. nobilis; Lindley, Bot. Regist. XXV, App. 33 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,572					
D. mucronulata; R. Brown in Transact. Linn. Soc. X, 213 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,573					
D. formosa; R. Brown in Transact. Linn. Soc. X, 213, t. 3 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,573	M.fr.VII,51.				
D. Baxteri; R. Brown, prot. nov. 38 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,573	M.fr.VII,51.				
D. nivea; R. Brown in Transact. Linn. Soc. X, 214 (1809)...	W.A.	—	—	—	—	—	—	B.fl.V,574	M.fr.VII,52.				
D. Arctotidis; R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,574					
D. nana; Meissner in Hook. Kew Misc. VII, 121 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,575	M.fr.VII,52.				
D. Preissii; Meissner in Lehm. pl. Preiss. I, 599 (1845)	W.A.	—	—	—	—	—	—	B.fl.V,575	M.fr.VII,54.				
D. sclerophylla; Meissner in Hook. Kew Misc. VII, 123 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,576	M.fr.VII,52.				
D. pulchella; Meissner in Hook. Kew Misc. IV, 211 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,576					
D. plumosa; R. Brown in Transact. Linn. Soc. X, 214 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,576	M.fr.VII,51.				
D. senecionifolia; R. Brown. prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.				
D. vestita; Kippist in Hook. Kew Misc. VII, 121 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.				
D. cirsioides; Meissner in Hook. Kew Misc. IV, 211 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,52.				
D. Hewardiana; Meissner in Hook. Kew Misc. IV, 210 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,577	M.fr.VII,53.				
D. patens; Bentham, Fl. Austr. V, 578 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,578					
D. conferta; Bentham, Fl. Austr. V, 578 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,578					
D. horrida; Meissner in Hook. Kew Misc. IV, 211 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,52.				
D. serratuloides; Meissner in Hook. Kew Misc. VII, 123 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,52.				
D. comosa; Meissner in Hook. Kew Misc. IV, 211 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,579	M.fr.VII,53.				
D. Shuttleworthiana; Meissner in Hook. Kew Misc. VII, 122 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,580	M.fr.VII,52.				
D. speciosa; Meissner in Hook. Kew Misc. IV, 211 (1852)...	W.A.	—	—	—	—	—	—	B.fl.V,580					
D. tridentata; Meissner in Hook. Kew Misc. VII, 120 (1855)	W.A.	—	—	—	—	—	—	B.fl.V,580	M.fr.VII,53.				
D. tenuifolia; R. Brown in Transact. Linn. Soc. X, 215 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,581	M.fr.VII,53.				
D. proteoides; Lindley, Bot. Regist. XXV, App. 33 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,582					
D. runcinata; Meissner in Hook. Kew Misc. IV, 210 (1852)	W.A.	—	—	—	—	—	—	B.fl.V,582					
D. obtusa; R. Brown in Transact. Linn. Soc. X, 214 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,582	M.fr.V,185.				
D. bipinnatifida; R. Brown, prot. nov. 39 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,583					
D. pteridifolia; R. Brown in Transact. Linn. Soc. X, 215 (1809)	W.A.	—	—	—	—	—	—	B.fl.V,583	M.fr.VII,54.				
D. calophylla; R. Brown, prot. nov. 40 (1830)	W.A.	—	—	—	—	—	—	B.fl.V,583	M.fr.VII,54.				

CORNACEAE.

HUMBOLDT, Bonpland & Kunth, nov. gen. Americ. III, 430 (1818).—Genus 1, Species 1.

STYLIDIUM; Loureiro, Fl. Cochinch. I, 220 (1790). (Marlea, Rhytidandra, Pseudalangium.)

S. Vitiense; F. v. M., Cens. 74 (1882) from Asa Gray (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,386	M.fr.II,84,176	...	...	P.	

RUBIACEAE.

A. L. de Jussieu, gen. plant. 196 (1789), from B. de Jussieu (1759).—Genera 30, Species 127.

SARCOCEPHALUS ; Afzelius in Transact. hort. soc. Lond. V, 422, t. 18 (1824).																	
S. cordatus ; Miquel, Fl. Ind. Batav. II, 133 (1856)...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,402	...	...	...	As., P.	
OLDENLANDIA ; Plumier, nov. plant. Americ. gen. 42, t. 36 (1703). (Hedyotia, Synaptantha, Anotis.)																	
O. corymbosa ; Linné, sp. pl. 119 (1753)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VII,45	...	...	As., Afr.
O. paniculata ; Linné, spec. plant. edit. sec. 1667 (1763)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VII,45; VIII,47	...	As., P.
O. psychotrioides ; F. v. M. in Viet. Naturalist VI, 54 (1889)	...	...	...	...	—	—	—	—	—	Q.	—	—	...	...	...	...	As., P.
O. auricularia ; F. v. M., Cens. of Austr. pl. 74 (1882) from Linné (1753)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,404					
O. polyclada ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.VIII,146.				
O. coerulescens ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,404	M.fr.IV,38.				
O. spermacocoides ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.VIII,146.				
O. Cronchiana ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	—	—	—	—	—	—	N.A.	—	M.fr.X,85.				
O. mitrasaemoides ; F. v. M., Cens. of Austr. pl. 74 (1882)...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,404	M.fr.IV,37.				
O. galioides ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	W.A.	—	—	—	—	Q.	N.A.	B.fl.III,404	M.fr.IV,38.				
O. seleranthoides ; Bentham & J. Hooker, gen. pl. II, 88 (1873)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,405	M.fr.IV,39.				
O. elatinoides ; F. v. M., Cens. of Austr. pl. 74 (1882) from Benth. (1866)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,405	M.fr.VIII,147.				
O. tillaeacea ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,405	M.fr.IV,39.				
O. trachymenioides ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,405	M.fr.IV,40.				
O. pterospora ; F. v. M., Cens. of Austr. pl. 74 (1882)	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,406	M.fr.IV,40,177.				
DENTELLA ; R. et G. Forster, char. gen. 25, t. 13 (1776). (Lippaya.)																	
D. repens ; R. & G. Forster, char. gen. 25, t. 13 (1776)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,406	M.fr.IX,187	...	...	As., P.	
OPHIORRHIZA ; Linné, Fl. Zeylan. 190 et 239 (1747).																	
O. Australiana ; Bentham, Fl. Austr. III, 407 (1866)	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,407	M.fr.IX,187.				
ABBOTTIA ; F. v. M., fragm. IX, 181 (1875).																	
A. singularis ; F. v. M., fragm. IX, 181 (1875)	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,181.			
GARDENIA ; Ellis in Phil. Transact. LI, 935, t. 23 (1761).																	
G. edulis ; F. v. M., fragm. I, 54 (1858)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,408	M.fr.I,54.		
G. resinosa ; F. v. M., fragm. I, 54 (1858)	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,408	M.fr.I,54.		
G. pyriformis ; Cunningham in Benth. Fl. Austr. III, 409 (1866)...	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III.409			
G. megasperma ; F. v. M., fragm. I, 54 (1858)	...	...	...	...	...	...	—	—	—	—	—	—	N.A.	B.fl.III,409	M.fr.I,54.		
G. Macgillivraei ; Bentham, Fl. Austr. III, 409 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,409	M.fr.VII,46.		
G. ochreata ; F. v. M., fragm. I, 55 (1858)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,409	M.fr.155;VII,46.		
G. auffrutieosa ; R. Brown in Benth. Fl. Austr. III, 410 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,410			
G. fucata ; R. Brown in Benth. Fl. Austr. III, 410 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.III,410			
G. Jardinei ; F. v. M. in Benth. Fl. Austr. III, 410 (1866)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,410	M.fr.VII,46.		
RANDIA ; Houston in Linné, hort. Cliffort. 485 (1737).																	
R. hirta ; F. v. M., fragm. VII, 46 (1869)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	—	M.fr.IX,181.		
R. chartacea ; F. v. M., fragm. IX, 180 (1875)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,410	M.fr.VII,47.		
R. Benthamiana ; F. v. M., fragm. IX, 180 (1875)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,180.		
R. Moorei ; F. v. M. in Benth. Fl. Austr. III, 411 (1866)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,411			
R. stipularis ; F. v. M., Papuan plants I, 69 (1876)	...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,47;IX,70,181.		
R. Fitzalani ; F. v. M., Rep. Burdek. Exped. 12 (1860)	...	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.III,411	M.fr.V,19;IX,180.		
R. sessilis ; F. v. M., fragm. VII, 47 (1869)	...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.VII,47.		
R. densiflora ; Bentham, Fl. Hongk. 155 (1861)	...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,412	M.fr.II,132; III,166; IX,180	As.	

DIPLOSPORA ; De Candolle, prodr. IV, 477 (1830). (Discospermum.)

D. Australis ; Bentham, Fl. Austr. III, 413 (1866) ...	—	—	—	—	—					Q.	—	B.fl.III,413					
D. ixoroides ; F. v. M., fragm. IX, 182 (1875) ...	—	—	—	—	—					Q.	—	B.fl.III,416	M.fr.IX,182.				

IXORA ; Linné, syst. nat. 8 (1735), gen. pl. 27 (1737). (Pavetta, Tarenna, Webera, Stylocoryne.)

I. Pavetta ; Roxburgh, Fl. Ind. I, 395, ed. Carey (1820) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,414	M.fr.IX,182	...	...	As., P.	
I. tomentosa ; Roxburgh, Fl. Ind. I, 396, ed. Carey (1820)...	—	—	—	—	—					—	N.A.	B.fl.III,414	...	...	...	As.	
I. coccinea ; Linné, sp. pl. 110 (1753) ...	—	—	—	—	—					—	N.A.	B.fl.III,415	...	...	...	As.	
I. Timorensis ; Decaisne in Nouv. Ann. du Mus. III, 419 (1834) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,415	M.fr.V,18; IX,183	...	As.		
I. Becklerii ; Bentham, Fl. Austr. III, 415 (1866) ...	—	—	—	—	—	N.S.W.				Q.	—	B.fl.III,415	M.fr.IX,183.				
I. pentamera ; Bentham, Fl. Austr. III, 416 (1866) ...	—	—	—	—	—					—	N.A.	B.fl.III,416					
I. Dallachyana ; F. v. M., Cens. of Austr. pl. 75 (1882) from Bentham (1866) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,412					
I. expandens ; F. v. M., Cens. of Austr. pl. 75 (1882) ...	—	—	—	—	—					—	N.A.	—	M.fr.VI,25.				

COFFEA ; Linné, syst. nat. 8 (1735); gen. pl. 55 (1737) from Ray (1691).

C. Bengalensis ; Roxburgh, hort. Bengal. 15 (1814) ...	—	—	—	—	—					Q.	—	—	M.fr.VIII,147	...	...	As.	

TIMONIUS ; Rumphius, herb. Amboin. III, 216, t. 140 (1743). (Polyphragmon.)

T. Rumphii ; De Candolle, prodr. IV, 461 (1830) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,417	M.fr.II,134; IX,187	...	As., P.		

SCYPHIPHORA ; K. F. Gaertner, de fruct. III, 91, t. 196 (1805). (Epithinia.)

S. hydrophylacea ; K. F. Gaertner, de fruct. III, 91, t. 196 (1805) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,418	M.fr.VII,46; IX,187	...	As., P.		

GUETTARDA ; Osbeck in Linné, spec. plant. 991 (1753). (Antirrhoea, Guettardella, Bobea, Haplocaryon).

G. speciosa ; Linné, spec. plant. 991 (1753) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,419	M.fr.IX,183	...	...	As., Afr., P.	
G. tenuiflora ; F. v. M., fragm. VII, 48 (1869) ...	—	—	—	—	—					Q.	—	B.fl.III,418	M.fr.IX,183.				
G. putaminosa ; F. v. M., fragm. IX, 183 (1875) ...	—	—	—	—	—					Q.	—	B.fl.III,419	M.fr.IV,92; V,212.				
G. myrtoides ; F. v. M., fragm. IX, 184 (1875) ...	—	—	—	—	—					Q.	—	—	M.fr.IX,184.				

HODGKINSONIA ; F. v. M., fragm. II, 132 (1861).

H. ovatiflora ; F. v. M., fragm. II, 132 (1861) ...	—	—	—	—	—	N.S.W.				Q.	—	B.fl.III,420	M.fr.III,166.				

CANTHIUM ; Lamarck, Encycl. méthod. I, 602 (1783). (Psydrax, Plectronia partly.)

C. cymosum ; Persoon, synops. plant. I, 200 (1805) ...	—	—	—	—	—					Q.	—	—	M.fr.IX,185	...	...	As., P.	
C. latifolium ; F. v. M. in Benth. Fl. Austr. III, 421 (1866) ...	W.A.	S.A.	—	—	N.S.W.					Q.	—	B.fl.III,421	M.fr.IX,185				
C. attenuatum ; R. Brown in Benth. Fl. Austr. III, 421 (1866) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,421					
C. lucidum ; Hooker & Arnott, Bot. of Beechey's voy. 65 (1841) ...	—	—	—	—	N.S.W.					Q.	N.A.	B.fl.III,421	M.fr.II,133; IX,185; XI,135		P.		
C. oleifolium ; Hooker in Mitch. Trop. Austr. 397 (1848) ...	—	—	—	—	N.S.W.					Q.	—	B.fl.III,422	M.fr.II,134; IX,185.				
C. buxifolium ; Bentham, Fl. Austr. III, 422 (1866)...	—	—	—	—	N.S.W.					Q.	—	B.fl.III,422					
C. vacciniifolium ; F. v. M. in Transact. phil. Inst. Vict. III, 47 (1858) ...	—	—	—	—	N.S.W.					Q.	—	B.fl.III,422	M.fr.II,134; IX,186.				
C. coprosmoides ; F. v. M. in Transact. phil. Inst. Vict. III, 47 (1858) ...	—	—	—	—	N.S.W.					Q.	N.A.	B.fl.III,422					

MORINDA ; Vaillant in act. Acad. Par. 202 (1722).

M. citrifolia ; Linné, spec. plant. 176 (1753) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,422	..	...	...	As., Afr., P.	
M. acutifolia ; F. v. M., fragm. IX, 179 (1875) ...	—	—	—	—	—					Q.	—	B.fl.III,429	M.fr.IX,179.				
M. jasminoides ; Cunningham in Hook. Bot. Mag. t. 3351 (1834)...	—	—	—	V.	N.S.W.					Q.	—	B.fl.III,423					
M. umbellata ; Linné, spec. plant. 176 (1753)...	—	—	—	—	—					Q.	—	B.fl.III,423	...	...	...	As., P.	
M. reticulata ; Bentham, Fl. Austr. III, 424 (1866) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,423	M.fr.IX,180.				
M. hypotephra ; F. v. M. in Vict. Nat. III, 170 (1887) ...	—	—	—	—	—					Q.	—	—					

COELOSPERMUM ; Blume, Bijdr. 994 (1826). (Pogonolobus.)

C. paniculatum ; F. v. M., fragm. V, 19 (1865) ...	—	—	—	—	N.S.W.					Q.	—	B.fl.III,425	M.fr.VII,46; IX,185.				
C. reticulatum ; Bentham, Fl. Austr. III, 425 (1866) ...	—	—	—	—	—					Q.	N.A.	B.fl.III,425	M.fr.I,56; IX,185.				

HYDNOPHYTUM ; Jack in Transact. Linn. Soc. XIV, 124 (1823).

H. formicarum ; Jack in Transact. Linn. Soc. XIV, 124 (1823)	—	—	—	—	—	Q.	N.A.	—	M.fr.VII,45	..	...	As., P.

MYRMECODIA ; Jack in Transact. Linn. Soc. XIV, 122 (1823).

M. Antoinii ; Beccari, Malesia II, 116, t. 14 (1884)	—	—	—	—	—	Q.	—	—	M.fr.VII,45.			

LASIANTHUS ; Jack in Transact. Linn. Soc. XIV, 125 (1823).

L. strigosus ; Wight in Calc. Journ. nat. hist. VI, 512 (1846)	—	—	—	—	—	Q.	—	B.fl.III,426		...	...	...	As.

PSYCHOTRIA ; Linné, syst. ed. decim. 929 (1759). (Uragoga, 1737, Grumilia.)

P. nesophila ; F. v. M., fragm. II, 135 (1861)...	—	—	—	—	—	Q.	N.A.	B.fl.III,427	M.fr.IX,184.			
P. Dallachyana ; Bentham, Fl. Austr. III, 427 (1866)	—	—	—	—	—	Q.	—	B.fl.III,427				
P. nematopoda ; F. v. M., fragm. VII, 48 (1869)	—	—	—	—	—	Q.	—	—	M.fr.IX,184.			
P. loniceroides ; Sieber in De Candolle, prodr. IV, 523 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,427	M.fr.IX,184.			
P. daphnoides ; Cunningham in Hook. Bot. Mag. 3228 (1833)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,428	M.fr.II,135; IX,184.			
P. poliostemma ; Bentham, Fl. Austr. III, 427 (1866)	—	—	—	—	—	Q.	—	B.fl.III,428				
P. Fitzalani ; Bentham, Fl. Austr. III, 427 (1866)	—	—	—	—	—	Q.	—	B.fl.III,428				
P. Carronis ; Moore & F. v. M., fragm. VII, 49 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,70,185.			

GEOPHILA ; D. Don, prodr. fl. Nepal. 136 (1825). (Cephaelis partly.)

G. reniformis ; D. Don, prodr. fl. Nepal. 136 (1825)...	—	—	—	—	—	Q.	—	—	M.fr.VII,45; VIII,147; IX,187.			[Am., P. As., Afr.,

COPROSMA ; R. et G. Forster, char. gen. 137, t. 69 (1776). (Nertera, Marquisia.)

C. Nertera ; F. v. M., fragm. IX, 186 (1875)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,431	M.fr.IX,186	..	...	Am., P., N.Z.
C. reptans ; F. v. M., fragm. IX, 186 (1875)	—	—	—	V.	N.S.W.	—	—	B.fl.III,431	M.fr.VIII,147.			
C. pumila ; J. Hooker in Lond. Journ. VI, 465 (1847)	—	—	T.	V.	—	—	—	B.fl.III,430	M.fr.IX,186	...	...	N.Z.
C. nitida ; J. Hooker in Lond. Journ. VI, 465 (1847)	—	—	T.	V.	—	—	—	B.fl.III,430	M.fr.IX,186.			
C. Billardieri ; J. Hooker in Lond. Journ. VI, 465 (1847)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,430				
C. hirtella ; Labillardière, Nov. Holl. pl. spec. I, 70, t. 95 (1804)...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,429	M.fr.IX,186.			
C. Baueri ; Endlicher, iconogr. gen. pl. XI, t. 111 (1838)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,69.			
C. putida ; C. Moore & F. v. M., fragm. VII, 45 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,69.			
C. lanceolaris ; F. v. M., fragm. IX, 70 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,70.			
C. pilosa ; Endlicher, floral. ins. Norf. prodr. 60 (1833)	—	—	—	—	N.S.W.	—	—	—				

OPERCULARIA ; J. Gaertner, de fruct. I, 111, t. 24 (1788).

O. vaginata ; Labillardière, Nov. Holl. pl. spec. I, 34, t. 46 (1804)	W.A.	—	—	—	—	—	—	B.fl.III,433				
O. spermacocea ; Labillardière, Nov. Holl. pl. spec. I, 35, t. 47 (1804)	W.A.	—	—	—	—	—	—	B.fl.III,433				
O. scabrida ; Schlechtendal, Linnaea XX, 604 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,433				
O. hirsuta ; F. v. M. in Benth. Fl. Austr. III, 434 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,434				
O. aspera ; Gaertner, de fruct. I, 112, t. 24 (1788)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,434				
O. hispida ; Sprengel, syst. veg. I, 385 (1825)...	—	—	—	V.	N.S.W.	—	—	B.fl.III,434				
O. diphylla ; Gaertner, de fruct. I, 113 (1788)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,434				
O. ovata ; J. Hooker in Lond. Journ. VI, 465 (1847)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,435				
O. varia ; J. Hooker in Lond. Journ. VI, 466 (1847)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,435				
O. rubioides ; A. L. de Jussieu in Annal. du Mus. IV, 428 (1804)...	W.A.	—	—	—	—	—	—	B.fl.III,435				
O. volubilis ; R. Brown in Benth. Fl. Austr. III, 435 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,435				
O. hispidula ; Endlicher in Hueg. enum. 58 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,436				
O. echinocephala ; Bentham, Fl. Austr. III, 436 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,436				
O. apiciflora ; Labillardière, Nov. Holl. pl. spec. I, 35, t. 48 (1804)	W.A.	—	—	—	—	—	—	B.fl.III,436				

POMAX ; Solander in Gaertner, de fruct. I, 112 (1788).

P. umbellata ; Solander in Gaertner, de fruct. I, 112 (1788)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,437	M.fr.IX,187.			

ELEUTHRANTHES ; F. v. M., fragm. IV, 92 (1864).

E. opercularina ; F. v. M., fragm. IV, 92 (1864)	W.A.	—	—	—	—	—	—	B.fl.III,437	M.fr.IV,92.			

KNOXIA; Linné, Fl. Zeyl. 189 (1747).

K. corymbosa; Willdenow, spec. pl. I, 582 (1798)	—	—	—	—	—	Q.	N.A.	B.fl.III,438	M.fr.IV,41; IX,187	...	As., P.

SPERMACOCE; Dillenius, hort. Elth. 369, t. 277 (1732).

S. brachystoma; R. Brown in Benth. Fl. Austr. III, 439 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,439	M.fr.IV,41.		
S. pogostoma; Bentham, Fl. Austr. III, 440 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,440			
S. leptoloba; Bentham, Fl. Austr. III, 440 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,440			
S. marginata; Bentham, Fl. Austr. III, 440 (1866)	—	S.A.	—	—	—	Q.	N.A.	B.fl.III,440			
S. multicaulis; Bentham, Fl. Austr. III, 440 (1866)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,440			
S. exserta; Bentham, Fl. Austr. III, 441 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,441			
S. membranacea; R. Brown in Benth. Fl. Austr. III, 441 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,441			
S. debilis; Bentham, Fl. Austr. III, 441 (1866)	—	—	—	—	—	Q.	—	B.fl.III,441			
S. inaperta; F. v. M., fragm. IV, 43 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,441	M.fr.IV,43.		
S. stenophylla; F. v. M., fragm. IV, 43 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.III,442	M.fr.IV,43.		
S. laevigata; F. v. M., fragm. IV, 41 (1863)	—	—	—	—	—	Q.	N.A.	B.fl.III,442	M.fr.IV,41.		
S. breviflora; F. v. M. in Benth. Fl. Austr. III, 442 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,442			
S. auriculata; F. v. M., fragm. IV, 42 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,442	M.fr.IV,42.		
S. suffruticosa; R. Brown in Benth. Fl. Austr. III, 443 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,443			

ASPERULA; Dodoens, pemptad. 355 (1583) from Dalechamps (1554).

A. geminifolia; F. v. M., fragm. V, 147 (1865)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,443	M.fr.IX,188; XI,27.		
A. oligantha; F. v. M. in Neerl. Kruidk. Arch. IV, 111 et 112 (1859)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,444	M.fr.IX,187; XI,27.		

GALIUM; Dodoens, pemptad. 335 (1583) from Dioscorides.

G. umbrosum; Solander in G. Forster prodr. 89 (1786)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,446	M.fr.IX,188...	...	N.Z.
G. Australe; De Candolle, prodr. IV, 608 (1830)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,446			

CAPRIFOLIACEAE.

ADANSON, Familles des plantes II, 133 (1763).—Genus 1, Species 2.

SAMBUCUS; Tournefort, inst. 606, t. 396 (1700) from Dodoens (1583) and Plinius. (Tripetelus.)

S. xanthocarpa; F. v. M. in Transact. phil. Soc. Vict. I, 42 (1855)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,398			
S. Gaudichaudiana; De Candolle, prodr. IV, 322 (1830)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,398			

PASSIFLOREAE.

A. L. DE JUSSIEU in Ann. du Mus. VI, 102 (1805).—Genera 2, Species 6.

PASSIFLORA; Plukenet, phytograph. 202 et 282, t. 104, 210, 211, 212 (1692). (Disemma).

P. Herbertiana; Lindley, Bot. Regist. t. 737 (1823)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,311	M.fr.IX,68.		
P. cinnabarina; Lindley in Gard. Chron. 724 (1855)...	—	—	—	V.	N.S.W.	—	—	—	M.fr.IX,68.		
P. glabra; Wendland, collect. pl. I, 55, t. 17 (1808)...	—	—	—	—	N.S.W.	—	—	—			
P. aurantia; G. Forster, fl. ins. Austr. prodr. 62 (1786)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,312	M.fr.IX,68 ...	...	P.
P. brachystephanea; F. v. M. in Benth. Fl. Austr. III, 312 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,312	M.fr.I,56.		

MODECCA, Lamarck, Encycl. méthod. IV, 208 (1797).

M. Australis; R. Brown in De Candolle prodr. III, 337 (1828)	—	—	—	—	—	Q.	N.A.	B.fl.III,313	M.fr.IX,69.		

CUCURBITACEAE.

HALLER, enum. stirp. Helv. praef. 34 (1742).—Genera 10, Species 27.

TRICHOSANTHES; Linné, gen. plant. 295 (1737).

T. Holtzei; F. v. M. in Austr. Journ. Pharm. I, 447 (1886)	—	—	—	—	—	—	N.A.	—			
T. pentaphylla; F. v. M. in Benth. Fl. Aust. III, 314 (1866)	—	—	—	—	—	Q.	—	B.fl.III,314			

T. cucumerina; Linné, spec. plant. 1008 (1753) ... ... — — — — — Q. N.A. B.fl.III,314 ... ... As.
T. subvelutina; F. v. M. in A. & C. De Candolle, mon. phan. III, 366 (1881) ... — — — — N.S.W. Q. — B.fl.III,315 M.fr.VI,187.
T. Hearnii; F. v. M. in Benth. Fl. Austr. III, 315 (1866) ... ... — — — — — Q. — B.fl.III,315
T. Muelleri; Cogniaux in F. v. M., Cens. fourth suppl. 4 (1889) ... ... — — — — — — N.A. —

LAGENARIA; Séringe in Mém. Soc. Genèv. III, 25, t. 2 (1825).
L. vulgaris; Séringe in De Candolle, prodr. III, 279 (1828) ... ... — — — — — Q. — B.fl.III,316 M.fr.VII,62... ... As., Afr.

LUFFA; Tournefort in act. Acad. Paris. 84, t. 2 (1706), from Vesling (1638).
L. Aegyptiaca; Miller, Gardener's Dictionary (1731) ... ... — — — — — Q. N.A. B.fl.III,316 M.fr.III,107 ... As., Afr., P.
L. graveolens; Roxburgh, Fl. Ind. III, 716 (1832) ... ... — — — — — — N.A. B.fl.III,316 M.fr.III106;IV,175 ... As., P.
L. foetida; Cavanilles, icon. I, 7, t. 9—10 (1791) ... ... — — — — — — N.A. — M.fr.XI,138... ... As., Afr., P.

ZANONIA; Linné, coroll. 19 (1737). (Alsomitra.)
Z. Capricornica; F. v. M., fragm. VII, 61 (1870) ... ... — — — — — Q. — — M.fr.VII,61
Z. Hookeri; F. v. M., fragm. VI, 188 (1868) ... ... — — — — — Q. — — M.fr.V,181;VII,62.
Z. Stephensiana; F. v. M., fragm. VIII, 181 (1874)... ... ... — — — — — Q. — — M.fr.VIII,181.

CUCUMIS; Tournefort, inst. 104, t. 31 (1700) from Plinius.
C. trigonus; Roxburgh, Fl. Ind. III, 722 (1832) ... ... W.A. — — — N.S.W. Q. N.A. B.fl.III,317 M.fr.VI,187... ... As.
C. Chate; Linné, syst. nat. edit. decim. 5 (1759) ... ... W.A. S.A. — — N.S.W. Q. N.A. — ... ... As., Afr.

BENINCASA; Savi in Bibl. Ital. IX, 158 (1818).
B. vacua; F. v. M., Cens. of Austr. pl. 76 (1882) ... ... — — — — — Q. — — M.fr.VI,186.

MOMORDICA; Tournefort, institut. rei herbar. 103, t. 29 et 30 (1700).
M. Balsamina; Linné, sp. pl. 1009 (1753) ... ... — — — — N.S.W. Q. — B.fl.III,318 M.fr.VI,187... .. As., Afr.
M. Charantia; Linné, sp. pl. 1009 (1753) ... ... — S.A. — — N.S.W. Q. N.A. — ... ... As., Afr.

BRYONOPSIS; Arnott in Hook. Journ. of Bot. III, 274 (1841). (Bryonia partly.)
B. laciniosa; Naudin in Annal. des sc. nat. sér. cinq. VI, 30 (1869) ... ... — — — — N.S.W. Q. N.A. B.fl.III,319 ... ... As., Afr.
B. Pancheri; Naudin in Annal. des sc. nat. sér. cinq. V, 30 (1869) ... ... — — — — N.S.W. — — — ... ... P.

MELOTHRIA; Linné, coroll. 1 (1737). (Zehneria, Mukia.)
M. Cunninghamii; Bentham, Fl. Austr. III, 320 (1866) ... ... — — — — N.S.W. Q. N.A. B.fl.III,320 M.fr.VI,188;VII,62.
M. Baueriana; F. v. M., fragm. VI, 188 (1866) ... ... — — — — N.S.W. — — — M.fr.VI,188.
M. Muelleri; Bentham, Fl. Austr. III, 320 (1866) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.III,320 M.fr.I,182; II,180; III,107.
M. Maderaspatana; Cogniaux in De Candolle mon. phan. III, 623 (1881) ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.III,321 M.fr.VI,187 ... ... As., Afr., P.
M. subpellucida; Cogniaux in F. v. M., Cens. fourth suppl. 4 (1889) ... ... — — — — — Q. N.A. —
M. Celebica; Cogniaux in De Candolle, mon. phan. III, 625 (1881) ... ... — — — — — Q. N.A. — ... ... As.

SICYOS; Linné, syst. nat. 9 (1735); gen. plant. 297 (1737). (Sycios.)
S. angulata; Linné, spec. plant. 1013 (1753) ... ... — — T. V. N.S.W. Q. — B.fl.III,322 M.fr.VI,187; VII,62 ... Afr., Am., N.Z., P.

COMPOSITAE.

VAILLANT in act. Acad. Paris. 143 (1718).—Genera 90, Species 539.

ETHULIA; Linné, fil. decas. prima plant. rar. hort. Upsal. 1, t. 1 (1762).
E. conyzoides; Linné, fil. dec. prim. 1, t. 1 (1762) ... ... — — — — N.S.W. Q. — — .. .. As., Afr.

CENTRATHERUM; Cassini in Bull. de la Soc. philom. 31 (1817).
C. muticum; Lessing in Schlecht. Linnaea II, 320 (1829) ... ... — — — — N.S.W. Q. — B.fl.III,460 ... ... Am., P.

PLEUROCARPAEA; Bentham, Fl. Austr. III, 460 (1866).
P. denticulata; Bentham, Fl. Austr. III, 460 (1866) ... ... — — — — — Q. N.A. B.fl.III,460

VERNONIA ; Schreber, gen. plant. II, 541 (1791).

V. cinerea ; Lessing in Schlecht. Linnaea II, 291 (1829)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.III,459	...	...	...	As., Afr., P.
V. vagans ; De Candolle, prodr. V, 32 (1843)	—	—	—	—	—	Q.	—	—	M.fr.VI,234		...	As.

ELEPHANTOPUS ; Vaillant in act. Acad. Par. 309 (1719).

E. scaber ; Linné, spec. plant. 814 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.III,461	...	...	...	As., Afr.

ADENOSTEMMA ; R. et G. Forster, char. gen. 89, t. 45 (1776).

A. viscosum ; R. & G. Forster, char. gen. 89, t. 45 (1776)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,462	...	...	...	As., Afr., Am., P

AGERATUM ; Linné, gen. plant. 247 (1737).

A. conyzoides ; Linné, spec. plant. 839 (1753)	—	—	—	—	—	Q.	—	B.fl.III,462	...	...	...	As., Afr., Am., P.

EUPATORIUM ; Tournefort, inst. 455, t. 259 (1700), from C. Bauhin (1623), from Gerard (1597).

E. cannabinum ; Linné, spec. plant. 838 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,462	M.fr.V,62	...	...	E., As., Afr.

LAGENOPHORA ; Cassini in Bull. de la Soc. philom. 34 (1818). (Ixauchenus.)

L. Billardieri ; Cassini in Diction. XXV, 111 (1822)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,507	...	...	...	As.
L. Huegelii ; Bentham in Hueg. enum. 59 (1837)	W.A.	S.A.	T.	V.	—	—	—	B.fl.III,507				

SOLENOGYNE ; Cassini in Diction. des sc. nat. V, 174 (1828). (Emphysopus.)

S. bellioides ; Cassini in Dict. V, 174 (1828)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,508	M.fr.V,62.					
S. Emphysopus ; F. v. M., Key to syst. Vict. pl. I, 310 (1888)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,508	M.fr.V,62.					

BRACHYCOME ; Cassini in Bull. de la Soc. philom. 199 (1816). (Bellidis subgenus, Pacquerina, Brachystephium, Steiroglossa, Silphiospermum.)

B. diversifolia ; Fischer & Meyer, ind. sem. hort. Petrop. 31 (1835)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,511						
B. segmentosa ; Moore & F. v. M., fragm. VIII, 144 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,144.					
B. melanocarpa ; Sonder & F. v. M. in Schl. Linn. XXV, 476 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,511						
B. radicans ; Steetz in Lehm. pl. Preiss. I, 429 (1845)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,512						
B. goniocarpa ; Sonder & F. v. M. in Schl. Linn. XXV, 474 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,512						
B. pachyptera ; Turczaninow in Bull. de Mosc. XXIV, 175 (1851)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,512						
B. iberidifolia ; Bentham in Hueg. enum. 59 (1837)	W.A.	—	—	—	—	—	N.A.	B.fl.III,512	M.fr.XI,123.					
B. pusilla ; Steetz in Lehm. pl. Preiss. I, 427 (1845)	W.A.	—	—	—	—	—	—	B.fl.III,513						
B. microcarpa ; F. v. M., fragm. I, 50 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,513	M.fr.I,50.					
B. Stuartii ; Bentham, Fl. Austr. III, 513 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,513						
B. scapigera ; De Candolle, prodr. VII, 277 (1838)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,513						
B. Muelleri ; Sonder in Schlecht. Linnaea XXV, 475 (1852)	—	S.A.	—	—	—	—	—	B.fl.III,514						
B. graminea ; F. v. M., first gen. Rep. 14 (1853)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,514	M.fr.I,49; VIII,142.					
B. angustifolia ; Cunningham in De Candolle, prodr. V, 306 (1836)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,514	M.fr.I,49.					
B. linearifolia ; De Candolle, prodr. V, 306 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,515						
B. basaltica ; F. v. M., fragm. I, 50 (1858)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,515	M.fr.I,50.					
B. trachycarpa ; F. v. M. in Schlecht. Linnaea XXV, 339 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,515						
B. exilis ; Sonder in Linnaea XXV, 473 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,516						
B. ptychocarpa ; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.III,516						
B. debilis ; Sonder in Linnaea XXV, 477 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,516						
B. cuneifolia ; Tate in Transact. R. S. of S. Austr. XI, 83 (1888)	—	S.A.	—	—	—	—	—	—						
B. decipiens ; J. Hooker in Lond. Journ. VI, 114 (1847)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,516						
B. cardiocarpa ; F. v. M. in Benth. Fl. Austr. III, 516 (1866)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,517						
B. nivalis ; F. v. M. in Transact. phil. Soc. Vict. I, 43 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.III,517						
B. scapiformis ; De Candolle, prodr. V, 306 (1836)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,517						
B. stricta ; De Candolle, prodr. V, 305 (1836)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,518						
B. heterodonta ; De Candolle, prodr. V, 305 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,518						
B. Billardieri ; Bentham, Fl. Austr. III, 518 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,518						

B. cheilocarpa ; F. v. M. in Wing'a South. Sc. Record, Aug. (1882)	W.A.	—	—	—	—	—	—	—		
B. ciliaris ; Lessing, syn. comp. 172 (1832)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,518		
B. calocarpa ; F. v. M. in Schlecht. Linnaea XXV, 399 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,519	M.fr.XI,27.	
B. chrysoglossa ; F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854)	—	S.A.	—	V.	N.S.W.	—	—	—		
B. marginata ; Bentham in Hueg. enum. 60 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,519		
B. Sieberi ; De Candolle, prodr. V, 306 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,520		
B. discolor ; C. Stuart in Benth. Fl. Austr. III, 520 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,520		
B. multifida ; De Candolle, prodr. V, 306 (1836)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,520	M.fr.VIII,145.	
B. latisquamea ; F. v. M., fragm. XI, 16 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,16.	
B. glandulosa ; Bentham, Fl. Austr. III, 520 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,520		
B. collina ; Bentham, Fl. Austr. III, 521 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,521		
ERODIOPHYLLUM ; F. v. M., fragm. IX, 119 (1875). (Subgenus of Erigeron, sect. Trimorphaea.)										
E. Elderi ; F. v. M., fragm. IX, 120 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,119.	
MINURIA ; De Candolle, prodr. V, 298 (1836). (Therogeron, Elachothamnua, Kippistia.)										
M. leptophylla ; De Candolle, prodr. V, 298 (1836)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,498	M.fr.IX,119; X,56.	
M. Cunninghamii ; Bentham, Fl. Austr. III, 498 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,498		
M. Candollei ; F. v. M., fragm. IX, 119 (1875)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,499	M.fr.V,87; X,56.	
M. suaedifolia ; F. v. M. in Bentham, Fl. Austr. III, 499 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,499	M.fr.X,56.	
ACHNOPHORA ; F. v. M. in Proceed. Roy. Soc. South Austr. VI, 32 (1883).										
A. Tatei ; F. v. M. in Proc. Roy. Soc. S. Austr. VI, 33 (1883)	—	S.A.	—	—	—	—	—	—		
CALOTIS ; R. Brown in Bot. Regist. t. 504 (1820). (Huenefeldia, Goniopogon, Cheiroloma.)										
C. dentex ; R. Brown in Bot. Regiat. 504 (1820)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,501		
C. cuneifolia ; R. Brown in Bot. Regist. 504 (1820)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,501		
C. glandulosa ; F. v. M. in Transact. Vict. Inst. 129 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.III,502		
C. cymbacantha ; F. v. M. in Schlecht. Linnaea XXV, 400 (1852)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,502		
C. erinacea ; Steetz in Lehm. pl. Preiss. I, 424 (1845)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,502		
C. acabiosifolia ; Sonder & F. v. M. in Linnaea XXV, 471 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,503	M.fr.XI,27.	
C. acapigera ; Hooker in Mitch. Trop. Austr. 75 (1848)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,503	M.fr.XI,27.	
C. anthemoides ; F. v. M. in Transact. phil. soc. Vict. I, 44 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.III,504		
C. lappulacea ; Bentham in Hueg. enum. 60 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,504		
C. microcephala ; Bentham, Fl. Austr. III, 504 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,504		
C. breviseta ; Bentham in Hueg. enum. 60 (1837)	—	—	—	—	—	Q.	N.A.	B.fl.III,505		
C. plumulifera ; F. v. M. in Transact. phil. Inst. Vict. III, 57 (1859)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,505		
C. porphyroglossa ; F. v. M. in Benth. Fl. Austr. III, 505 (1866)	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.III,505		
C. pteroaperma ; R. Brown in Benth. Fl. Austr. III, 505 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,505		
C. hispidula ; F. v. M. in Transact. Vict. Inat. 130 (1855)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,506		
C. Kempei ; F. v. M. in Transact. Roy. Soc. S. Austr. IV, 112 (1881)	—	S.A.	—	—	—	—	—	—	M.fr.XI,139.	
ASTER ; Tournefort, inst. 481, t. 274 (1700), from Dioscorides. (Olcaria, Celmisia, Eurybia, Steetzia, Diplopappus, Diplostephium.)										
A. megalophyllus ; F. v. M., fragm. V, 70 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.III,468	M.fr.V,70.	
A. chrysophyllus ; Cunningham in De Candolle, prodr. V, 266 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,468	M.fr.II,88; V,71.	
A. alpicola ; F. v. M., fragm V, 70 (1865)	—	—	—	V.	N.S.W.	—	—	B.fl.III,468	M.fr.V,70.	
A. rosmarinifolius ; Cunningham in De Candolle, prodr. V, 268 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,468		
A. viscosus ; Labillardière, Nov. Holl. pl. spec. II, 53, t. 203 (1806)	—	—	T.	V.	—	—	—	B.fl.III,468	M.fr.V,71.	
A. Sonderi ; F. v. M., fragm. V, 83 (1865)	—	S.A.	—	—	—	—	—	B.fl.III,469	M.fr.V,83.	
A. pannosus ; F. v. M., fragm. V, 83 (1865)	—	S.A.	—	V.	—	—	—	B.fl.III,469	M.fr.V,83.	
A. oliganthemus ; F. v. M., Cens. of Austr. pl. 78 (1882)	—	—	—	—	N.S.W.	—	—	B.fl.III,469		
A. argophyllus ; Labillardière, Nov. Holl. pl. spec. II, 52, t. 201 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,470	M.fr.V,68.	
A. cydonifolius ; Cunningham in De Candolle, prodr. V, 267 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.III,470		

A. myrsinoides ; Labillardière, Nov. Holl. pl. spec. II, 53, t. 202 (1806)...	..	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,471 M.fr.V,69.
A. persoonioides ; Cunningham in De Candolle, prodr. V, 267 (1836) ...	...	—	—	T.	—	—	—	—	B.fl.III,471 M.fr.V,69.
A. obcordatus ; F. v. M., fragm. V, 69 (1865)	...	—	—	T.	—	—	—	—	B.fl.III,471 M.fr.V,69.
A. pinifolius ; F. v. M., fragm. V, 71 (1865)	...	—	—	T.	—	—	—	—	B.fl.III,472 M.fr.V,71.
A. ledifolius ; Cunningham in De Candolle, prodr. V, 269 (1836)... ...	...	—	—	T.	—	—	—	—	B.fl.III,472
A. dentatus ; Andrews, Bot. Reposit. t. 61 (1799)	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,472 M.fr.V,82.
A. stellulatus ; Labillardière, Nov. Holl. pl. spec. II, 50, t. 196 (1806) ...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,473 M.fr.V,81.
A. asterotrichus ; F. v. M., fragm. V, 79 (1865)	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,473 M.fr.I,111; II,180.
A. gravis ; F. v. M., fragm. V, 82 (1865)	...	—	—	—	—	N.S.W.	—	—	B.fl.III,474 M.fr.V,82.
A. Nernstii ; F. v. M., fragm. V, 81 (1865)	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,474 M.fr.V,81.
A. hygrophilus ; Cunningham in De Candolle, prodr. V, 269 (1836) ...	...	—	—	—	—	—	Q.	—	B.fl.III,474
A. Siemssenii ; F. v. M., fragm. V, 71 (1865)...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,475 M.fr.I,50.
A. tubuliflorus ; F. v. M., fragm. V, 65 (1865)	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,475 M.fr.V,65.
A. axillaris ; F. v. M., fragm. V, 64 (1865)	...	W.A.	S.A.	T.	V.	N.S.W.	—	N.A.	B.fl.III,475 M.fr.V,64.
A. revolutus ; F. v. M., Cens. of Austr. pl. 78 (1882)	...	W.A.	—	—	—	—	—	—	B.fl.III,476
A. exilifolius ; F. v. M., fragm. V, 69 (1865)...	...	W.A.	—	—	—	—	—	—	B.fl.III,476 M.fr.V,69.
A. ramulosus ; Labillardière, Nov. Holl. plant. spec. II. 51, t. 198 (1806)	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,476
A. florulentus ; F. v. M., fragm. V, 70 (1865)	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,477 M.fr.V,82.
A. microphyllus ; Ventenat, Jard. de Malmais. 83 (1803)	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,478 M.fr.V,70.
A. lepidophyllus ; Persoon, synop. plant. II, 442 (1807)	...	—	S.A.	T.	V.	N.S.W.	—	—	—
A. Mitchelli ; F. v. M., fragm. V, 78 (1865)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,478 M.fr.V,78.
A. exiguifolius ; F. v. M., fragm. V, 67 (1865)	...	W.A.	S.A.	—	—	—	—	—	B.fl.III,478 M.fr.V,67.
A. Cassiniae ; F. v. M., fragm. V, 68 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,479 M.fr.V,68.
A. cyanodiscus ; F. v. M., fragm. V, 82 (1865)	...	—	—	—	—	N.S.W.	—	—	B.fl.III,479 M.fr.V,82.
A. pimeleoides ; Cunningham in De Candolle, prodr. V, 268 (1836) ...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,479
A. iodochrus ; F. v. M., fragm. V, 81 (1865)...	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,480 M.fr.II,110.
A. adenolasius ; F. v. M., fragm. V, 67 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,480 M.fr.V,67.
A. calcareus ; F. v. M., Cens. of Austr. pl. 78 (1882)	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481
A. magniflorus ; F. v. M., fragm. V, 80 (1865)	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,481 M.fr.V,80.
A. Muelleri ; Cens. 78 (1882), from Sonder (1852)	...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481
A. Stuartii ; F. v. M., fragm. V, 76 (1865)	...	—	S.A.	—	—	—	—	—	B.fl.III,481 M.fr.I,202.
A. decurrens ; Cunningham in De Candolle, prodr. V, 269 (1836) ...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,481
A. glutescens ; F. v. M., fragm. V, 77 (1865)	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,482 M.fr.V,77.
A. orarius ; F. v. M., fragm. V, 78 (1865)	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,482 M.fr.V,78.
A. vernicosus ; F. v. M., fragm. V, 67 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,482 M.fr.V,67.
A. teretifolius ; F. v. M., fragm. V, 77 (1865)	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,483 M.fr.V,77.
A. Turczaninowii ; F. v. M., fragm. V, 67 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,483 M.fr.V,67.
A. ellipticus ; Cunningham in De Candolle, prodr. V, 271 (1836)	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,483 M.fr.I,16,242.
A. glandulosus ; Labillardière, Nov. Holl. pl. spec. II, 50, t. 197 (1806)...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,484
A. heleophilus ; F. v. M., fragm. V, 66 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,484 M.fr.V,66.
A. muricatus ; F. v. M., fragm. V, 66 (1865)...	...	W.A.	—	—	—	—	—	—	B.fl.III,484 M.fr.V,66.
A. Benthami ; F. v. M., Cens. of Austr. pl. 78 (1882)	...	—	—	—	V.	—	—	—	B.fl.III,485
A. Steetzii ; F. v. M., fragm. V, 66 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,485 M.fr.V,66.
A. Cunninghami ; F. v. M., Cens. 78 (1882), from De Candolle (1836) ..	...	—	—	—	—	N.S.W.	—	—	B.fl.III,486
A. paucidentatus ; F. v. M., fragm. V, 66 (1865)	...	W.A.	—	—	—	—	—	—	B.fl.III,485 M.fr.V,66.
A. megalodontus ; F. v. M., fragm. VIII, 244 (1874)	...	W.A.	S.A.	—	—	—	—	—	— M.fr.VIII,244.
A. adenophorus ; F. v. M., fragm. V, 78 (1865)	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,486 M.fr.I,111.
A. homolepis ; F. v. M., fragm. V, 65 (1865)...	...	W.A.	—	—	—	—	—	—	B.fl.III,486 M.fr.V,65.
A. xerophilus ; F. v. M., fragm. V, 76 (1865)	...	—	—	—	—	—	Q.	—	B.fl.III,486 M.fr.I,51,76&86.
A. Ferresii ; F. v. M., fragm. V, 75 (1865)	...	—	S.A.	—	—	—	—	N.A.	B.fl.III,487 M.fr.III,18.

A. exul; Lindley, Bot. Regist. XXV, App. 24 (1839)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,487	M.fr.I,51; V,75.			
A. argutus; R. Brown in Benth. Fl. Austr. III, 488 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,488				
A. Huegelii; F. v. M., fragm. V, 79 (1865)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,488	M.fr.V,79.			
A. Balli; F. v. M., fragm. VIII, 143 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,143.			
A. Mooneyi; F. v. M., fragm. VIII, 144 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,144.			
A. Celmisia; F. v. M., fragm. V, 84 (1865)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,489	M.fr.V,84	...	N.Z.	
VITTADINIA; Ach. Richard, Voy. de l'Astrol. bot. 250 (1832). (Eurybiopsis, Microgyne.)												
V. brachycomoides; F. v. M., fragm. V, 86 (1865)	—	—	—	—	—	Q.	N.A.	B.fl.III,490	M.fr.V,86.			
V. Australis; A. Richard, Voy. de l'Astrol. 251 (1832)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,490	M.fr.V,87; XI,20		N.Z.	
V. scabra; De Candolle, prodr. V, 281 (1836)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,491				
V. macrorrhiza; A. Gray in Proc. Am. Acad. V, 118 (1863)	—	—	—	—	—	—	N.A.	B.fl.III,492				
PODOCOMA; Cassini in Bull. de la Soc. philom. 137 (1817). (Ixiochlamys.)												
P. cuneifolia; R. Brown, App. to Sturt, Centr. Austr. 17 (1849)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,493				
ERIGERON; Linné, hort. Cliffort. 407 (1737). (Haplopappus partly.)												
E. pappochromus; Labillardière, Nov. Holl. pl. spec. II, 47, t. 193 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,494	M.fr.V,87.			
E. ambiguus; F. v. M. in Transact. phil. Inst. Vict. III, 58 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.III,494				
E. sessilifolius; F. v. M., fragm. X, 100 (1881)	—	—	—	—	—	—	N.A.	—	M.fr.X,137.			
E. minuriioides; Bentham, Fl. Aust. III, 495 (1866)	—	—	—	V.	—	—	—	B.fl.III,495				
E. conyzoides; F. v. M. in Transact. phil. Soc. Vict. I, 105 (1855)	—	—	—	V.	N.S.W.	—	—	B.fl.III,495	M.fr.V,87.			
CONYZA; Lamarck, Encycl. méthod. II, 87, 89 (1786).												
C. viscidula; Wallich, numer. list 3006 (1828)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,496	...	...	As., P.	
C. Aegyptiaca; Aiton, hort. Kew. III, 183 (1789)	—	—	—	—	—	Q.	N.A.	B.fl.III,497	...	...	As., Afr.	
BLUMEA; De Candolle in Guillemin, Arch. de Bot. II, 514 (1833).												
B. glandulosa; De Candolle in Wight, contrib. 14 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.III,525	..	...	As., Afr.	
B. lacera; De Candolle in Wight, contrib. 14 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.III,526	...	...	As., Afr.	
B. amplectens; De Candolle in Wight, contrib. 13 (1834)	—	—	—	—	—	Q.	—	B.fl.III,525	...	...	As.	
B. integrifolia; De Candolle, prodr. V, 433 (1836)	—	—	—	—	—	Q.	N.A.	B.fl.III,525				
B. diffusa; R. Brown in Benth. Fl. Austr. III, 525 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,525				
B. hieracifolia; De Candolle in Wight, contrib. 15 (1834)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,526	...	...	As.	
B. Cunninghamii; De Candolle, prodr. V, 435 (1836)	—	—	—	—	—	Q.	N.A.	B.fl.III,526				
B. acutata; De Candolle in Ann. du Mus. d'hist. nat. Par. III, 409 (1834)	—	—	—	—	—	Q.	—	—	M.fr.IX,160.	...	As.	
PLUCHEA; Cassini in Bull. de la Soc. philom. 31 (1817). (Spiropodium, Eyrea.)												
P. Indica; Lessing in Schlecht. Linnaea VI, 150 (1831)	—	—	—	—	—	Q.	N.A.	B.fl.III,527	M.fr.X,101	...	As., P.	
P. Dioscoridis; De Candolle, prodr. V, 450 (1836)	—	—	—	—	—	Q.	—	—	...	...	As., Afr.	
P. tetranthera; F. v. M. Rep. Babb. Exped. 12 (1860)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,528				
P. conocephala; F. v. M. in Transact. Roy. Soc. Vict. 24, 138 (1887)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,480	M.fr.V,79.			
P. baccharoides; F. v. M. in Benth. Fl. Austr. III, 528 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,528	M.fr.I,34.			
P. Eyrea; F. v. M. Rep. Babb. Exped. 11, 12 (1860)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,528				
P. squarrosa; Bentham, Fl. Austr. III, 529 (1866)	W.A.	—	—	—	—	—	N.A.	B.fl.III,529				
P. dentex; R. Brown in Benth. Fl. Austr. III, 529 (1866)	—	—	—	—	—	Q.	—	B.fl.III,529				
PTERIGERON; De Candolle, prodr. V, 293 (1836). (Streptoglossa, Oliganthemum.)												
P. decurrens; De Candolle, prodr. V, 293 (1836)	W.A.	—	—	—	—	—	N.A.	B.fl.III,531				
P. liatroides; Bentham, Fl. Austr. III, 532 (1866)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,532				
P. odorus; Bentham, Fl. Austr. III, 532 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,532				
P. macrocephalus; Bentham, Fl. Austr. 532 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,532				
P. microglossus; Bentham, Fl. Austr. III, 532 (1866)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,532				

P. dentatifolius; F. v. M., fragm. IX, 119 (1875) ... ... — S.A. — — — — — — M.fr.X,119.
P. adscendens; Bentham Fl. Austr. III, 533 (1866) ... ... — S.A. — — — Q. N.A. B.fl.III,533
P. filifolius; Bentham, Fl. Austr. III, 533 (1866) ... ... — — — — — — N.A. B.fl.III,533

THESPIDIUM; F. v. M. in Journ. of Landsborough s Exped. app. (1862).
T. basiflorum; F. v. M. in Journ. of Landsb. Exped. App. (1862) ... ... — — — — — Q. N.A. B.fl.III,534 M.fr.XI,101.

COLEOCOMA; F. v. M. in Hook. Kew Misc. IX, 19 (1857).
C. Centaurea; F. v. M. in Hook. Kew Misc. IX, 19 (1857) ... ... — — — — — — N.A. B.fl.III,533 M.fr.X,101.

EPALTES; Cassini in Bull. de la Soc. philom. 139 (1818). (Sphaeromorphaea partly, Ethuliopsis, Gymnaphanes.)
E. Cunninghamii; Bentham, Fl. Austr. III, 530 (1866) ... ... — S.A. — V. N.S.W. Q. — B.fl.III,530 M.fr.II,155.
E. Australis; Lessing in Linnaea V, 148 (1831) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.III,530 M.fr.XI,101 ... As.
E. pleiochaeta; F. v. M., fragm. X, 100 (1877) ... ... — — — — N.S.W. — — — M.fr.X,100.
E. Harrisii; F. v. M., fragm. XI, 101 (1881) ... ... — — — — — Q. — — M.fr.XI,101.
E. Tatei; F. v. M. in Transact. Roy. Soc. of S.A. VI, 31 (1883) ... ... W.A. S.A. — — — — — —

SPHAERANTHOS; Vaillant in Act. Acad. Par. 289 (1719). (Sphaeranthus.)
S. Indicus; Linné, sp. pl., 927 (1753) ... ... — — — — — Q. N.A. B.fl.III,521 M.fr.III,138 ... As., Afr.
S. Africanus; Linné, sp. pl. ed. sec. 13, 14 (1762) ... ... — — — — — — N.A. B.fl.III,522 M.fr.III,138 ... As., Afr.

PTEROCAULON; Elliot, Sketch of the Bot. of S. Carolina and Georgia II, 323 (1824). (Monenteles.)
P. verbascifolius; Bentham & J. Hooker, gen. pl. II, 294 (1873) ... ... — — — — — Q. N.A. B.fl.III,523
P. Billardieri; F. v. M., Papuan plants, 43 (1876) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.III,523 M.fr.XI,101 ... As., P.
P. sphacelatus; Bentham & J. Hooker, gen. plant. II, 94 (1873) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.III,523 .. .. P.
P. glandulosus; Bentham & J. Hooker, gen. plant. II, 94 (1873) ... ... — — — — — Q. N.A. B.fl.III,523
P. sphaeranthoides; Bentham & J. Hooker, gen. plant. II, 94 (1873) ... ... — — — — — — N.A. B.fl.III,524

STUARTINA; Sonder in Schlechtendal's Linnaea XXV, 521 (1852).
S. Mnelleri; Sonder in Schlecht. Linnaea XXV. 522 (1852) ... ... — S.A. — V. N.S.W. Q. — B.fl.III,657 M.fr.VIII,145; XI,27.

GNAPHALIUM; Linné, gen. plant. 250 (1737), from J. & C. Bauhin (1619). (Euchiton.)
G. luteo-album; Linné, spec. plant. 851 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.III,653 M.fr.V,150 E., As., Afr., Am., P., N.Z.
G. Japonicum; Thunberg, Fl. Japon. 311 (1784) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.III,653 M.fr.V,150; XI,27. As., N.Z.
G. alpigenum; F. v. M., second gen. Report 12 (1854) ... ... — — T. V. — — — B.fl.III,654 M.fr.V,150.
G. purpureum; Linné, spec. plant. 854 (1753) ... ... — — — — N.S.W. Q. — B.fl III,655... ... ... Am.
G. Indicum; Linné, spec. plant. 852 (1753) ... . — S.A. — — N.S.W. Q. N.A. B.fl.III,655 M.fr.V,149 ... As., Afr., P.
G. indutum; J. Hooker in Lond. Journ. of Bot. VI, 121 (1847) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.III,655 M.fr.V,150.
G. Traversii; J. Hooker, Handb. N. Zeal. Fl. 154 ... ... — — — V. N.S.W. — — B.fl.III,656... ... ... N.Z.

ANTENNARIA; Gaertner, de fruct. II, 410, t. 167 (1791). (Raoulia partly.)
A. Planchoni; F. v. M., Cens. 80 (1882), from J. Hook. (1860) ... ... — — T. — — — — B.fl.III,651
A. uniceps; F. v. M. in Transact. phil. Soc. Vict. t. 105 (1854) ... ... — — — V. N.S.W. — — B.fl.III,652 M.fr.VIII,145.

LEONTOPODIUM; R. Brown in Transact. Linn. Soc. XII, 124 (1817). (Raoulia partly.)
L. Catipes; F. v. M. in Papers of the Roy. Soc. of Tasm. 44 (1882) ... ... — — T. V. N.S.W. — — B.fl.III,651
L. Meredithae; F. v. M., Cens. of Austr. pl. 80 (1882) ... ... — — T. — — — — —

PTERYGOPAPPUS; J. Hooker in Lond. Journ. of Bot. VII, 120 (1847).
P. Lawrencii; J. Hooker in Lond. Journ. VII, 120 (1847) ... ... — — T. — — — — B.fl.III,656

PODOSPERMA; Labillardière, Nov. Holl. plant. specim. II, 35, t. 177 (1806). (Podospermum, Podotheca, Phaenopoda, Lophoclinium.)
P. gnaphalioides; F. v. M., fragm. XII, 22 (1882) from Graham (1842) ... ... W.A. — — — — — — B.fl.III,601 M.fr.XII,21.
P. angustifolium; Labillardière, Nov. Holl. pl. specim. II, 35, t.177 (1806) ... W.A. S.A. T. V. N.S.W. — — B.fl.III,601 M.fr.XI,20; XII,21.
P. pygmaeum; F. v. M., fragm. XII, 22 (1882) implied (1852) ... ... W.A. — — — — — — B.fl.III,602

P. chrysantha; F. v. M., fragm. XII, 22 (1882)	W.A.	—	—	—	—	—	—				B.fl.III,602	M.fr.XII,22.
P. Polakii; F. v. M., fragm. XII, 21 (1882)	W.A.	—	—	—	—	—	—				—	M.fr.XII,21.
IXIOLAENA; Bentham in Hueg. enum. 66 (1837).												
I. brevicompta; F. v. M., fragm. I, 53 (1858)	—	—	—	—	N.S.W.	Q.	—				B.fl.III,597	M.fr.I,53.
I. leptolepis; Bentham, Fl. Austr. III, 597 (1866)	—	S.A.	—	V.	N.S.W.	Q.	N.A.				B.fl.III,597	
I. supina; F. v. M. in Transact. Vict. Inst. 37 (1855)	—	S.A.	T.	—	—	—	—				B.fl.III,598	
I. tomentosa; Sonder & F. v. M. in Linnaea XXV, 504 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,598	M.fr.I,53; XI,27.
I. viscosa; Bentham in Hueg. enum. 66 (1837)	W.A.	—	—	—	—	—	—				B.fl.III,598	
PODOLEPIS; Labillardière, Nov. Holl. pl. specim. II, 57, t. 28 (1806). (Scalia, Panaetia, Scaliopsis, Siemssenia, Stylolepis, Rutidochlamys.)												
P. rhytidochlamys; F. v. M., fragm. IV, 79 (1864)	—	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,603	M.fr.IV,79.
P. longipedata; Cunningham in De Candolle, prodr. VI, 163 (1837)	—	—	—	V.	N.S.W.	Q.	—				B.fl.III,604	M.fr.I,112.
P. acuminata; R. Brown in Ait. hort. Kew. ed. sec. V, 82 (1813)	—	S.A.	T.	V.	N.S.W.	Q.	—				B.fl.III,604	M.fr.XI,27.
P. canescens; Cunningham in De Candolle, prodr. VI, 163 (1837)	—	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,605	
P. aristata; Bentham in Hueg. enum. 64 (1837)	W.A.	—	—	—	—	—	—				B.fl.III,605	
P. pallida; Turczaninow in Bull. de Mosc. XXIV, 78 (1851)	W.A.	—	—	—	—	—	N.A.				B.fl.III,605	
P. nutans; Steetz in Lehm. pl. Preiss. I, 464 (1845)	W.A.	—	—	—	—	—	—				B.fl.III,605	
P. gracilis; Graham in Edinb. New phil. journ. V, 379 (1828)	W.A.	—	—	—	—	—	—				B.fl.III,606	
P. rugata; Labillardière, Nov. Holl. pl. spec. II, 57, t. 208 (1806)	W.A.	S.A.	—	V.	—	—	N.A.				B.fl.III,606	
P. Lessoni; Bentham, Fl. Austr. III, 606 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,606	M.fr.XI,27.
P. Siemssenii; F. v. M., fragm. V, 200 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,607	M.fr.V,200.
P. microcephala; Bentham, Fl. Austr. 607 (1866)	W.A.	—	—	—	—	—	—				B.fl.III,607	
P. Kendalli; F. v. M. in Wing's S. Sc. Rec. III, 68 (1883)	W.A.	—	—	—	—	—	—				—	M.fr.VIII,168; X,108.
ATHRIXIA; Ker in Bot. Regist. VIII, 681 (1823). (Asteridea, Chrysodiscus, Trichostegia.)												
A. Australis; Steetz in Lehm. pl. Preiss. I, 482 (1845)	W.A.	—	—	—	—	—	—				B.fl.III,599	
A. gracilis; Bentham, Fl. Austr. III, 599 (1866)	W.A.	—	—	—	—	—	—				B.fl.III,599	
A. multiceps; Bentham, Fl. Austr. III, 599 (1866)	W.A.	—	—	—	—	—	—				B.fl.III,599	
A. stricta; Bentham, Fl. Austr. III, 600 (1866)	W.A.	—	—	—	—	—	—				B.fl.III,600	
A. tenella; Bentham, Fl. Austr. III, 600 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—				B.fl.III,600	
A. Croniniana; F. v. M. in Vict. Nat. V, 54 (1888)	W.A.	—	—	—	—	—	—				—	
A. chaetopoda; F. v. M., fragm. X, 56 (1876)	W.A.	—	—	—	—	—	—				—	M.fr.X,56.
LEPTORRHYNCHOS; Lessing, syn. composit. 273 (1832). (Rhytidanthe, Doratolepis, Waitziae subgenus.)												
L. squamatus; Lessing, syn. comp. 273 (1832)	—	S.A.	T.	V.	N.S.W.	Q.	—				B.fl.III,608	M.fr.I,52.
L. panaetioides; Bentham, Fl. Austr. III, 609 (1866)	—	S.A.	—	V.	N.S.W.	—	—				B.fl.III,609	
L. tenuifolius; F. v. M., fragm. I, 52 (1858)	—	S.A.	—	V.	—	—	—				B.fl.III,609	M.fr.I,52.
L. pulchellus; F. v. M. in Linnaea XXV, 500 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,610	M.fr.I,53; XI,27.
L. Baileyi; F. v. M., fragm. X, 101 (1877)	—	—	—	—	—	Q.	—				—	M.fr.X,101.
L. elongatus; De Candolle, prodr. VI, 160 (1837)	W.A.	S.A.	T.	V.	N.S.W.	—	—				B.fl.III,610	M.fr.XI,27,28.
L. medius; Cunningham in De Candolle, prodr. VI, 160 (1837)	W.A.	S.A.	—	V.	N.S.W.	—	—				—	M.fr.XI,85.
L. Waitzia; Sonder in Schlecht. Linnaea XXV, 501 (1852)	—	S.A.	—	V.	N.S.W.	—	—				B.fl.III,610	
L. nitidulus; De Candolle, prodr. VI, 160 (1837)	—	—	T.	V.	N.S.W.	—	—				B.fl.III,611	
WAITZIA; Wendland, Collect. pl. II, 13, t. 42 (1808). (Viraya, Morna, Pterochaete.)												
W. corymbosa; Wendland, Coll. pl. II, 13, t. 42 (1808)	W.A.	S.A.	—	V.	N.S.W.	Q.	—				B.fl.III,635	
W. aurea; Steetz in Lehm. pl. Preiss. I, 452 (1845)	W.A.	—	—	—	—	—	—				B.fl.III,636	
W. nivea; Bentham, Fl. Austr. III, 636 (1866)	W.A.	—	—	—	—	—	—				B.fl.III,636	
W. Steetziana; Lehmann, pl. Preiss. I, 454 (1845)	W.A.	—	—	—	—	—	—				B.fl.III,636	
W. podolepis; Steetz in Lehm. pl. Preiss. I, 450 (1845)	W.A.	—	—	—	—	—	—				B.fl.III,637	
W. paniculata; F. v. M. in Benth. Fl. Austr. III, 637 (1866)	W.A.	—	—	—	—	—	—				B.fl.III,637	

HELIPTERUM ; De Candolle, prodr. VI, 211 (1837). (Argyrocome partly, Pteropogon, Rodanthe, Xyridanthe, Anisolepis, Hyalosperma, Triptilodiscus, Acroclinium, Monencyanthes, Dimorpholepis, Duttonia, Cassiniola.)

H. Manglesii ; F. v. M., fragm. V, 200 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,640	M.fr.X,108.
H. Margaritae ; F. v. M., fragm. XI, 48 (1878)	—	—	—	—	—	—	N.A.	—	M.fr.XI,48.
H. roseum ; Bentham, Fl. Austr. III, 640 (1886)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.III,640	M.fr.X,108.
H. fuscescens ; Turczaninow in Bull. de Mosc., 80 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,602	M.fr.XII,22.
H. rubellum ; Bentham, Fl. Austr. III, 641 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,641	M.fr.X,108.
H. chlorocephalum ; Bentham, Fl. Austr. III, 641 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,641	
H. anthemoides ; De Candolle, prodr. VI, 216 (1837)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,641	M.fr.X,108.
H. polygalifolium ; De Candolle, prodr. VI, 216 (1837)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,641	M.fr.X,107,108.
H. floribundum ; De Candolle, prodr. VI, 217 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,642	M.fr.X,108.
H. Pyrethrum ; Bentham, Fl. Austr. III, 642 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,642	M.fr.X,110.
H. heteranthum ; Turczaninow in Bull. de Mosc. XXIV, 198 (1851)	W.A.	S.A.	—	—	—	—	—	B.fl.III,642	M.fr.III,137; X,108.
H. stipitatum ; F. v. M. in Bentham, Fl. Austr. III, 643 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,643	M.fr.III,133; X,109.
H. incanum ; De Candolle, prodr. VI, 215 (1837)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,643	
H. Cotula ; De Candolle, prodr. VI, 215 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,644	M.fr.II,157; III,134; X,110.
H. hyalospermum ; F. v. M. in Benth. Fl. Austr. III, 644 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,644	M.fr.X,109.
H. condensatum ; F. v. M., fragm. III, 136 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.III,645	M.fr.III,136.
H. polyphyllum ; F. v. M., fragm. I, 35 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,645	M.fr.I,35.
H. Humboldtianum ; De Candolle, prodr. VI, 216 (1866)	W.A.	—	—	—	—	—	N.A.	B.fl.III,645	M.fr.III,135; X,109.
H. Haigii ; F. v. M., fragm. X, 107 (1877)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,107.
H. tenellum ; Turczaninow in Bull. de Mosc. XXIV, 198 (1851)	W.A.	S.A.	—	—	—	—	—	B.fl.III,646	M.fr.III,135; X,109; XI,20.
H. strictum ; Bentham, Fl. Aust. III, 646 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,646	M.fr.X,109.
H. corymbiflorum ; Schlechtendal, Linnaea XXI, 448 (1848)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,647	M.fr.X,109.
H. sterilescens ; F. v. M. in Wing's S. Sc. Rec. II, 273 (1882)	W.A.	—	—	—	—	—	—	—	
H. pygmaeum ; Bentham, Fl. Austr. III, 647 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,647	M.fr.X,109.
H. spicatum ; F. v. M. in Bentham, Fl. Austr. III, 647 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,647	M.fr.X,109.
H. Charsleyae ; F. v. M., fragm. VIII, 168 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.XI,134.
H. moschatum ; Bentham, Fl. Austr. III, 648 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,648	M.fr.X,107.
H. Tietkensii ; F. v. M., fragm. VIII, 227 (1874)	—	S.A.	—	—	N.S.W.	—	—	—	M.fr.VIII,227.
H. pterochaetum ; Bentham, Fl. Austr. III, 648 (1866)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,648	M.fr.X,109.
H. Frenchii ; F. v. M. in Wing's S. Sc. Rec. III, 2 (1883)	W.A.	—	—	—	—	—	—	—	
H. Forrestii ; F. v. M. in Wing's S. Sc. Rec. II, 273 (1882)	W.A.	—	—	—	—	—	—	—	
H. polycephalum ; Bentham, Fl. Austr. III, 649 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,649	M.fr.III,139; X,109.
H. corymbosum ; Bentham, Fl. Austr. III, 649 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,649	M.fr.V,200.
H. laeve ; Bentham, Fl. Austr. III, 649 (1866)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.III,649	M.fr.V,200.
H. exiguum ; F. v. M. in Transact. Vict. Inst. 39 (1855)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,644	M.fr.X,109.
H. Dimorpholepis ; Bentham, Fl. Austr. III, 650 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,650	M.fr.X,109; XI,27.

HELICHRYSUM ; Vaillant in Act. Acad. Par. 290 (1719), from Theophrastos and Dioscorides. (Elichrysum, Schoenia, Petalolepis, Faustula, Ozothamnus, Swammerdamia, Lawrencella, Argyrophanes, Chrysocephalum, Conanthodium, Xanthochrysum, Argyroglottis, Acanthocladium, Raoulia partly.)

H. Cassinianum ; Gaudichaud in Freye. Voy. Bot. 466, t. 87 (1826)	W.A.	S.A.	—	—	—	—	—	B.fl.III,616	
H. Lawrencella ; F. v. M. in Benth. Fl. Austr. III, 616 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.III,616	
H. subnlifolium ; F. v. M., fragm. III, 134 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,616	M.fr.III,134.
H. filifolium ; F. v. M., fragm. III, 134 (1863)	W.A.	—	—	—	—	—	—	B.fl.III,617	M.fr.III,134.
H. MacIvorii ; F. v. M. in Wing's S. Sc. Rec. III, 99 (1883)	W.A.	—	—	—	—	—	—	—	
H. semifertile ; F. v. M., Rep. Babb. Exped. 14 (1858)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,617	
H. Baxteri ; Cunningham in De Candolle, prodr. VI, 193 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,617	
H. Ayersii ; F. v. M., fragm. VIII, 167 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.IX,195.
H. rutidolepis ; De Candolle, prodr. VI, 194 (1837)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,618	
H. scorpioides ; Labillardière, Nov. Holl. pl. spec. II, 45, t. 191 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,618	M.fr.VIII,167.

H. oligochaetum; F. v. M., fragm. VI, 235 (1868)	W.A.	—	—	—	—	—	N.A.	—	M.fr.VI,235.
H. Tepperi; F. v. M. in Wing's South. Science Record II, 1 (1882)	—	S.A.	—	—	—	—	—	—	
H. obtusifolium; Sonder & F. v. M. in Linnaea XXV, 515 (1852)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,619	
H. Calvertianum; F. v. M., fragm. X, 108 (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,136.
H. Spiceri; F. v. M., fragm. XI, 47 (1878)	—	—	T.	—	—	—	—	—	M.fr.XI,47.
H. dealbatum; Labillardière, Nov. Holl. pl. spec. II, 45, t. 190 (1806)	—	—	T.	V.	—	—	—	B.fl.III,619	
H. pumilum; J. Hooker, Fl. Tasman. I, 213, t. 60 (1860)	—	—	T.	—	—	—	—	B.fl.III,619	
H. lucidum; Henckel, adumbr. pl. hort. Hal. 5 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,620	M.fr.XI,48.
H. elatum; Cunningham in De Candolle, prodr. VI, 193 (1837)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,621	
H. adenophorum; F. v. M. in Transact. Vict. Inst. 38 (1855)	—	S.A.	—	V.	—	—	—	B.fl.III,622	
H. leucopsidium; De Candolle, prodr. VI, 193 (1837)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,622	M.fr.VI,255.
H. Blandowskianum; Steetz in Schlecht. Linnaea XXV, 512 (1852)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,622	
H. oxylepis; F. v. M., fragm. I, 35 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,623	M.fr.I,35.
H. collinum; De Candolle, prodr. VI, 190 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,623	
H. rupicola; De Candolle, prodr. VI, 190 (1837)	—	—	—	—	—	Q.	—	B.fl.III,623	
H. podolepideum; F. v. M., Rep. Babb. Exped. 13 (1858)...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,624	M.fr.III,134.
H. Gilesii; F. v. M., fragm. X, 85 (1876)	W.A.	—	—	—	—	—	N.A.	—	M.fr.X,86.
H. ambiguum; Turczaninow in Bull. de Mosc. 195 (1851)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,609	
H. apiculatum; De Candolle, prodr. VI, 195 (1837)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,624	
H. semipapposum; De Candolle, prodr. VI, 195 (1837)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,625	
H. Dockerii; F. v. M. in Benth. Fl. Austr. III, 626 (1866)...	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,626	M.fr.II,156.
H. Agyroglottis; Bentham Fl. Austr. III, 626 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,626	
H. ramosum; De Candolle, prodr. VI, 181 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,626	
H. cordatum; De Candolle, prodr. VI, 180 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,627	
H. obovatum; De Candolle, prodr. VI, 180 (1837)	—	—	—	—	N.S.W.	—	—	B.fl.III,627	M.fr.II,89.
H. Bidwillii; Bentham, Fl. Austr. III, 627 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,627	
H. Becklerii; F. v. M. in Benth. Fl. Austr. III, 627 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,627	M.fr.I,183,244.
H. diotophyllum; F. v. M., frsgm. V, 150 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,628	M.fr.VIII,46.
H. Thomsoni; F. v. M., fragm. VIII, 45 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,45.
H. diosmifolium; Don in Steud. Nom. Bot. ed. sec. 738 (1841)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,628	
H. retusum; F. v. M., fragm. VIII, 46 (1873)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,628	M.fr.VIII,46.
H. decurrens; F. v. M., fragm. VIII, 46 (1873)	—	S.A.	—	V.	N.S.W.	—	—	—	M.fr.VIII,46.
H. Cunninghamii; Bentham, Fl. Austr. III, 629 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,629	
H. reticulatum; Lessing in Steud. Nom. Bot. ed. sec. 740 (1841)...	—	—	T.	—	—	—	—	B.fl.III,629	
H. cinereum; F. v. M. in Benth. Fl. Austr. III, 629 (1866)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,629	M.fr.V,199.
H. bracteolatum; Bentham, Fl. Austr. III, 630 (1866)	—	—	T.	—	—	—	—	B.fl.III,630	
H. Kempei; F. v. M. in Melb. Chemist, Jan. (1882)	—	S.A.	—	—	—	—	—	—	
H. cassinioides; Bentham, Fl. Austr. III, 630 (1866)	—	—	—	—	—	Q.	—	B.fl.III,630	
H. Gunnii; F. v. M. in Benth. Fl. Austr. III, 630 (1866)	—	—	T.	—	—	—	—	B.fl.III,630	M.fr.V,199.
H. rosmarinifolium; Lessing, synops. gen. comp. 274 (1832)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,631	
H. ledifolium; F. v. M. in Benth. Fl. Austr. III, 630 (1866)	—	T.	—	—	—	—	—	B.fl.III,630	M.fr.V,199.
H. ferrugineum; Lessing, synops. gen. comp. 274 (1832)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,631	
H. antennarium; F. v. M. in Benth. Fl. Austr. III, 632 (1866)	—	—	T.	—	—	—	—	B.fl.III,632	M.fr.V,199.
H. obcordatum; F. v. M. in Benth. Fl. Austr. III, 632 (1866)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,632	M.fr.V,199.
H. Backhousii; F. v. M. in Benth. Fl. Austr. III, 632 (1866)	—	—	T.	V.	—	—	—	B.fl.III,632	M.fr.V,199.
H. cuneifolium; F. v. M. in Benth. Fl. Austr. III, 633 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,633	
H. baccharoides; F. v. M. in Benth. Fl. Austr. III, 633 (1866)	—	—	T.	V.	N.S.W.	—	—	B.fl.III,633	M.fr.V,199.
H. lepidophyllum; F. v. M. in Benth. Fl. Austr. III, 633 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,633	M.fr.V,199.
H. scutellifolium; Bentham, Fl. Austr. III, 633 (1866)	—	—	T.	—	—	—	—	B.fl.III,633	M.fr.V,199.
H. lycopodioides; Bentham, Fl. Austr. III, 634 (1866)	—	—	T.	—	—	—	—	B.fl.III,634	M.fr.V,199.
H. selaginoides; F. v. M. in Benth. Fl. Austr. III, 634 (1866)	—	—	T.	—	—	—	—	B.fl.III,634	M.fr.V,199.

CASSINIA ; R. Brown in Transact. Linn. Soc. XII, 126 (1817). (Non zoologorum, Apolochlamys, Achromolaena, Chromochiton.)

C. leptocephala ; F. v. M., fragm. III, 138 (1863)	—	—	—	—	N.S.W.	—	—	B.fl.III,585	M.fr.III,138.
C. compacta ; F. v. M., fragm. I, 18 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,585	M.fr.I,18.
C. denticulata ; R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	—	—	N.S.W.	—	—	B.fl.III,586	
C. longifolia ; R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,586	
C. aurea ; R. Brown in Transact. Linn. Soc. XII, 127 (1817)	—	—	—	—	N.S.W.	—	—	B.fl.III,586	
C. aculeata ; R. Brown in Transact. Linn. Soc. XII, 127	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,586	
C. laevis ; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,587	
C. tenuifolia ; Bentham, Fl. Austr. III, 585 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,585	
C. quinquefaria ; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,587	
C. arcuata ; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,587	
C. subtropica ; F. v. M., fragm. I, 17 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,588	M.fr.I,17.
C. Theodori ; F. v. M., fragm. V, 148 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,588	M.fr.V,148.
C. spectabilis ; R. Brown in Transact. Linn. Soc. XII, 128 (1817)	—	S.A.	T.	V.	—	—	—	B.fl.III,588	

HUMEA ; Smith, Exot. bot. I, t. 1 (1804). (Haeckeria, Calomeria.)

H. elegans ; Smith, Exot. bot. I, t. 1 (1804)	—	—	—	V.	N.S.W.	—	—	B.fl.III,589	M.fr.I,17.
H. cassiniacea ; F. v. M., fragm. I, 17 (1858)...	—	S.A.	—	—	—	—	—	B.fl.III,589	M.fr.I,17.
H. ozothamnoides ; F. v. M., fragm. I, 17 (1858)	—	—	—	V.	N.S.W.	—	—	B.fl.III,590	M.fr.VIII,142.
H. squamata ; F. v. M., fragm. XI, 86 (1880)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,634	M.fr.II,131; V,190.

ACOMIS ; F. v. M., fragm. II, 89 (1860).

A. Rutidosis ; F. v. M., fragm. II, 89 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,591	M.fr.II,89.
A. macra ; F. v. M., fragm. IV, 145 (1864)	—	—	—	—	—	Q.	—	B.fl.III,591	M.fr.IV,145.

RUTIDOSIS ; De Candolle, prodr. VI, 158 (1837). (Pumilo, Actinopappus, Lepidocoma.)

R. leiolepis ; F. v. M. in Transact. Vict. Inst. 131 (1855)	—	—	—	V.	—	—	—	B.fl.III,593	M.fr.I,34.
R. leptorrhynchoides ; F. v. M., fragm. V, 148 (1866)	—	—	—	V.	N.S.W.	—	—	B.fl.III,593	M.fr.V,148.
R. helichrysoides ; De Candolle, prodr. VI, 159 (1837)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,594	
R. leucantha ; F. v. M., fragm. I, 35 (1858)	—	—	—	—	—	Q.	—	B.fl.III,594	M.fr.I,35.
R. Murchisonii ; F. v. M., fragm. I, 34 (1858)	—	—	—	—	—	Q.	—	B.fl.III,594	M.fr.I,35.
R. Pumilo ; Bentham, Fl. Austr. III, 595 (1866)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,595	

PITHOCARPA ; Lindley, Bot. Regist. XXV, App. 23 (1839).

P. corymbosa ; Lindley, Bot. Regist. XXV, App. 23 (1839)	W.A.	—	—	—	—	—	—	B.fl.III,590	

AMMOBIUM ; R. Brown in Bot. Magaz. t. 2459 (1824).

A. alatum ; R. Brown in Bot. Magaz. t. 2459 (1824)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,583	
A. craspedioides ; Bentham, Fl. Austr. III, 584 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.III,584	

IXODIA ; R. Brown in Ait. hort. Kew. sec. ed. IV, 17 (1812).

I. achilleoides ; R. Brown in Ait. hort. Kew, sec. ed. IV, 17 (1812)	—	S.A.	—	V.	—	—	—	B.fl.III,583	

MILLOTIA ; Cassini in Ann. sc. nat. XVII, 416 (1829).

M. tenuifolia ; Cassini in Ann. sc. nat. XVII, 416 (1829)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,596	
M. Greevesii ; F. v. M., fragm. III, 18, t. 19 (1863)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,596	M.fr.III,18.
M. Kempei ; F. v. M. in Wing's South. Sc. Record II, 1 (1882)	—	S.A.	—	—	—	—	—	—	

TOXANTHUS ; Turczaninow in Bull. de Mosc. XXIV, 177 (1851). (Anthocerastes, Scyphocoronis.)

T. perpusillus ; Turczaninow in Bull. Mosc. XXIV, 177 (1851)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,592	M.fr.XI,49.
T. Muelleri ; Bentham, Fl. Austr. III, 592 (1866)	W.A.	S.A.	—	V.	—	—	—	B.fl.III,592	M.fr.XI,49.
T. major ; Turczaninow in Bull. de Mosc. XXIV, 64 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,592	M.fr.XI,49.

QUINETIA; Cassini in Diction. sc. nat. LX, 579 (1830).

Q. Urvillei; Cassini in Diction. LX, 579 (1830)	W.A.	S.A.	—	—	—	—	—	B.fl.III,595	

PHACELOTHRIX; F. v. M., fragm. XI, 49 (1878).

P. cladochaeta; F. v. M., fragm. XI, 49 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.III,594	M.fr.V,199.

DECAZESIA; F. v. M., fragm. XI, 71 (1879).

D. hecatocephala; F. v. M., fragm. XI, 72 (1879)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,72.

MYRIOCEPHALUS; Bentham in Hueg. enum. 61 (1837). (Hyalolepis, Antheidosorus, Gilberta, Lamprochlaena, Elachopappus, Polycalymma; some readily to be restituted.)

M. rhizocephalus; Bentham, Fl. Austr. III, 557 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,557	M.fr.III,155.
M. nudus; A. Gray in Hook. Kew Misc. III, 174 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,558	M.fr.III,157.
M. appendiculatus; Bentham in Hueg. enum. 61 (1837)	W.A.	—	—	—	—	—	—	B.fl.III,558	
M. Rudallii; F. v. M., fragm. III, 157 (1863)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.III,559	M.fr.VI,235; VII,145.
M. gracilis; Bentham, Fl. Austr. III, 559 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,559	M.fr.VIII,169.
M. helichrysoides; A. Gray in Hook. Kew Misc. III, 175 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,559	M.fr.III,159.
M. suffruticosus; Bentham, Fl. Austr. III, 559 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,559	
M. Stuartii; Bentham, Fl. Austr. III, 560 (1866)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,560	
M. Guerinae; F. v. M., fragm. VIII, 169 (1874)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,169.

ANGIANTHUS; Wendland, Collect. pl. II, 31 (1809). (Siloxerus, Styloncerus, Ogcerostylus, Cylindorosorus, Phyllocalymma, Skirrophorus, Chrysocoryne, Eriocladium, Pogonolepis, Piptostemma, Epitriche, Gamozygis, Cephalosorus partly, Hyalochlamys, Dithyrostegia, Pleuropappus; some readily to be restituted.)

A. humifusus; Bentham, Fl. Austr. III, 563 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,563	
A. tomentosus; Wendland, Collect. pl. II, 31, t. 48 (1809)	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.III,562	
A. pleuropappus; Bentham, Fl. Austr. III, 563 (1866)	—	S.A.	—	—	—	—	—	B.fl.III,563	
A. brachypappus; F. v. M. in Transact. phil. Soc. Vict. I, 44 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,563	
A. myosuroides; Bentham, Fl. Austr. III, 563 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,563	
A. tenellus; Bentham, Fl. Austr. III, 564 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.III,564	
A. pusillus; Bentham, Fl. Austr. III, 564 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,564	
A. Milnei; Bentham, Fl. Austr. III, 564 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,564	
A. Cunninghamii; Bentham, Fl. Austr. III, 565 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,565	
A. phyllocephalus; Bentham, Fl. Austr. III, 565 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,565	M.fr.III,159.
A. micropoides; Bentham, Fl. Austr. III, 565 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,565	
A. microcephalus; Bentham, Fl. Austr. III, 566 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,566	M.fr.III,158.
A. platycephalus; Bentham, Fl. Austr. III, 566 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,566	
A. Drummondii; Bentham, Fl. Austr. III, 566 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,566	
A. Preissianus; Bentham, Fl. Austr. III, 566 (1866)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,566	
A. pygmaeus; Bentham, Fl. Austr. III, 567 (1868)	W.A.	—	—	—	—	—	—	B.fl.III,567	
A. globifer; Bentham, Fl. Austr. 567 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,567	
A. demissus; Bentham, Fl. Austr. III, 567 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,567	
A. strictus; Bentham, Fl. Austr. III, 568 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,568	
A. plumiger; Bentham, Fl. Austr. III, 568 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,568	
A. amplexicaulis; Bentham, Fl. Austr. III, 568 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,568	M.fr.III,159.

GNEPHOSIS; Cassini in Bull. de la Soc. philom. 45 (1820). (Cephalosorus partly, Nematopus, Crossolepis, Leptotriche, Trichanthodium, Cyathopappus; some readily to be restituted.)

G. Burkittii; Bentham, Fl. Austr. III, 570 (1866)	—	S.A.	—	—	N.S.W.	—	—	B.fl.III,570	
G. eriocarpa; Bentham, Fl. Austr. III, 570 (1866)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,570	M.fr.III,156.
G. macrocephala; Turczaninow in Bull. Soc. Mosc. XXIV, 190 (1851)	W.A.	—	—	—	—	—	—	B.fl.III,570	M.fr.II,158.
G. skirrophora; Bentham, Fl. Austr. III, 570 (1866)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.III,570	

G. cyathopappa ; Bentham, Fl. Austr. III, 571 (1866)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,571	M.fr.II,158.
G. codonopappa ; F. v. M. in Giles, geogr. trav. 217 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,2.
G. leptoclada ; Bentham, Fl. Austr. III, 571 (1866)...	W.A.	—	—	—	—	—	—	B.fl.III,571	M.fr.III,158.
G. arachnoidea ; Turczaninow in Bull. Soc. Mosc. XXIV, 189 (1851)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,571	M.fr.II,158.
G. tenuissima ; Cassini, in Bull. de la Soc. philom. 43 (1820)	W.A.	—	—	—	—	—	—	B.fl.III,572	
G. acicularis ; Bentham, Fl. Austr. III, 572 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,572	
G. pygmaea ; Bentham, Fl. Austr. III, 572 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,572	
G. brevifolia ; Bentham, Fl. Austr. III, 572 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,572	
G. eriocephala ; Bentham, Fl. Austr. III, 573 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,573	
ERIOCHLAMYS ; Sonder & F. v. M. in Schlecht. Linnaea XXV, 488 (1852).									
E. Behrii ; Sonder & F. v. M. in Linnaea XXV, 488 (1852)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,591	
E. Knappii ; F. v. M. in Melb. Chem. VI, 61, p. 4 (1883)	W.A.	S.A.	—	—	—	—	N.A.	—	
CALOCEPHALUS ; R. Brown in Transact. Linn. Soc. XII, 106 (1817). (Leucophyta, Pachysura, Blennospora, Achrysum.)									
C. Drummondii ; Bentham, Fl. Austr. III, 574 (1866)	W.A.	S.A.	—	—	—	—	—	B.fl.III,574	
C. Brownii ; F. v. M., Rep. Babb. Exped. 13 (1858)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,574	
C. Sonderi ; F. v. M., sec. gen. Report 12 (1854)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,775	
C. lacteus ; Lessing, syn. gen. comp. 271 (1832)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,575	
C. citreus ; Lessing, syn. gen. comp. 271 (1832)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,575	
C. angianthoides ; Bentham, Fl. Austr. III, 575 (1866)	W.A.	—	T.	—	—	—	—	B.fl.III,575	
C. Francisii ; Bentham, Fl. Austr. III, 576 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,576	M.fr.III,155.
C. Dittrichii ; F. v. M. in Uhlworm's bot. Centralbl. XXVII, 300 (1886) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	—	
C. platycephalus ; Bentham, Fl. Austr. 115, 576 (1866)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,576	M.fr.III,154.
C. multiflorus ; Bentham, Fl. Austr. III, 576 (1866)	W.A.	—	—	—	—	—	N.A.	B.fl.III,576	
C. aeruoides ; Bentham, Fl. Austr. III, 576 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,576	M.fr.III,154.
CEPHALIPTERUM ; A. Gray in Hook. Kew Misc. IV, 271 (1852).									
C. Drummondii ; A. Gray in Hook. Kew Misc. IV, 272 (1852)	W.A.	S.A.	—	—	—	—	—	B.fl.III,577	
GNAPHALODES ; A. Gray in Hook. Kew Misc. IV, 228 (1852).									
G. uliginosum ; A. Gray in Hook. Kew Misc. IV, 228 (1852)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,578	M.fr.XI,27.
G. condensatum ; A. Gray in Hook. Kew Misc. IV, 228 (1852)	W.A.	—	—	—	—	—	—	B.fl.III,578	
G. filifolium ; Bentham, Fl. Austr. III, 578 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,578	
CRASPEDIA ; G. Forster, florul. insul. Austr. prodr. 306 (1786). (Richea, Pycnosorus.)									
C. Richea ; Cassini, in Diction. XI, 353 (1818)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,579	... ... N.Z.
C. pleiocephala ; F. v. M. in Schlecht. Linnaea XXV, 404 (1852)...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580	M.fr.XI,27.
C. chrysantha ; Bentham, Fl. Austr. III, 580 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580	
C. globosa ; Bentham, Fl. Austr. III, 580 (1866)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,580	
CHTHONOCEPHALUS ; Steetz in Lehm. pl. Preiss. I, 444 (1845). (Chamaesphaerion, Gyrostephium, Lachnothalamus.)									
C. tomentellus ; Bentham, Fl. Austr. III, 581 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,581	M.fr.III,156; IV,176.
C. Pseudevax ; Steetz in Lehm. pl. Preiss. I, 445 (1845)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,582	
C. pygmaeus ; Bentham, Fl. Austr. III, 582 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,582	
NABLONIUM ; Cassini in Diction. XXXIV, 101 (1825).									
N. calyceroides ; Cassini, Dict. XXXIV, 101 (1825)...	—	—	T.	—	—	—	—	B.fl.III,545	
CHRYSOGONUM ; Linné, hort. Cliffort. 424 (1737). (Moonia, Pentalepis).									
C. trichodesmoides ; Bentham & J. Hooker, gen. pl. II, 350 (1873)	—	—	—	—	—	—	N.A.	B.fl.III,540	M.fr.VIII,145.
C. ecliptoides ; Bentham & J. Hooker, gen. pl. II, 350 (1873)	—	—	—	—	—	—	N.A.	B.fl.III,540	M.fr.VIII,145.
C. procumbens ; Bentham & J. Hooker, gen. pl. II, 350 (1873)	—	—	—	—	—	—	N.A.	B.fl.III,540	M.fr.VIII,145.

SIEGESBECKIA ; Linné, hort. Cliffort. 412 (1737).

S. orientalis ; Linné, spec. plant. 900 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,535	...	...	...	As., Afr., Am., P.

ENHYDRA ; Loureiro, Flor. Cochinch. II, 510 (1790). (Enydra.)

E. fluctuans ; Loureiro, fl. Cochin. II, 511 (1799)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,546	M.fr.III,139		...	As., Afr., Am.

ECLIPTA ; Linné, mantissa altera, 157 (1771). (Blainvillea.)

E. alba ; Hasskarl, plant. Javan. rar. 528 (1848)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,536	...	...	...	As., Afr., Am.
E. latifolia ; Linné, fil. suppl. 378 (1781)	—	—	—	—	—	Q.	N.A.	—	M.fr.VIII,145		...	As., Afr., Am.
E. platyglossa ; F. v. M., fragm. II, 135 (1861)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,536	M.fr.XI,27.			

WEDELIA ; Jacquin, stirp. Amer. hist. 217, t. 130 (1763). (Wollastonia.)

W. calendulacea ; Lessing, synops. gen. comp. 222 (1832)	—	—	—	—	—	Q.	—	B.fl.III,537	...	...	...	As.
W. urticifolia ; De Candolle in Wight, Contrib. Bot. Ind. 18 (1834)	—	—	—	—	—	Q.	N.A.	B.fl.III,538	...	...	...	As.
W. spilanthoides ; F. v. M., fragm. V, 64 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,538	M.fr.V,64.			
W. verbesinoides ; F. v. M. in Benth. Fl. Austr. III, 538 (1866)	—	S.A.	—	—	—	—	N.A.	B.fl.III,538				
W. biflora ; De Candolle in Wight, Contrib. Bot. Ind. 18 (1834)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,539	...	...	...	As., Afr., P.
W. asperrima ; Bentham, Fl. Austr. III, 539 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.III,539				

SPILANTHES ; Jacquin, stirp. Amer. hist. 214, t. 126 (1763).

S. grandiflora; Turczaninow in Bull. de Mosc. XXIV, 185 (1851)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,541	M.fr.V,63.			
S. anactina ; F. v. M., fragm. V, 63 (1865)	—	—	—	—	—	Q.	N.A.	B.fl.III,541	M.fr.V,63...		...	As.

BIDENS ; Dillenius, hort. Eltham. 51 et 52, t. 43 et 44 (1732), from Cesalpini (1583). (Possibly immigrated.)

B. tripartitus ; Linné, spec. plant. 831 (1753)...	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,543	...	...	...	E., As.
B. pilosus ; Linné, spec. plant. 832 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,543	...	...	...	As., Afr., Am., P.
B. bipinnatus ; Linné, spec. plant. 832 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,543	...	...	...	As., Afr., Am.

GLOSSOGYNE ; Cassini in Diction. LI, 475 (1827). (Diodontium.)

G. tenuifolia ; Cassini in Dict. LI, 475 (1827)...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.III,544	M.fr.I,51 ...		...	As., P.
G. retroflexa ; F. v. M., fragm. I, 51 (1858)	—	—	—	—	—	Q.	—	B.fl.III,544				
G. filifolia ; F. v. M. in Benth. Fl. Austr. III, 544 (1866)	—	—	—	—	—	—	N.A.	B.fl.III,544				

FLAVERIA ; A. L. de Jussieu, gen plant. 186 (1789).

F. Australasica ; Hooker in Mitch. Trop. Austr. 118 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.III,546	M.fr.I,183.			

SOLIVA ; Ruiz et Pavon, prodr. 113, t. 24 (1794). (Gymnostyles ; possibly immigrated.)

S. anthemifolia ; R. Brown in Transact. Linn. Soc. XII, 102 (1817)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,552	...	...	...	Am.

COTULA ; Linné, syst. nat. 9 (1735); Linné, gen. pl. 256 (1737). (Gymnogyne, Strongylospermum, Pleiogyne, Symphyomera, Ctenosperma, Leptinella.)

C. filifolia ; Thunberg, prodr. pl. Capens. 161 (1800)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,548	M.fr.VI,114		...	Afr.
C. coronopifolia ; Linné, sp. pl. 892 (1753)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,549	M.fr.VI,114		...	E., Afr., Am., N.Z.
C. Gymnogyne ; F. v. M. in Benth. Fl. Austr. III, 549 (1866)	W.A.	—	—	—	—	—	—	B.fl.III.549				
C. Australis ; J. Hooker, Fl. N. Zeal. I, 128 (1853)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,550	...	...	...	N.Z.
C. alpina ; J. Hooker, Fl. Tasman. I, 192, t. 51 (1860)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,550				
C. Drummondii ; Bentham, Fl. Austr. III, 550 (1866)	W.A.	—	—	—	—	—	—	B.fl.III,550				
C. reptans ; Bentham, Fl. Austr. III, 551 (1866)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,551				
C. filicula ; J. Hooker in Benth. Fl. Austr. III, 551 (1866)...	—	—	T.	V.	N.S.W.	—	—	B,fl.III,551				

CENTIPEDA ; Loureiro, Fl. Cochinch. II, 492 (1790). (Myriogyne, Sphaeromorphaea partly.)

C. racemosa ; F. v. M., Cens. 84 (1882) from Hook. (1848)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.III,553				
C. orbicularis ; Loureiro, Fl. Cochinch. II, 492 (1790)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,553	M.fr.VIII,142		...	As., Am., P., N.Z.
C. Cunninghami ; F. v. M., fragm. VIII, 143 (1874)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	—				
C. thespidioides ; F. v. M., fragm. VIII, 143 (1874)	—	S.A.	—	V.	N.S.W.	Q.	—	—				

ABROTANELLA ; Cassini in Dict. XXXVI, 27 (1825). (Scleroleima, Trineuron.)													
A. fosterioides ; J. Hooker, Hand. N. Zeal. Fl. 139 (1864)...	...	...	...	—	—	T.	—	—	—	—	B.fl.III,554		
A. nivigena ; F. v. M. in Benth. Fl. Austr. III, 554 (1866)...	...	...	...	—	—	—	V.	—	—	—	B.fl.III,554		
A. scapigera ; F. v. M. in Benth. Fl. Austr. III, 554 (1866)	...	...	...	—	—	T.	—	—	—	—	B.fl.III,554		
ELACHANTHUS ; F. v. M. in Schlecht. Linnaea XXV, 410 (1852).													
E. pusillus ; F. v. M. in Schlecht. Linnaea XXV, 411 (1852)	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,555		
DIMORPHOCOMA ; F. v. M. & Tate in Trans. Roy. Soc. of S. Austr. VI, 107 (1883).													
D. minutula ; F. v. M. & Tate in Transact. Roy. Soc. of S.A. VI, 107 (1883)	...	—	S.A.	—	—	—	—	—	—		—		
CERATOGYNE ; Turczaninow in Bull. de Mosc. XXIV, 68 (1851). (Diotosperma.)													
C. obionoides ; Turczaninow in Bull. de Mosc. XXIV, 68 (1851)	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,555	M.fr.III,159.	
ISOETOPSIS ; Turczaninow in Bull. de Mosc. XXIV, 175 (1851).													
I. graminifolia ; Turczaninow in Bull. de Mosc. XXIV, 175, t. 3 (1851)	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,556			
EMILIA ; Cassini in Bulletin de la Soc. philomat. 68 (1817).													
E. purpurea ; Cassini in Diction. des sc. nat. XXXIV, 393 (1825)	...	...	—	—	—	—	—	Q.	—	—	M.fr.XII ...	...	As., Afr.
GYNURA ; Cassini in Diction. des sc. nat. XXXIV, 391 (1825).													
G. Pseudo-China ; De Candolle, prodr. VI, 299 (1837)	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,661	...	...	...	As.
SENECIO ; Tournefort, inst. 456, t. 260 (1700), from Plinius. (Bedfordia, Centropappus.)													
S. Gregorii ; F. v. M., pl. of Greg. Search. Exped. for Leichh. 7 (1858)	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,663			
S. platylepis ; De Candolle, prodr. VI, 371 (1837)	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,664	M.fr.XI,27.		
S. papillosus ; F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857)	...	...	...	—	—	T.	—	—	—	—	B.fl.III,664		
S. primulifolius ; F. v. M. in Transact. phil. Inst. Vict. II, 69 (1857)	...	...	...	—	—	T.	—	—	—	—	B.fl.III,664		
S. pectinatus ; De Candolle, prodr. VI, 372 (1837)	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,664		
S. spathulatus ; A. Richard, sert. Astrol. 125 (1833)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,665		
S. megaglossus ; F. v. M. in Linnaea XXV, 419 (1852)	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.III,665	M.fr.VI,255.		
S. magnificus ; F. v. M. in Linnaea XXV, 418 (1852)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,665		
S. insularis ; Bentham, Fl. Austr. III, 666 (1866)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,666		
S. Centropappus ; F. v. M., annual report 26 (1858)	...	...	...	...	—	—	T.	—	—	—	—	B.fl.III,666		
S. macranthus ; A. Richard, sert. Astrol. 126 (1833)	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.III,666		
S. Daltoni ; F. v. M., fragm. VI, 27 (1867)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.VI,27.		
S. lautus ; Solander in G. Forst. prodr. 91 (1786)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.III,667	M.fr.XI,20	...	N.Z.
S. capillifolius ; J. Hooker in Lond. Journ. VI. 123 (1847)	...	...	...	...	—	—	T.	—	—	—	—	B.fl.III,667		
S. vagus ; F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854)	...	..	...	—	—	—	V.	N.S.W.	—	—	B.fl.III,667		
S. amygdalifolius ; F. v. M., fragm. I, 232 (1859)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.III,668	M.fr.I,232.		
S. velleioides ; Cunningham in De Candolle, prodr. VI, 374 (1837)	...	...	...	...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.III,668		
S. dryadens ; Sieber in Spreng. syst. veg. III, 562 (1826)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,668		
S. Behrianus ; Sonder & F. v. M. in Linnaea XXV, 527 (1852)	...	...	...	...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,669		
S. leucoglossus ; F. v. M., fragm. II, 15 (1861)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,669	M.fr.II,15.		
S. brachyglossus ; F. v. M. in Schlecht. Linnaea XXV, 525 (1852)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,669	M.fr.XI,27.		
S. Georgianus ; De Candolle, prodr. VI, 371 (1837)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,670		
S. Gilberti ; Turczaninow in Bull. Soc. Mosc. XXIV, 208 (1851)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,670		
S. ramosissimus ; De Candolle, prodr. VI, 371 (1837)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.III,671		
S. odoratus ; Hornemann, hort. reg. bot. Hafn. II, 809 (1815)	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,671		
S. Cunninghamii ; De Candolle, prodr. VI, 371 (1837)	...	...	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.III,671		
S. anethifolius ; Cunningham in De Candolle, prodr. VI, 371 (1837)	...	...	...	...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.III,672		
S. Bedfordii ; F. v. M., annual report 26 (1858)	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.III,673		
S. Billardieri ; F. v. M., annual report 26 (1858)	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.III,673		

ERECHTITES ; Rafinesque, Fl. Ludov. 65 (1817).

E. prenanthoides ; De Candolle, prodr. VI, 296 (1837)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,658				N.Z.
E. Atkinsoniae ; F. v. M., fragm. V, 88 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,658	M.fr.V,88.			
E. arguta ; De Candolle, prodr. VI, 296 (1837)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,659				N.Z.
E. picridioides ; Turczaninow in Bullet. de Mosc. XXIV, 200 (1851)	W.A.	S.A.	—	V.	—	—	—	—				
R. mixta ; De Candolle, prodr. VI, 297 (1837)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.III,659				
E. quadridentata ; De Candolle, prodr. VI, 295 (1837)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,660				N.Z.
E. hispidula ; De Candolle, prodr. VI, 296 (1837)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.III,660				

CYMBONOTUS ; Cassini in Diction. XXXV, 397 (1825).

C. Lawsonianus ; Gaudichaud in Freyc. Voy. Bot. 462, t. 86 (1826)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,674	M.fr.XI,27.			

SAUSSUREA ; De Candolle in Ann. du Mus. XVI, 156 (1810). (Aplotaxis, Haplotaxis.)

S. affinis ; Sprengel in De Candolle, prodr. VI, 540 (1837)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,456	M.fr.I,36			As.

CENTAUREA ; Linné, hort. Cliffort. 420 (1737). (Leuzea partly.)

C. Australis ; Bentham & J. Hooker, gen. plant. II, 479 (1873)	—	—	—	V.	N.S.W.	Q.	—	B.fl.III,457				

TRICHOCLINE ; Cassini in Bull. de la Soc. philom. 13 (1817). (Amblysperma, Gerberae subgenus.)

T. scapigera ; Bentham & J. Hooker, gen. plant. II, 479 (1873)	W.A.	—	—	—	—	—	—	B.fl.III,676				

MICROSERIS ; D. Don in Edinb. phil. Mag. XI, 388 (1832). (Phyllopappus, Scozonera partly.)

M. Forsteri ; J. Hooker, Fl. N. Zeal. I, 151 (1853)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.III,676	M.fr.XI,27			N.Z.

CREPIS ; Linné, gen. pl. 240 (1737). (Youngia.)

C. Japonica ; Bentham, Fl. Hongk. 194 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.III,679				As., Afr., P.

CAMPANULACEAE.

A. L. DE JUSSIEU, gen. plant. 163 (1789), from B. de Jussieu (1759).—Genera 4, Species 35.

LOBELIA ; Linné, Fl. Lappon. 227 (1737). (Rapuntium, Pratia, Grammatotheca, Holostigma, Isolobus.)

L. heterophylla ; Labillardière, Nov. Holl. pl. spec. I, 52, t. 74 (1804)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,124	M.fr.IV,184; X,41.			
L. Browniana ; Roemer & Schultes, syst. veg. V, 71 (1819)	—	S.A.	T.	V.	N.S.W.	Q.	—	—	M.fr.X,41.			
L. simplicicaulis ; R. Brown, prodr. 564 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,124	M.fr.IV,184; X,41.			
L. microsperma ; F. v. M., fragm. X, 41 (1876)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	—	M.fr.X,41.			
L. dentata ; Cavanilles, icon. pl. VI, 14, t. 522 (1801)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,125	M.fr.IV,183; X,42.			
L. gracilis ; Andrews, Bot. Reposit. t. 340 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,125	M.fr.IV,183; X,42.			
L. rhytidosperma ; Bentham, Fl. Austr. IV, 126 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,126	M.fr.X,40.			
L. tenuior ; R. Brown, prodr. 564 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,126	M.fr.IV,183; X,42.			
L. rhombifolia ; De Vriese in Lehm. pl. Preiss. I, 397 (1845)	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,126	M.fr.X,42.			
L. parvifolia ; R. Brown, prodr. 564 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,127	M.fr.IV,183.			
L. trigonocaulis ; F. v. M., fragm. I, 18 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,127	M.fr.I,18,242.			
L. Bergiana ; Chamisso in Schlecht. Linnaea VIII, 217 (1835)	W.A.	—	—	—	—	—	—	B.fl.IV,127	M.fr.II,20; IV,171; X,43.			Afr.
L. anceps ; Thunberg. prodr. plant. Cap. 40 (1794)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,128	M.fr.IV,184; VI,114; X,43			Afr., Am., N.Z.
L. surrepens ; J. Hooker, Fl. Tasman. I, 237, t. 69 (1860)	—	—	T.	—	—	—	—	B.fl.IV,129	M.fr.IV,183.			
L. membranacea ; R. Brown, prodr. 563 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,129	M.fr.IV,171; X,42.			
L. stenophylla ; Bentham, Fl. Austr. III, 130 (1869)	—	—	—	—	—	Q.	N.A.	B.fl.IV,130	M.fr.X,42.			
L. humistrata ; F. v. M. in Benth. Fl. Austr. III, 130 (1869)	W.A.	—	—	—	—	Q.	N.A.	B.fl.IV,130	M.fr.IV,171; X,42.			
L. dioica ; R. Brown, prodr. 565 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,130	M.fr.IV,183; X,43.			
L. purpurascens ; R. Brown, prodr. 563 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,131	M.fr.IV,183; X,42.			
L. pratioides ; Bentham, Fl. Austr. IV, 131 (1869)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,131	M.fr.X,43.			
L. irrigua ; R. Brown, prodr. 563 (1810)	—	—	T.	—	—	—	—	B.fl.IV,132				

L. gelida; F. v. M., fragm. IV, 183 (1864)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,132	M.fr.IV,183.	
L. platycalyx; F. v. M., fragm. IV, 183 (1864)	—	S.A.	T.	V.	—	—	—	B.fl.IV,133	M.fr.X,43.	
L. concolor; R. Brown, prodr. 563 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,133	M.fr.X,43.	
L. Benthami; F. v. M., Cens. of Austr. pl. 85 (1882)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,133		
L. pedunculata; R. Brown, prodr. 563 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,133	M.fr.X,42.	
ISOTOMA; R. Brown, prodr. 565 (1810). (Lobelia partly, Enchysia partly, Laurencia partly.)										
I. Brownii; G. Don, gen. syst. III, 716 (1834)	W.A.	—	—	—	—	—	—	B.fl.IV,134	M.fr.X,40.	
I. axillaris; Lindley, Bot. Regist. t. 964 (1826)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,135	M.fr.X,39.	
I. petraea; F. v. M. in Linnaea XXV, 420 (1852)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,135	M.fr.X,39.	
I. pusilla; Bentham in Hueg. enum. 75 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,135		
I. Gulliverii; F. v. M., fragm. X, 39 (1876)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,39.	
I. scapigera; G. Don, gen. syst. III, 716 (1834)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,136	M.fr.IV,183; X,39.	
I. fluviatilis; F. v. M. in Bentham, Fl. Austr. IV, 136 (1869)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,136	M.fr.X,39.	
WAHLENBERGIA; Schrader, cat. hort. bot. Goetting. (1814). (Strzeleckia.)										
W. gracilis; A. de Candolle, Monogr. comp. 142 (1830)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,137	...	As., N.Z.
SPHENOCLEA; Gaertner, de fructibus I, 113 (1788).										
S. Zeilanica; Gaertner de fruct. I, 113, t. 24 (1788)	—	—	—	—	—	—	N.A.	—	...	As., Afr., Am.

CANDOLLEACEAE.

F. v. M., fragm. VIII, 41 (1873). (Stylideae; R. Brown, prodr. 565 (1810).—Genera 4, Species 97.

CANDOLLEA; Labillardière in Ann. du Mus. VI, 451 (1805.) (Stylidium 1805, Ventenata, Forsteropsis.)

C. carnosa; F. v. M., Cens. 85 (1882) from Benth. (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,6		
C. pilosa; Labillardière in. Annal. du Mus. VI, 453, t. 63 (1805)	W.A.	—	—	—	—	—	—	B.fl.IV,7		
C. reduplicata; F. v. M., Cens. 85 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,7		
C. scabrida; F. v. M., Cens. 85 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,8		
C. hirsuta; F. v. M., Cens. 85 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,8		
C. crossocephala; F. v. M., Cens. 85 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,8	M.fr.VI,5,254; X,58.	
C. juncea; F. v. M., Cens. 85 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,9	M.fr.X,58.	
C. guttata; F. v. M., Cens. 85 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,9		
C. repens; F. v. M., Cens. 85 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,9	M.fr.X,58.	
C. serrulata; Labillardière in Annal. du Mus. VI, 454, t. 63 (1805)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,10	M.fr.IV,94; VI,6; X,58.	
C. linearis; F. v. M., Cens. 85 (1882) from Swartz (1807)	—	—	—	—	N.S.W.	—	—	B.fl.IV,11	M.fr.IV,95.	
C. elongata; F. v. M., Cens. 85 (1882) from Benth. (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,583	M.fr.X,86.	
C. spinulosa; F. v. M., Cens. 85 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,11		
C. Tepperiana; F. v. M., in Melb. Chem. and Drugg. II, 15 (1887)	—	S.A.	—	—	—	—	—	—		
C. limbata; F. v. M., Cens. 86 (1882)	W.A.	—	—	—	—	—	—	—	M.fr.X,57.	
C. caespitosa; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,11	M.fr.X,57.	
C. squamellosa; F. v. M., Cens. 86 (1882) from De Candolle (1838)	W.A.	—	—	—	—	—	—	—	M.fr.X,56.	
C. violacea; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,12		
C. lutea; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,12		
C. filifera; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,12	M.fr.X,58,86.	
C. ciliata; F. v. M., Cens. 86 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,13		
C. sobolifera; F. v. M., Cens. 86 (1882)	—	—	—	V.	—	—	—	B.fl.IV,13	M.fr.IV,94.	
C. Flondii; F. v. M., Cens. 86 (1882)	—	—	—	—	—	Q.	N.A.	B.fl.IV,13	M.fr.I,149; VI,6; X,58.	
C. disperma; F. v. M., Cens. 86 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,14	M.fr.IV,93.	
C. calcarata; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	S.A.	—	V.	—	—	N.A.	B.fl.IV,14	M.fr.IV,94; VI,6.	
C. perpusilla; F. v. M., Cens. 86 (1882) from J. Hooker (1847)	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,15	M.fr.VI,78; X,58.	

C. eriorrhiza ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	—	B.fl.IV,15	M.fr.I,147; VI,6; X,58.
C. debilis ; F. v. M., Cens. 86 (1882)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,15	M.fr.I,149; VI,6; X,58.
C. floribunda ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,15	M.fr.I,148; VI,6; X,58.
C. leptorrhiza ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,16	M.fr.I,148.
C. assimilis ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,16	M.fr.X,57.
C. rupestris ; F. v. M., Cens. 86 (1882) from Sonder (1845)...	W.A.	—	—	—	—	—	—	B.fl.IV,16	
C. spathulata ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,17	
C. Barleei ; F. v. M., Cens. 86 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,17	M.fr.VI,5,249.
C. lineata ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,17	
C. glauca ; Labillardière in Annal. du Mus. VI, 454, t. 64 (1805)	W.A.	—	—	—	—	—	—	B.fl.IV,17	
C. amoena ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,18	
C. striata ; F. v. M., Cens. 86 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,18	
C. diversifolia ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,18	M.fr.X,58.
C. articulata ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,19	
C. Brunoniana ; F. v. M., Cens. 86 (1882) from Bentham (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,19	
C. diuroides ; F. v. M., Cens. 86 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,20	
C. scandens ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,20	M.fr.VI,78.
C. verticillata ; F. v. M., Cens. 86 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,20	M.fr.IV,94.
C. glandulosa ; F. v. M., Cens. 86 (1882) from Salisbury (1807)	W.A.	—	—	—	—	—	—	B.fl.IV,20	
C. laricifolia ; F. v. M., Cens. 86 (1882) from Richard (1807)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,21	
C. Preissii ; F. v. M., Cens. 86 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,21	M.fr.III,122.
C. imbricata ; F. v. M., Cens. 86 (1882) from Bentham (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,21	
C. adpressa ; F. v. M., Cens. 86 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,22	
C. despecta ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,22	M.fr.IV,94; VI,58.
C. utricularioides ; F. v. M., Cens. 86 (1882) from Bentham (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,23	
C. pygmaea ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,23	
C. longitubea ; F. v. M., Cens. 86 (1882) from Bentham (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,23	
C. diffusa ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	—	B.fl.IV,23	
C. fissiloba ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,24	M.fr.I,154.
C. trichopoda ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	Q.	—	—	M.fr.X,86.
C. alsinoides ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,24	M.fr.I,151; VI,6; X,58.
C. tenerrima ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,24	M.fr.I,150.
C. brachyphylla ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,24	
C. capillaris ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	—	B.fl.IV,25	
C. rotundifolia ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,25	M.fr.I,151.
C. schizantha ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,25	M.fr.I,152.
C. lobuliflora ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,25	M.fr.I,153.
C. uliginosa ; F. v. M., Cens. 86 (1882) from Swartz (1807)	—	—	—	—	—	Q.	—	B.fl.IV,26	... ... As.
C. pulchella ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,26	
C. petiolaris ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,26	
C. emarginata ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. corymbosa ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. lepida ; F. v. M., Cens. 86 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. streptocarpa ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,27	
C. uniflora ; F. v. M., Cens. 86 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,28	M.fr.VI,6.
C. pedunculata ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,28	M.fr.VI,91.
C. pachyrrhiza ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,28	M.fr.I,152.
C. muscicola ; F. v. M., Cens. 86 (1882)	—	—	—	—	—	—	N.A.	B.fl.IV,29	M.fr.I,153.
C. crassifolia ; F. v. M., Cens. 86 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,29	
C. pycnostachya ; F. v. M., Cens. 86 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,29	

C. pubigera; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30				
C. canaliculata; F. v. M., Cens. 86 (1882) from Lindley (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30				
C. leptophylla; F. v. M., Cens. 86 (1882) from De Candolle (1838) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30				
C. dichotoma; F. v. M., Cens. 86 (1882) from De Candolle (1838) ...	W.A.	—	—	—	—	—	—	B.fl.IV,30				
C. bulbifera; F. v. M., Cens. 86 (1882) from Bentham (1837) ...	W.A.	—	—	—	—	—	—	B.fl.IV,31				
C. Merralliana; F. v. M. in Vict. Naturalist V, 76 (1888) ...	W.A.	—	—	—	—	—	—	—				
C. breviscapa; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,31				
C. eglandulosa; F. v. M., Cens. 86 (1882) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,31	M.fr.I,150; VI,6; X,58.			
C. fasciculata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,32				
C. falcata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,32				
C. rhynchocarpa; F. v. M., Cens. 86 (1882) from Sonder (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,33				
C. adnata; F. v. M., Cens. 86 (1882) from R. Brown (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,33	M.fr.X,58.			
LEEWENHOEKIA; R. Brown, prodr. 572 (1810). (Levenhookia, Coleostylis.)												
L. pusilla; R. Brown, prodr. 573 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,34	M.fr.X,58.			
L. dubia; Sonder in Lehm. pl. Preiss. I, 392 (1845) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,34	M.fr.III,121; IV,94; X,58.			
L. Sonderi; F. v. M., fragm. I, 18 (1858) ...	—	—	—	V.	—	—	—	B.fl.IV,35				
L. pauciflora; Bentham in Hueg. enum. 74 (1837) ...	W.A.	—	—	—	—	—	—	B.fl.IV,35	M.fr.VI,77.			
L. leptantha; Bentham, Fl. Austr. IV, 35 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,35				
L. stipitata; F. v. M., fragm. IV, 94 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.			
L. Preissii; F. v. M., fragm. IV, 94 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,36	M.fr.IV,94.			
PHYLLACHNE; R. et G. Forster, char. gen. 115, t. 58 (1776). (Forstera, Helophyllum, Oreostylidium.)												
P. bellidifolia; F. v. M., fragm. VIII, 39 (1873) ...	—	—	T.	—	—	—	—	B.fl.IV,37	M.fr.X,58.			
DONATIA; R. et G. Forster, char. gen. 9, t. 5 (1776).												
D. Novae Zealandiae; J. Hooker, Fl. N. Zeal. I, 81, t. 20 (1853) ...	—	—	T.	—	—	—	—	B.fl.II,450	M.fr.IV,3; VII,150; VIII,41; X,58.	N.Z.		

GOODENIACEAE.

R. Brown, prodr. 573 (1810).—Genera 11, Species 220.

BRUNONIA; Smith in Transact. Linn. Soc. X, 366 (1809).												
B. Australis; Smith in Transact. Linn. Soc. X, 367, t. 28 (1809) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,121	M.fr.X,111.			
DAMPIERA; R. Brown, prodr. 587 (1810). (Linschottenia.)												
D. luteiflora; F. v. M., fragm. X, 11 (1876) ...	W.A.	—	—	—	—	—	—	—	M.fr.X,11.			
D. Linschotenii; F. v. M., fragm. VI, 28 (1866) ...	—	—	—	—	—	Q.	—	B.fl.IV,108	M.fr.VI,28.			
D. spicigera; Bentham, Fl. Austr. IV, 199 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,109				
D. candicans; F. v. M., fragm. X, 86 (1876) ...	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,86.			
D. teres; Lindley, Bot. Regist. XXV, App. 27 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,109				
D. trigona; De Vriese in Lehm. pl. Preiss. I, 401 (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,109	M.fr.II,17; VIII,245; X,11.			
D. prostrata; De Vriese, Goodenov. 83 (1854) ...	W.A.	—	—	—	—	—	—	B.fl.IV,110				
D. alata; Lindley, Bot. Regist. XXV, App. 27 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,110	M.fr.X,11.			
D. coronata; Lindley, Bot. Regist. XXV, App. 27 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.IV,110				
D. carinata; Bentham, Fl. Austr. IV, 111 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,111				
D. sacculata; F. v. M. in Benth. Fl. Austr. IV, 111 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,111				
D. incana; R. Brown, prodr. 588 (1810) ...	W.A.	—	—	—	—	—	N.A.	B.fl.IV,111	M.fr.II,17.			
D. hederacea; R. Brown, prodr. 588 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,112	M.fr.X,12.			
D. ferruginea; R. Brown, prodr. 588 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,112	M.fr.VI,29.			
D. Brownii; F. v. M., fragm. VI, 29 (1866) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,112	M.fr.VI,29.			
D. lanceolata; Cunningham in De Candolle, prodr. VII, 503 (1839) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,113	M.fr.VI,29.			
D. altissima; F. v. M. in Benth. Fl. Austr. IV, 113 (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,113				

D. marifolia ; Bentham, Fl. Austr. IV, 114 (1869)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,114	
D. rosmarinifolia ; Schlechtendal, Linnaea XX, 603 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,114	M.fr.VI,29; X,12.
D. lavandulacea ; Lindley, Bot. Regist. XXV, App. 27 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,114	
D. juncea ; Bentham, Fl. Austr. IV, 115 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,115	
D. oligophylla ; Bentham, Fl. Austr. IV, 115 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,115	
D. loranthifolia ; F. v. M. in Benth. Fl. Austr. IV, 115 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,115	
D. stricta ; R. Brown, prodr. 589 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,116	M.fr.VI,29; X,12; XI,121.
D. Scottiana ; F. v. M., fragm. XI, 120 (1881)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,120.
D. leptoclada ; Bentham, Fl. Austr. IV, 116 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,116	
D. fasciculata ; R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,116	M.fr.XI,121.
D. subspicata ; Bentham, Fl. Austr. IV, 117 (1869)...	W.A.	—	—	—	—	—	—	B.fl.IV,117	
D. triloba ; Lindley, Bot. Regist. XXV, App. 27 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,117	
D. linearis ; R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,117	
D. cuneata ; R. Brown, prodr. 588 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,118	
D. sericantha ; F. v. M. in Benth. Fl. Austr. IV, 118 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,118	
D. parvifolia ; R. Brown, prodr. 589 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,118	
D. glabrescens ; Bentham, Fl. Austr. IV, 119 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,119	
D. adpressa ; Cunningham in De Candolle, prodr. VII, 503 (1839)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,119	
D. diversifolia ; De Vriese in Lehm. pl. Preiss. I, 403 (1845)	W.A.	—	—	—	—	Q.	—	B.fl.IV,119	M.fr.I,20.
D. eriocephala ; De Vriese, Goodenov. 118, t. 21 (1854)	W.A.	—	—	—	—	—	—	B.fl.IV,120	M.fr.VI,29; X,12.
D. Wellsiana ; F. v. M., fragm. X, 12 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,12.
DIASPASIS ; R. Brown, prodr. 586 (1810).									
D. filifolia ; R. Brown, prodr. 587 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.I,206; II,181; VIII,58.
LESCHENAULTIA ; R. Brown, prodr. 581 (1810). (Latourea.)									
L. formosa ; R. Brown, prodr. 581 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,40	M.fr.VI,226.
L. oblata ; Sweet, flor. Australas. 26, 46, t. 46 (1828)	W.A.	—	—	—	—	—	—	—	M.fr.VI,226.
L. chlorantha ; F. v. M., fragm. II, 20 (1860)	W.A.	—	—	—	—	—	—	B.fl.IV,40	M.fr.II,20.
L. linarioides ; De Candolle, prodr. VII, 519 (1838)...	W.A.	—	—	—	—	—	—	B.fl.IV,40	
L. tubiflora ; R. Brown, prodr. 581 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,41	
L. superba ; F. v. M., fragm. VI, 10 (1866)	W.A.	—	—	—	—	—	—	B.fl.IV,41	M.fr.VI,10.
L. striata ; F. v. M., fragm. VIII, 245 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,245.
L. acutiloba ; Bentham, Fl. Austr. IV, 41 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,41	
L. laricina ; Lindley, Bot. Regist. XXV, App. 27 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,41	
L. hirsuta ; F. v. M., fragm. VI, 9 (1866)	W.A.	—	—	—	—	—	—	B.fl.IV,42	M.fr.XII,23.
L. longiloba ; F. v. M., fragm. VI, 10 (1866)	W.A.	—	—	—	—	—	—	B.fl.IV,42	M.fr.XII,23.
L. biloba ; Lindley, Bot. Regist. XXV, App. 27 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,42	
L. expansa ; R. Brown, prodr. 581 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,42	
L. floribunda ; Bentham in Hueg. enum. 70 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,43	M.fr.II,20.
L. heteromera ; Bentham, Fl. Austr. IV, 43 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,43	
L. divaricata ; F. v. M., fragm. III, 33, 167 (1863)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,43	M.fr.VIII,246.
L. filiformis ; R. Brown, prodr. 581 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,44	M.fr.VI,9,249.
L. agrostophylla ; F. v. M., fragm. VI, 8, pl. (1866)	—	—	—	—	—	—	N.A.	B.fl.IV,44	M.fr.VI,8; 249.
ANTHOTIUM ; R. Brown, prodr. 582 (1810).									
A. humile ; R. Brown, prodr. 582 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,44	M.fr.VIII,58.
A. rubriflorum ; F. v. M. in Benth. Fl. Austr. IV, 45 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,45	
CATOSPERMA ; Bentham in Hooker, icon. pl. t. 1028 (1868).									
C. Muelleri ; Bentham in Hook. icon. pl. t. 1028 (1868)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,83	M.fr.I,121; VIII,58,245.
C. Careyi ; F. v. M. in Melb. Chemist, Apr. (1884)	W.A.	—	—	—	—	—	—	—	

SCAEVOLA; Linné, mantiss. II, 145 (1771). (Pogonetes, Crossotoma, Temminckia, Kamphusia, Molkenboeria, Merkusia, Verreauxia.)

S. Koenigii; Vahl, symb. bot. III, 36 (1794)	—	—	—	—	—	Q. N.A.	B.fl.IV,86	M.fr.XI,76	As., Afr.
S. enantophylla; F. v. M., fragm. VIII, 58 (1873)	—	—	—	—	—	Q. —	—	M.fr.VI,225.	
S. spinescens, R. Brown, prodr. 586 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q. N.A.	B.fl.IV,87	M.fr.XI,20,27,76.	
S. Groeneri; F. v. M., fragm. VI, 15 (1866)	W.A.	S.A.	—	—	—	— —	B.fl.IV,88	M.fr.VIII,58.	
S. tomentosa; Gaudichaud in Freyc. Voy. Bot. 460, t. 81 (1826)	W.A.	—	—	—	—	— —	B.fl.IV,88		
S. atriplicina; F. v. M., fragm. II, 18 (1860)	W.A.	—	—	—	—	— —	B.fl.IV,88	M.fr.III,165,181.	
S. striata; R. Brown, prodr. 586 (1810)	W.A.	—	—	—	—	— —	B.fl.IV,89		
S. phlebopetala; F. v. M., fragm. II, 18 (1860)	W.A.	—	—	—	—	— —	B.fl.IV,89	M.fr.II,VI,16; XI,76.	
S. pilosa; Bentham in Hueg. enum. 69 (1837)	W.A.	—	—	—	—	— —	B.fl.IV,89		
S. hispida; Cavanilles, icon. et descr. pl. VI, 7, t. 510 (1801)	—	—	—	V.	N.S.W.	Q. —	B.fl.IV,90	M.fr.I,121,207; XI,76.	
S. apterantha; F. v. M., fragm. I, 121 (1859)	—	—	—	V.	—	— —	B.fl.IV,90	M.fr.I,121.	
S. Hookeri; F. v. M., gen. report, 15 (1853)	—	—	T.	V.	N.S.W.	— —	B.fl.IV,90	M.fr.XI,76.	
S. parvifolia; F. v. M. in Benth. Fl. Austr. IV, 91 (1869)	W.A.	S.A.	—	—	—	Q. N.A.	B.fl.IV,91	M.fr.VIII,58; X,59; XI,76.	
S. oxyclona; F. v. M., fragm. X, 58 (1876)	W.A.	—	—	—	—	— —	—	M.fr.X,58.	
S. restiacea; Bentham, Fl. Austr. IV, 91 (1869)	W.A.	—	—	—	—	— —	B.fl.IV,91		
S. depauperata; R. Brown, App. Sturt's Exped. 20 (1849)	—	S.A.	—	—	—	— N.A.	B.fl.IV,91	M.fr.III,33; VIII,58; XI,76.	
S. tortuosa; Bentham, Fl. Austr. IV, 91 (1869)	W.A.	—	—	—	—	— —	B.fl.IV,91		
S. Cunninghamii; De Candolle, prodr. VII, 508 (1838)	—	—	—	—	—	— N.A.	B.fl.IV,92		
S. collaris; F. v. M., Rep. Babb. Exped. 15 (1858)	—	S.A.	—	—	—	— —	B.fl.IV,92	M.fr.II,20; VIII,58; XI,76.	
S. angulata; R. Brown, prodr. 586 (1810)	—	—	—	—	—	Q. N.A.	B.fl.IV,92		
S. Brooksiana; F. v. M. in Vict. Naturalist I, 122 (1884)	W.A.	—	—	—	—	— —	—		
S. nitida; R. Brown, prodr. 584 (1810)	W.A.	—	—	—	—	— —	B.fl.IV,93		
S. globuliflora; Labillardière, Nov. Holl. pl. spec. I, 55, t. 78 (1804)	W.A.	—	—	—	—	— —	B.fl.IV,93		
S. porocarya; F. v. M., fragm. II, 19 (1860)	W.A.	—	—	—	—	— —	B.fl.IV,94	M.fr.II,19; XI,76.	
S. attenuata; R. Brown, prodr. 583 (1810)	W.A.	—	—	—	—	— —	B.fl.IV,94		
S. glandulifera; De Candolle, prodr. VII, 510 (1838)	W.A.	—	—	—	—	— —	B.fl.IV,94		
S. anchusifolia; Bentham in Hueg. enum. 68 (1837)	W.A.	—	—	—	—	— —	B.fl.IV,94		
S. holosericea; De Vriese in Lehm. pl. Preiss. I, 408 (1845)	W.A.	—	—	—	—	— —	B.fl.IV,95		
S. suaveolens; R. Brown, prodr. 585 (1810)	—	S.A.	—	V.	N.S.W.	Q. —	B.fl.IV,95		
S. revoluta; R. Brown, prodr. 586 (1810)	—	—	—	—	—	Q. N.A.	B.fl.IV,96	M.fr.I,121.	
S. crassifolia; Labillardière, Nov. Holl. pl. spec. I, 56, t. 79 (1804)	W.A.	S.A.	—	V.	—	— —	B.fl.IV,96	M.fr.VIII,245; XI,20,76.	
S. macrostachya; Bentham, Fl. Austr. IV, 97 (1869)	—	—	—	—	—	— N.A.	B.fl.IV,97	M.fr.VIII,58; XI,76.	
S. longifolia; De Vriese in Lehm. pl. Preiss. I, 410 (1845)	W.A.	—	—	—	—	— —	B.fl.IV,97		
S. lanceolata; Bentham in Hueg. enum. 69 (1837)	W.A.	—	—	—	—	— —	B.fl.IV,97	M.fr.I,207; XI,76.	
S. thesioides; Bentham in Hueg. enum. 68 (1837)	W.A.	—	—	—	—	— —	B.fl.IV,98	M.fr.I,207; XI,77.	
S. macrophylla; Bentham, Fl. Austr. IV, 98 (1869)	W.A.	—	—	—	—	— —	B.fl.IV,98		
S. platyphylla; Lindley, Bot. Regist. XXV, App. 26 (1839)	W.A.	—	—	—	—	— —	B.fl.IV,98	M.fr.XI,77.	
S. auriculata; Bentham, Fl. Austr. IV, 99 (1869)	W.A.	—	—	—	—	— —	B.fl.IV,99	M.fr.XI,77.	
S. ovalifolia; R. Brown, prodr. 584 (1810)	W.A.	S.A.	—	—	N.S.W.	Q. N.A.	B.fl.IV,96	M.fr.VIII,58; XI,76.	
S. aemula; R. Brown, prodr. 584 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q. —	B.fl.IV,99	M.fr.XI,77.	
S. humilis; R. Brown, prodr. 585 (1810)	—	S.A.	—	—	—	— —	B.fl.IV,100		
S. amblyanthera; F. v. M., fragm. I, 121 (1859)	—	—	—	—	—	Q. N.A.	B.fl.IV,100	M.fr.I,121.	
S. microphylla; Bentham, Fl. Austr. IV, 100 (1869)	W.A.	—	—	—	—	— —	B.fl.IV,100		
S. cuneiformis; Labillardière, Nov. Holl. pl. spec. I, 56, t. 80 (1804)	W.A.	—	—	—	—	— —	B.fl.IV,101	M.fr.XI,77.	
S. microcarpa; Cavanilles, icon. et descr. pl. VI, 6, t. 509 (1801)	—	S.A.	T.	V.	N.S.W.	Q. —	B.fl.IV,101	M.fr.VI,16; XI,77.	
S. linearis; R. Brown, prodr. 586 (1810)	—	S.A.	—	—	—	— —	B.fl.IV,102		
S. Oldfieldii; F. v. M., fragm. II, 19 (1860)	W.A.	—	—	—	—	— —	B.fl.IV,102	M.fr.II,19; XI,77.	
S. paludosa; R. Brown, prodr. 586 (1810)	W.A.	—	—	—	—	— —	B.fl.IV,102	M.fr.XI,77.	
S. sericophylla; F. v. M. in Benth. Fl. Austr. IV, 102 (1869)	W.A.	—	—	—	—	— —	B.fl.IV,102	M.fr.XI,77.	

S. canescens ; Bentham in Hueg. enum. 69 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,103							
S. humifusa ; De Vriese in Lehm. pl. Preiss. I, 410 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,103	M.fr.XI,77.						
S. fasciculata ; Bentham in Hueg. enum. 68 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.XI,77.						
S. stenophylla ; Bentham, Fl. Austr. IV, 104 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,104	M.fr.I,113; XI,77.						
S. Reinwardtii ; De Vriese in Lehm. pl. Preiss. I, 409 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,106	M.fr.II,17.						
S. Verreauxii ; F. v. M., Bot. Teachings 65 (1877)	W.A.	—	—	—	—	—	—	B.fl.IV,105	M.fr.II,17.						
SELLIERA ; Cavanilles in Anal. Cienz. Nat. I, 41, t. 5 (1799).															
S. radicans ; Cavanilles, icon. et descr. pl. V, 49, t. 474 (1799)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,82	M.fr.VIII,58	...	...	Am., N.Z.			
S. exigua ; F. v. M., fragm. III, 142 (1862)	W.A.	—	—	—	—	—	—	B.fl.IV,82	M.fr.III,142.						
CALOGYNE ; R. Brown, prodr. 579 (1810). (Distylis.)															
C. pilosa ; R. Brown, prodr. 579 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,81	M.fr.VI,7,28.						
C. Berardiana ; F. v. M., fragm. VI, 7 (1867)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,81	M.fr.VI,7; VIII,58.						
C. heteroptera ; F. v. M., fragm. X, 43 (1876)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,43.						
C. purpurea ; F. v. M., fragm. VIII, 57 (1873)	—	—	—	—	—	—	N.A.	—	M.fr.VIII,57.						
GOODENIA ; Smith in Transact. Linn. Soc. II, 347 (1794). (Goodenoughia, Tetraphylax, Picrophyta, Stekhovia, Aillya.)															
G. phylicoides ; F. v. M., fragm. I, 206 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,55	M.fr.I,206.						
G. viscida ; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,55	M.fr.I,120; III,35,163.						
G. xanthotricha ; De Vriese, Gooden. 155 (1854)	W.A.	—	—	—	—	—	—	B.fl.IV,56	M.fr.VI,13.						
G. scapigera ; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,56	M.fr.I,114;III,163; XI,50.						
G. quadrilocularis ; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,57	M.fr.III.141; IV.13.						
G. Ramelii ; F. v. M., fragm. III, 20, t. 17 (1862)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,57	M.fr.VIII,245; X,13.						
G. pinifolia ; De Vriese, Gooden. 157, t. 30 (1854)	W.A.	—	—	—	—	—	—	B.fl.IV,58							
G. decurrens ; R. Brown, prodr. 575 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,58							
G. racemosa ; F. v. M., fragm. I, 114 (1859)	—	—	—	—	—	Q.	—	B.fl.IV,58	M.fr.I,114.						
G. bellidifolia ; Smith in Transact. Linn. Soc. II, 349 (1794)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,58	M.fr.I,117; XI,50.						
G. stelligera ; R. Brown, prodr. 575 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,59	M.fr.VI,14.						
G. Stevensoni ; F. v. M. in Vict. Naturalist III, 138 (1887)	—	—	—	—	N.S.W.	—	—	—							
G. ovata ; Smith in Transact. Linn. Soc. II, 347 (1794)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,59	M.fr.I,205; X,110.						
G. varia ; R. Brown, prodr. 576 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,61	M.fr.I,205.						
G. amplexans ; F. v. M. in Transact. phil. Inst. Vict. II, 70 (1859)	—	S.A.	—	V.	—	—	—	B.fl.IV,60	M.fr.VI,15.						
G. strophiolata ; F. v. M., fragm. I, 119 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,60	M.fr.I,119; II,111.						
G. laevis ; Bentham, Fl. Austr, IV, 61 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,61							
G. barbata ; R. Brown, prodr. 576 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,61	M.fr.II,110,176; VI,15.						
G. arthrotricha ; F. v. M. in Benth. Fl. Austr. IV, 62 (1869)	W.A.	—	—	—	—	—	—	B.fl.IVi62							
G. disperma ; F. v. M., fragm. I, 113 (1859)	—	—	—	—	—	Q.	—	B.fl.IV,62	M.fr.IV,145; VI,15.						
G. geniculata ; R. Brown, prodr. 577 (1810)	W.A.	S A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,62	M.fr.VI,14.						
G. hederacea ; Smith in Transact. Linn. Soc. II, 349 (1794)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,63	M.fr.VI,15,110.						
G. xanthosperma ; F. v. M., fragm. X, 12 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,12.						
G. hirsuta ; F. v. M., fragm. III, 35 (1862)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,64	M.fr.III,35.						
G. heterophylla ; Smith in Transact. Linn. Soc. II, 349 (1794)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,64	M.fr.I,117.						
G. deflexa ; F. v. M. in Melb. Chem. Jan. (1883)	—	—	—	—	—	—	N.A.	—							
G. glabra ; R. Brown, prodr. 577 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,64	M.fr.VI,14.						
G. strongylophylla ; F. v. M., fragm. VI, 12 (1867)	—	—	—	—	—	Q.	—	B.fl.IV,65	M.fr.VI,12.						
G. rotundifolia ; R. Brown, prodr. 576 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,65	M.fr.VI,15.						
G. azurea ; F. v. M., fragm. I, 117 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,66	M.fr.X,87.						
G. scaevolina ; F. v. M., fragm. I, 118 (1859)...	—	—	—	—	—	—	N.A.	B.fl.IV,66	M.fr.XI,50.						
G. Stobbsiana ; F. v. M., fragm. XI, 49 (1879)	—	—	—	—	—	—	N.A.	—	M.fr.XI,49,137.						
G. incana ; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,66	M.fr.I,155; III,141.						
G. leptoclada ; Bentham, Fl. Austr. IV, 67 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,67	M.fr.VI,227; VIII,186; XI,50.						

G. Eatoniana; F. v. M., fragm. VIII, 186 (1874)	W.A.	—	—	—	—	—	—	M.fr.XI,51.
G. coerulea; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	B.fl.IV,67	M.fr.III,140; XI,20.
G. trichophylla; De Vriese in Benth. Fl. Austr. IV, 67 (1869)	W.A.	—	—	—	—	—	B.fl.IV,67	
G. Hassallii; F. v. M., fragm. VI, 10, t. 51 (1867)	W.A.	—	—	—	—	—	B.fl.IV,68	M.fr.XI,50.
G. pterygosperma; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	B.fl.IV,68	M.fr.I,155; III,163.
G. Vilmoriniae; F. v. M., fragm. III, 19, t. 16 (1862)	—	S.A.	—	—	—	—	N.A. B.fl.IV,68	M.fr.VIII,245.
G. Bonneyana; F. v. M. in Benth. Fl. Austr. IV, 69 (1869)	W.A.	—	—	—	—	—	B.fl.IV,69	M.fr.VI,226,250.
G. calcarata; F. v. M., fragm. VI, 14 (1867)	—	S.A.	—	—	N.S.W.	Q.	— B.fl.IV,69	M.fr.I,119; VIII,245.
G. albiflora; Schlechtendal, Linnaea XX, 599 (1847)	—	S.A.	—	—	—	—	— B.fl.IV,70	M.fr.I,119.
G. Nicholsoni; F. v. M., fragm. I, 203, t. 4 (1859)	—	S.A.	—	—	—	—	— B.fl.IV,69	M.fr.I,203.
G. Macmillani; F. v. M., fragm. I, 119, t. 5 (1859)	—	—	—	V.	—	—	— B.fl.IV,69	M.fr.I,119.
G. grandiflora; Sims, Bot. Mag. t. 890 (1806)	W.A.	S.A.	—	—	N.S.W.	Q.	— B.fl.IV,69	M.fr.I,204; VI,14.
G. Chambersii; F. v. M., fragm. I, 204 (1859)	—	S.A.	—	—	—	—	N.A. B.fl.IV,70	M.fr.I,204.
G. Strangfordii; F. v. M., fragm. VI, 11, t. 52 (1867)	—	S.A.	—	—	—	Q.	N.A. B.fl.IV,70	
G. Mitchellii; Bentham, Fl. Austr. IV, 71 (1869)	—	S.A.	—	—	—	Q.	— B.fl.IV,71	M.fr.XI,75.
G. Mueckeana; F. v. M., fragm. VIII, 56 (1873)	W.A.	S.A.	—	—	—	—	— —	M.fr.IX,194; XI,133.
G. melanoptera; F. v. M., fragm. I, 115 (1859)	—	—	—	—	—	—	N.A. —	M.fr.I,115.
G. heterochila; F. v. M., fragm. III, 142 (1863)	—	S.A.	—	—	—	—	N.A. B.fl.IV,71	M.fr.VI,12; VIII,245.
G. sepalosa; F. v. M. in Benth. Fl. Austr. IV, 72 (1869)	—	—	—	—	—	—	N.A. B.fl.IV,72	
G. hispida; R. Brown, prodr. 577 (1810)	—	—	—	—	—	Q.	N.A. B.fl.IV,72	M.fr.VI,12.
G. auriculata; Bentham, Fl. Austr. IV, 72 (1869)	—	—	—	—	—	—	N.A. B.fl.IV,72	
G. Armstrongiana; De Vriese, Goodenov. 138, t. 24 (1854)	—	—	—	—	—	—	N.A. B.fl.IV,73	M.fr.I,205; VII,40.
G. corynocarpa; F. v. M., fragm. II, 16 (1860)	W.A.	—	—	—	—	—	— B.fl.IV,73	M.fr.VI,15.
G. mollissima; F. v. M. in Benth. Fl. Austr. IV, 73 (1869)	—	—	—	—	—	Q.	— B.fl.IV,73	
G. O'Donnelli; F. v. M. in Austr. Journ. of Pharm. I, 278 (1886)	—	—	—	—	—	—	N.A. —	
G. cycloptera; R. Brown in App. Sturt's Exped. 83 (1849)	W.A.	S.A.	—	V.	N.S.W.	Q.	— B.fl.IV,74	M.fr.VI,14; XI,51.
G. tenella; R. Brown, prodr. 577 (1810)	W.A.	—	—	—	—	—	— B.fl.IV,74	M.fr.XI,51.
G. elongata; Labillardière, Nov. Holl. pl. spec. I, 52, t. 75 (1804)	—	S.A.	T.	V.	N.S.W.	—	— B.fl.IV,74	M.fr.VI,14; VIII,142,245; X,110.
G. pinnatifida; Schlechtendal, Linnaea XXI, 440 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	— B.fl.IV,75	M.fr.VI,14; XI,51.
G. pusilliflora; F. v. M., Key to the Syst. of Vict. pl. I, 354 (1888)	—	S.A.	—	V.	N.S.W.	Q.	— —	
G. tenuiloba; F. v. M. in Wing's S. Sc. Rec. new ser. I, 25 (1885)	W.A.	—	—	—	—	—	— —	
G. coronopifolia; R. Brown, prodr. 577 (1810)	—	—	—	—	—	Q.	N.A. B.fl.IV,75	M.fr.X,110.
G. heteromera; F. v. M., Sec. Rep. 13 (1854)	—	S.A.	—	V.	N.S.W.	Q.	— B.fl.IV,76	M.fr.I,115; VI,14; XI,27.
G. concinna; Bentham, Fl. Austr. IV, 76 (1869)	W.A.	—	—	—	—	—	— B.fl.IV,76	
G. glauca; F. v. M. in Transact. Vict. Inst. 40 (1855)	W.A.	S.A.	—	V.	N.S.W.	Q	N.A. B.fl.IV,76	M.fr.VI,14; XI,27.
G. filiformis; R. Brown, prodr. 578 (1810)	W.A.	—	—	—	—	—	— B.fl.IV,77	M.fr.I,116; VI,14; VIII,245.
G. Armitiana; F. v. M., fragm. X, 110 (1877)	—	—	—	—	—	Q.	N.A. —	M.fr.X,110.
G. cirrifica; F. v. M. in Austr. Journ. of Pharm. I, 81 (1886)	—	—	—	—	—	—	N.A. —	
G. microptera; F. v. M., fragm. III, 34 (1862)	—	S.A.	—	—	—	—	N.A. B.fl.IV,77	M.fr.VI,227; XI,75.
G. bicolor; F. v. M. in Benth. Fl. Austr. IV, 80 (1869)	—	—	—	—	—	—	N.A. B.fl.IV,80	
G. humilis; R. Brown, prodr. 575 (1810)	—	S.A.	T.	V.	N.S.W.	—	— B.fl.IV,79	M.fr.VII,227; VIII,142,245; X,110.
G. paniculata; Smith in Transact. Linn. Soc. II, 348 (1794)	—	—	—	V.	N.S.W.	Q.	— B.fl.IV,78	M.fr.I,116.
G. purpurascens; R. Brown, prodr. 578 (1810)	—	—	—	—	—	Q.	N.A. B.fl.IV,78	M.fr.I,117.
G. gracilis; R. Brown, prodr. 575 (1810)	—	—	—	V.	N.S.W.	Q.	N.A. B.fl.IV,79	M.fr.XI,51.
G. lamprosperma; F. v. M., fragm. I, 116 (1859)	W.A.	—	—	—	—	—	N.A. B.fl.IV,79	M.fr.III,34.
G. minutiflora; F. v. M., fragm. VIII, 244 (1874)	—	—	—	—	—	—	N.A. —	M.fr.VIII,244.
G. claytoniacea; F. v. M. in Benth. Fl. Austr. IV, 79 (1869)	W.A.	—	—	—	—	—	— B.fl.IV,79	M.fr.II,111.
G. Pumilio; R. Brown, prodr. 579 (1810)	—	—	—	—	—	Q.	N.A. B.fl.IV,80	M.fr.X,111.

VELLEYA; Smith in Transact. Linn. Soc. IV, 217 (1798). (Velleia, Euthales.)

V. Daviesii; F. v M., fragm. X, 10 (1876)	W.A.	—	—	—	—	—	— —	M.fr.X,10,,54.

V. panduriformis ; Cunningham in Benth. Fl. Austr. IV, 46 (1869)	—	—	—	—	—	—	—	N.A.	B.fl.IV,46	M.fr.XI,73.		
V. connata ; F. v. M. in Transact. phil. Soc. Vict. I, 18 (1855)	W.A.	S.A.	—	V.	N.S.W.	—	N.A.	B.fl.IV,46	M.fr.VI,227; VIII,246.			
V. perfoliata ; R. Brown, prodr. 581 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,46				
V. discophora ; F. v. M., fragm. X, 10 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,10.			
V. trinervis ; Labillardière, Nov. Holl. pl. spec. I, 54, t. 77 (1804)	W.A.	—	—	—	—	—	—	B.fl.IV,47	M.fr.XI,75.			
V. macrophylla ; Bentham, Fl. Austr. IV, 47 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,47	M.fr.XI,75.			
V. montana ; J. Hooker in Lond. Journ. of Bot. VI, 265 (1847)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,50				
V. pubescens ; R. Brown, prodr. 581 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,49	M.fr.VI,11; XI,75.			
V. spathulata ; R. Brown, prodr. 580 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,50	M.fr.XI,75.			
V. macrocalyx ; De Vriese, Goodenov. 176, t. 34 (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,49	M.fr.X,75.			
V. lyrata ; R. Brown, prodr. 580 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,49				
V. cycnopotamica ; F. v. M., fragm. VI, 7 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,48	M.fr.VI,249; X,10.			
V. paradoxa ; R. Brown, prodr. 580 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,48	M.fr.VI,11; XI,75; XII,23.			
V. macroplectra ; F. v. M., fragm. XII, 22 (1882)	W.A.	—	—	—	—	—	—	—				

SYNPETALEAE HYPOGYNAE.

F. v. M. in Woolls' pl. of the neighb. of Sydney, 34 (1880).

GENTIANEAE.

Necker in Act. Acad. Theod. Pal. II, 477 (1770), from B. de Jussieu (1759).—Genera 5, Species 24.

LIMNANTHEMUM ; Gmelin, Nov. Comm. Acad. Petrop. XIV, 257 (1770). (Villarsia, Liparophyllum.)

L. exaltatum ; F. v. M., fragm. IX, 165 (1875)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,376	M.fr.II,21; VI,139,254.			
L. parnassifolium ; F. v. M., fragm. IX, 165 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,376	M.fr.VI,140.			
L. violifolium ; F. v. M., Cens. 90 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,377	M.fr.VI,138.			
L. lasiospermum ; F. v. M., fragm. IX, 165 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,377	M.fr.VI,137.			
L. albiflorum ; F. v. M., Cens. 90 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,377	M.fr.VI,138.			
L. latifolium ; F. v. M., fragm. IX, 164 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,375	M.fr.VI,141.			
L. calthifolium ; F. v. M., fragm. IX, 164 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,374	M.fr.VI,140.			
L. congestiflorum ; F. v. M., Cens. 90 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,375	M.fr.VI,141.			
L. capitatum ; F. v. M., fragm. IX, 164 (1875)	W.A.	—	—	—	—	—	—	B.fl.IV,575	M.fr.VI,142.			
L. Gunnii ; J. Hooker, Fl. Tasman. II, 368 (1860) implied	—	—	T.	—	—	—	—	B.fl.IV,382	M.fr.VI,137; IX,164.			
L. exiguum ; F. v. M., fragm. I, 40 (1858)	—	—	T.	—	—	—	—	B.fl.IV,381	M.fr.VI,137.			
L. exiliflorum ; F. v. M., fragm. V, 46 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,381	M.fr.VI,137.			
L. hydrocharoides ; F. v. M. in Benth. Fl. Austr. IV, 380 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,380	M.fr.VI,139; IX,165.			
L. geminatum ; Grisebach, gen. et spec. Gent. 346 (1839)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,380	M.fr.VI,137,254.			
L. crenatum ; F. v. M. in Transact. phil. Soc. Vict. I, 17 (1854)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,379	M.fr.IV,127; VI,137; IX,165.			
L. minimum ; F. v. M., fragm. I, 40 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,379	M.fr.VI,137; IX,164.			
L. Moonii ; Thwaites, enum. pl. Ceyl. 205 (1862)	—	—	—	—	—	Q.	—	—	M.fr.IX,163	...	...	As., P.
L. Indicum ; Thwaites, enum. pl. Ceyl. 205 (1862)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,378	M.fr.IV,128; VII,136; IX,164.			

SEBAEA ; Solander in R. Brown, prodr. 451 (1810).

S. ovata ; R. Brown, prodr. 452 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,371	M.fr,VI,136; IX,165	...	N.Z.	
S. albidiflora ; F. v. M. in Transact. phil. Soc. Vict. I, 46 (1854)	—	S.A.	T.	V.	—	—	—	B.fl.IV,371	M.fr.VI,136.			

ERYTHRAEA ; Reneaulme, specim. hist. plant. 77, t. 76 (1611).

E. spicata ; Persoon, syn. plant. I, 283 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,371	M.fr.VI,136; IX,165; XI,20.	E., As., Af., P.		

CANSCORA; Lamarck, Encycl. méthod. I, 602 (1783). (Orthostemon.)

C. diffusa; R. Brown, prodr. 451 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,372	M.fr.VI,136; IX,165	...	As., Afr., P.

GENTIANA; Tournefort, inst. 80, t. 40 (1700), from Dioscorides.

G. quadrifaria; Blume, Bijdr. 804 (1825)	—	—	—	V.	N.S.W.	—	—	—	...	...	...	...	As.
G. saxosa; Forster in Svensk. Kongl. Vet. Ac. Handl. 183, t. 5 (1777)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,373	M.fr.VI,136,255; IX,165	..	Am, N.Z.

LOGANIACEAE.

R. Brown in App. Flind. Voy. 564 (1814).—Genera 7, Species 52.

STRYCHNOS; Linné, syst. nat. 8 (1735); spec. plant. 189 (1753).

S. lucida; R. Brown, prodr. 469 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,369	M.fr.IV,44	...	...	As.
S. psilosperma; F. v. M., fragm. IV, 44 (1863)	—	—	—	—	—	Q.	—	B.fl.IV,369	M.fr.VI,131			

FAGRAEA; Thunberg in Kongl. Svensk. Vetensk. Acad. Handl. 125 (1782).

F. racemosa; Jack in Roxb. Fl. Ind. ed. Carey. II, 35 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.IV,367	M.fr.II,137	..	...	As.

GARDNERIA; Wallich in Roxb. Fl. Indic. I, 400 (1820). (Loranthophyllum.)

G. fagraeacea; F. v. M., fragm. VI, 130 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,368	M.fr.VI,130.			

MITREOLA; Linné, hort. Cliffort. 492 (1737).

M. oldenlandioides; Wallich, numeric list. 4350 (1828)	—	—	—	—	—	—	N.A.	B.fl.IV,349	M.fr.I,130; VI,131.	..	As.

MITRASACME; Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804). (Mitragyne.)

M. Archeri; J. Hooker, Fl. Tasman. II, 368 (1860)	—	—	T.	—	—	—	—	B.fl.IV,351				
M. montana; J. Hooker, Fl. Tasman. I, 274, t. 88 (1859)	—	—	T.	V.	—	—	—	B.fl.IV,351	M.fr.VI,131.			
M. serpillifolia; R. Brown, prodr. 454 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,352	M.fr.I,130.			
M. paludosa; R. Brown, prodr. 453 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,352	M.fr.I,134.			
M. pilosa; Labillardière, Nov. Holl. pl. spec. I, 36, t. 49 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,352	M.fr.I,130; VI,131.			
M. alsinoides; R. Brown, prodr. 453 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,353				
M. polymorpha; R. Brown, prodr. 452 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,353	M.fr.VI,131.			
M. longiflora; F. v. M. in Benth. Fl. Austr. IV, 353 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,353				
M. elata; R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,354	M.fr.I,132.			
M. exserta; F. v. M., fragm. I, 131 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,354	M.fr.I,131.			
M. tenuiflora; Bentham, Fl. Austr. IV, 354 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,354				
M. ambigua; R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,354				
M. nudicaulis; Reinwardt in Blume, Bijdr. 849 (1826)	—	—	—	—	—	—	N.A.	B.fl.IV,355	M.fr.I,132.	..	...	As.
M. connata; R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,355	M.fr.I,131.			
M. laevis; Bentham in Journ. Linn. Soc. I, 93 (1857)	—	—	—	—	—	—	N.A.	B.fl.IV,355	M.fr.I,133.			
M. Indica; Wight, icon. pl. Ind. or. IV, t. 1601 (1850)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,356	...	...	..	As.
M. stellata; R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,356				
M. Cunninghamii; Bentham, Fl. Austr. IV, 357 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,357				
M. pygmaea; R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,357				
M. lutea; F. v. M., fragm. I, 133 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,357	M.fr.IV,170.			
M. multicaulis; R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,357				
M. laricifolia; R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,358	M.fr.VI,131.			
M. prolifera; R. Brown, prodr. 453 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,358				
M. gentianea; F. v. M., fragm. I, 130 (1859)	—	—	—	—	—	—	N.A.	B.fl.IV,358	M.fr.I,130.			
M. phascoides; R. Brown, prodr. 454 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,359				
M. paradoxa; R. Brown, prodr. 454 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,359	M.fr.I,130; VI,131.			
M. distylis; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	—	S.A.	T.	V.	—	—	—	B.fl.IV,359	M.fr.VI,131.			

GENIOSTOMA ; R. & G. Forster, char. gen. 23, t. 12 (1776).

G. Australianum ; F. v. M., fragm. V, 19 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,367	M.fr.V,19.							
G. petiolosum ; Moore & F. v. M., fragm. VII, 28 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,193.							

LOGANIA ; R. Brown, prodr. 454 (1810). (Euosma 1808, restorable.)

L. longifolia ; R. Brown, prodr. 456 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,361	M.fr.VI,132.							
L. latifolia ; R. Brown, prodr. 455 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,361	M.fr.VI,132.							
L. crassifolia ; R. Brown, prodr. 455 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,362	M.fr.VI,132.							
L. ovata ; R. Brown, prodr. 455 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,362	M.fr.VI,132.							
L. buxifolia ; F. v. M., fragm. VI, 132 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,362	M.fr.VI,132.							
L. stenophylla ; F. v. M., fragm. I, 128 (1859)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,362	M.fr.I,128.							
L. micrantha ; Bentham in Journ. Linn. Soc. I, 94 (1857)	W.A.	—	—	—	—	—	—	B.fl.IV,363	M.fr.I,129; VI,132.							
L. linifolia ; Schlechtendal, Linnaea XX, 605 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,363	M.fr.I,129; VI,131.							
L. fasciculata ; R. Brown, prodr. 456 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,363	M.fr.I,129; VI,131.							
L. floribunda ; R. Brown, prodr. 456 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,364	M.fr.VI,131.							
L. cordifolia ; Hooker in Mitch. Trop. Austr. 341 (1848)	—	—	—	—	—	Q.	—	B.fl.IV,364	M.fr.VI,132.							
L. nuda ; F. v. M., fragm. I, 129 (1859)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,365	M.fr.III,163; VI,131.							
L. spermacocea ; F. v. M., fragm. VI, 134 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,134.							
L. flaviflora ; F. v. M. in Vict. Naturalist V, 165 (1889)	W.A.	—	—	—	—	—	—	—								
L. callosa ; F. v. M., fragm. VI, 134 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.VI,134.							
L. campanulata ; R. Brown, prodr. 456 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,365	M.fr.I,130; VI,133.							
L. serpillifolia ; R. Brown, prodr. 456 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,366	M.fr.I,130; VI,133.							
L. pusilla ; R. Brown, prodr. 456 (4810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,366	M.fr.VI,131.							

PLANTAGINEAE.

A. L. de Jussieu, Gen. plant. 89 (1789).—Genus 1, Species 4.

PLANTAGO ; Tournefort, inst. 126, t. 48 (1700), from l'Ecluse (1576), from W. Turner (1538).

P. varia ; R. Brown, prodr. 424 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,139	M.fr.XI,20.							
P. stellaris ; F. v. M., fragm. II, 23 (1860)	—	—	—	V.	N.S.W.	—	—	—	M.fr.VIII,148.							
P. Brownii ; Rapin in Mém. de la Soc. Linn. Par. VI, 484 (1827)	—	—	T.	—	—	—	—	B.fl.V,141	M.fr.II,23	...	...	N.Z.				
P. Gunnii ; J. Hooker in Lond. Journ. V. 446, t. 13 (1846)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,142	M.fr.VIII,148; XI,134.							

PRIMULACEAE.

Ventenat, Tabl. du régn. végét. II, 285 (1799).—Genera 3, Species 6.

ANAGALLIS ; Tournefort, inst. 142, t. 59 (1700) from Hippocrates and Dioscorides. (Centunculus, Micropyxis.)

A. pumila ; Swartz, Fl. Ind. occ. I, 345 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.IV,270	...	...	...	As., Afr., Am.				
A. Centunculus ; Afzelius, de Vegetab. Suecan. (1785)	—	S.A.	—	V.	—	—	—	—	...	...	...	E., As., Afr., Am.				

LYSIMACHIA ; Tournefort, inst. 141, t. 59 (1700) from Dioscorides.

L. salicifolia ; F. v. M. in Benth. Fl. Austr. IV, 269 (1869)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,269								
L. Japonica ; Thunberg, Fl. Japon. 83 (1784)	—	—	—	—	N.S.W.	—	—	B.fl.IV,269	...	...	..	As.				

SAMOLUS ; Tournefort, inst. 143, t. 60 (1700). (Sheffieldia.)

S. Valerandi ; Linné, spec. plant. 171 (1753)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,271	...	...	...	E., As., Afr., Am.				
S. repens ; Persoon, synops. plant. I, 171 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,271	M.fr.XI,20	...	...	Afr., Am., P., N.Z.				

MYRSINACEAE.

R. Brown, prodr. 532 (1810). (Primulacearum subordo.)—Genera 5, Species 12.

MAESA; Forskael, Fl. Aegypt. Arab. 66 (1775). (Baeobotrys.)

M. dependens; F. v. M., fragm. V, 107 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,273	M.fr.V,107.					
M. haplobotrys; F. v. M., fragm. V, 161 (1866)	—	—	—	—	—	Q.	—	B.fl.IV,273	M.fr.V,161.					

SAMARA; Linné, mantiss. II, 144 (1771). (Choripetalum.)

S. Australiana; F. v. M., fragm. VI, 164 (1868)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,274	M.fr.III,36; IV,82,174.					

MYRSINE; Linné, syst. nat. 8 (1735); Linné, gen. plant. 54 (1737).

M. urceolata; R. Brown, prodr. 534 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,274						
M. campanulata; F. v. M., fragm. VI, 235 (1868)	—	—	—	—	—	Q.	—	—	M.fr.VI,235.					
M. crassifolia; R. Brown, prodr. 534 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,275	M.fr.IV,81; V,108; VI,164; VIII,48.					
M. platystigma; F. v. M., fragm. VIII, 48 (1873)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,48.					
M. variabilis; R. Brown, prodr. 534 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,275	M.fr.IV,81.					
M. achradifolia; F. v. M., fragm. VI, 164 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,275	M.fr.VI,164.					

ARDISIA; Swartz, nov. gen. et spec. 3 et 48 (1788). (Labisia partly.)

A. Pseudo-Jambosa; F. v. M., fragm. IV, 81 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,276	M.fr.IV,81.					
A. brevipedata; F. v. M., fragm VI, 163 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,276	M.fr.VI,163.					

AEGICERAS; Gaertner, de fruct. I, 216, t. 46 (1788).

A. majus; Gaertner, de fruct. I, 216, t. 46 (1788)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,277	...	...	...	As., P.		

SAPOTACEAE.

A. L. de Jussieu, Gen. plant. 151 (1789), from B. de Jussieu (1759).—Genera 6, Species 19.

NIEMEYERA; F. v. M., fragm. VII, 114 (1870). (Chrysophyllum partly.)

N. prunifera; F. v. M., fragm. VII, 114 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,278	M.fr.VI,26.					

AMORPHOSPERMUM; F. v. M., fragm. VII, 112 (1870). (Sersalisia partly.)

A. antilogum; F. v. M., fragm. VII, 113 (1870)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.X,119.					

LUCUMA; Molina, Saggio 186 (1782). (Sersalisia partly.)

L. sericea; Bentham & J. Hooker, gen. pl. II, 654 (1876)	—	—	—	—	—	Q.	N.A.	B.fl.IV,279	M.fr.V,162; VII,112.					
L. Galactoxylon; Bentham & J. Hooker, gen. pl. II, 654 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,279						

SIDEROXYLON; Dillenius, hort. Eltham. 357, t. 265 (1732). (Achras partly, Sapota partly, Sersalisia partly.)

S. Arnhemicum; Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	—	—	N.A.	B.fl.IV,280						
S. Pohlmanianum; Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,281	M.fr.V,184.					
S. xerocarpum; Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,281						
S. chartaceum; Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	—	Q.	—	B.fl.IV,281						
S. Richardi; F. v. M., Cens. of Austr. pl. 92 (1882)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,282						
S. Australe; Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,282	M.fr.V,162; IX,73.					
S. Brownii; F. v. M., Cens. of Austr. pl. 92 (1882)	—	—	—	—	—	Q.	—	B.fl.IV,283						
S. myrsinoides; Bentham & J. Hooker, gen. pl. II, 655 (1876)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,283	M.fr.IX,73.					
S. euphlebium; F. v. M., Cens. of Austr. pl. 92 (1882)	—	—	—	—	—	Q.	—	—	M.fr.VII,110.					
S. Brownlessianum; F. v. M., fragm. VIII, 111 (1870)	—	—	—	—	—	Q.	—	—	M.fr.VII,111.					
S. Howeanum; F. v. M., Cens. of Austr. pl. 92 (1882)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,72.					
S. costatum; F. v. M., Cens. 92 (1882) from Endlicher (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,72.					

HORMOGYNE; A. de Candolle, prodr. VIII, 176 (1844).

H. cotinifolia; A. De Candolle, prodr. VIII, 176 (1844)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,284	M.fr.V,161; VII,111.					

MIMUSOPS; Linné, Fl. Zeil. 57 (1747).

M. parvifolia; R. Brown, prodr. 531 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,284	M.fr.V,162; VI,178; VII,114.					
M. Kauki; Linné, spec. plant. 349 (1753)	—	—	—	—	—	Q.	—	B.fl.IV,285	M.fr.V,162	...	...	As.		

EBENACEAE.

VENTENAT, Tabl. II, 443 (1799).—Genera 2, Species 15.

DIOSPYROS; Linné, gen. plant. 143 (1737). (Cargillia.)

D. cordifolia; Roxburgh, pl. Corom. I, 38, t. 50 (1795)	—	—	—	—	—	—	—	—	—	—	N.A.	B.fl.IV,286	...	...	As.
D. hebecarpa; Cunningham in Benth. Fl. Austr. IV, 286 (1869)	—	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl.IV,286			
D. maritima; Blume, Bijdrag. 669 (1825)	—	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl.IV,287	M.fr.V,163; X,75	..	As.
D. mabacea; F. v. M., Docum. intercol. Exhib. 35 (1866)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,288	M.fr.V,162; X,75.		
D. Cargillia; F. v. M., Docum. intercol. Exhib. 35 (1866)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,288	M.fr.IV,82; V,162; X,74.		
D. pentamera; F. v. M. & Woolls, Doc. intercol. Exhib. 35 (1866)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,288	M.fr.IV,82; V,163; X,75.		

MABA; R. & G. Forster, char. gen. 121, t. 61 (1776).

M. laurina; R. Brown, prodr. 527 (1810)	—	—	—	—	—	—	—	—	—	Q.	—	B.fl.IV,289			
M. rufa; Labillardière, sert. Austr. Caled. 33, t. 36 (1824)	—	—	—	—	—	—	—	—	—	Q.	—	B.fl.IV,289	M.fr.V,164; VI,253; X,74	...	P.
M. hemicycloides; F. v. M. in Benth. Fl. Austr. IV, 290 (1869)	—	—	—	—	—	—	—	—	—	Q.	—	B.fl.IV,290			
M. fasciculosa; F. v. M., fragm. V, 163 (1866)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,290	M.fr.X,74.		
M. compacta; R. Brown, prodr. 528 (1810)	—	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl.IV,290	M.fr.V,165.		
M. reticulata; R. Brown, prodr. 528 (1810)	—	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl.IV,291			
M. geminata; R. Brown, prodr. 527 (1810)	—	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl.IV,291	M.fr.V,164.		
M. humilis; R. Brown, prodr. 527 (1810)	—	—	—	—	—	—	—	—	—	Q.	N.A.	B.fl.IV,291	M.fr.V,164; X,74.		
M. buxifolia; Persoon, synops. pl. II, 602 (1807)	—	—	—	—	—	—	—	—	—	—	N.A.	—	M.fr.X,74	...	As., Afr.

AQUIFOLIACEAE.

DE CANDOLLE, Théor. élém. I, 217 (1813).—Genera 2, Species 2.

BRYONIA; Endlicher in den Ann. des Wien. Mus. I, 184 (1836).

B. Arnhemensis; F. v. M., fragm. II, 119 (1861)	—	—	—	—	—	—	—	—	—	—	N.A.	B.fl.I,397	M.fr.II,119.		

ILEX; Linné, gen. plant. 33 (1737) from C. Bauhin (1623).

I. peduncularis; F. v. M., fragm. VII, 105 (1871)	—	—	—	—	—	—	—	—	—	Q.	—	—	M.fr.VII,105.		

STYRACEAE.

L. C. RICHARD, Analyse du fruit, 48 (1808).—Genus 1, Species 2.

SYMPLOCOS; N. J. Jacquin, enum. plant. Carib. 24 (1760).

S. Stawellii; F. v. M., fragm. V, 60 (1865)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,292	M.fr.V,60.		
S. Thwaitesii; F. v. M., fragm. III, 22 (1862)	—	—	—	—	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,293	M.fr.V,211.		

JASMINEAE.

A. L. DE JUSSIEU, Gen. plant. 104 (1789) from Necker (1770).—Genera 5, Species 21.

JASMINUM; Tournefort, inst. 597, t. 368 (1700), from l'Ecluse (1611), from W. Turner (1538).

J. didymum; G. Forster, prodr. 3 (1786)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,294	M.fr.IV,150; VI,87; VIII,43.						As., P.
J. racemosum; F. v. M., fragm. I, 19 (1858)	—	—	—	—	—	Q.	—	B.fl.IV,295	M.fr.VI,87; VIII,43						
J. lineare; R. Brown, prodr. 521 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,295	M.fr.VI,87; VIII,43.						
J. simplicifolium; G. Forster, prodr. 3 (1786)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,296	M.fr.VI,87; VIII,43					...	As., P.
J. aemulum; R. Brown, prodr. 521 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,296	M.fr.VI,86.						
J. calcareum; F. v. M., fragm. I, 212 (1859)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,297	M.fr.VI,87; VIII,43.						
J. suavissimum; Lindley in Mitch. Trop. Austr. 355 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,297	M.fr.I,183; II,181; VI,87.						
J. singuliflorum; F. v. M. & Bailey in suppl. to Queensl. Fl. 38 (1886)	—	—	—	—	—	Q.	—	—							

OLEA; Tournefort, inst. 598, t. 370 (1700) from Plinius.

O. paniculata; R. Brown, prodr. 523 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,297	M.fr.IV,83; VIII,43					..	P.
O. Endlicheri; F. v. M., fragm. VIII, 43 (1872)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,43; IX,169.						

LIGUSTRUM; Tournefort. inst. 596, t. 367 (1700) from C. Bauhin (1623).

L. Australianum; F. v. M., fragm. V, 20 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,298	M.fr.V,20.			

NOTELAEA; Ventenat, Choix des pl. dans le jard. de Cels. t. 25 (1803).

N. ovata; R. Brown, prodr. 524 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,299	M.fr.VIII,43.			
N. longifolia; Ventenat, Choix t. 25 (1803)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,299	M.fr.VIII,43.			
N. punctata; R. Brown, prodr. 524 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,300				
N. microcarpa; R. Brown, prodr. 524 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,300	M.fr.VIII,43.			
N. ligustrina; Ventenat, Choix t. 25 (1803)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,300	M.fr.VIII,43,142.			
N. linearis; Bentham, Fl. Austr. IV, 300 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,300				

MAYEPEA; Aublet, Hist. des pl. de la Guian. I, 81, t. 31 (1775). (Chionanthus partly, Linociera partly, Ceranthes.)

M. ramiflora; F. v. M., Cens. 92 (1882) from Roxburgh (1820)	—	—	—	—	—	Q.	—	B.fl.IV,301	M.fr.IV,83; VIII,42.			
M. picrophloia; F. v. M., Cens. of Austr. plant. 92 (1882)...	—	—	—	—	—	Q.	—	B.fl.IV,301	M.fr.III,139,24; IV,176.			
M. axillaris; F. v. M., Cens. 92 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,301	M.fr.VIII,42.			
M. quadristaminea; F. v. M., fragm. X, 89 (1876)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,41.			

APOCYNEAE.

A. L. de Jussieu, Gen. plant. 143 (1789).—Genera 13, Species 47.

CHILOCARPUS; Blume, catal. hort. Buitenz. (1823).

C. Australis; F. v. M., fragm. II, 90 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,303	M.fr.VI,118.			

MELODINUS; R. & G. Forster, char. gen. 37, t. 19 (1776).

M. acutiflorus; F. v. M. in Transact. phil. Soc. Vict. II, 71 (1857)	—	—	—	—	—	Q.	—	B.fl.IV,304				
M. Guilfoylei; F. v. M., fragm. VI, 118 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,304	M.fr.VI,118.			
M. Baueri; Endlicher, prodr. pl. Norfolk. 59 (1833)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.			

CARISSA; Linné, mantiss, pl. 7 (1767).

C. laxiflora; Bentham, Fl. Austr. IV, 305 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,305				
C. Brownii; F. v. M., fragm. IV, 45 (1863)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,305	M.fr.IV,45.			

CERBERA; Linné, gen. plant. 62 (1737).

C. Odollam; Gaertner, de fruct. 193, t. 124 (1791)	—	—	—	—	—	Q.	N.A.	B.fl.IV,306	M.fr.VI,117	...	As.,	P.

ALYXIA; Banks in R. Brown, prodr. 469 (1810). (Gynopogon, 1776.)

A. buxifolia; R. Brown, prodr. 470 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,307	M.fr.VI,117; VII,131.			
A. ruscifolia; R. Brown, prodr. 470 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,308	M.fr.VI,117; VII,131.			P.
A. ilicifolia; F. v. M., fragm. IV, 149 (1864)...	—	—	—	—	—	Q.	—	B.fl.IV,308	M.fr.IV,149.			
A. squamulosa; C. Moore & F. v. M., fragm. VIII, 47 (1873)	—	—	—	—	N.S.W.	—	—	—	M.fr.VI,117; VIII,47.			
A. obtusifolia; R. Brown, prodr. 470 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,308				
A. Gynopogon; Roemer & Schultes, syst. veg. IV, 440 (1819)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,47.			
A. spicata; R. Brown, prodr. 470 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,308	M.fr.VI,117.			
A. Lindii; F. v. M., fragm. VIII, 46 (1873)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,46.			
A. thyrsiflora; Bentham, Fl. Austr. IV, 309 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,309				
A. Thozetii; F. v. M., fragm. X, 103 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,103.			

OCHROSIA; A. L. de Jussieu, gen. plant. 143 (1789). (Lactaria.)

O. elliptica; Labillardière, sert. Austr. Caled. 27, t. 30 (1824)	—	—	—	—	—	Q.	—	B.fl.IV,310	M.fr.III,110; VI,118.		As.,	P.
O. Moorei; F. v. M. in Benth. Fl. Austr. IV, 310 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,310	M.fr.III,110; VI,118; VII,131.			
O. Kilneri; F. v. M., fragm. VII, 129 (1871)...	—	—	—	—	—	Q.	—	—	M.fr.VIII,129.			

NOTONERIUM ; Bentham in B. & J. H. gen. II, 698 (1876).

N. Gossei ; Bentham in B. & J. H. gen. pl. 698 (1876)	—	S.A.	—	—	—	—	—	—	M.fr.X,103.					

TABERNAEMONTANA ; Plumier, nov. pl. Amer. gen. 18, t. 30 (1703).

T. orientalis ; R. Brown, prodr. 468 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,311	M.fr.VI,117				...	P.
T. pubescens ; R. Brown, prodr. 468 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,311	M.fr.VI,118				...	P.

ALSTONIA ; R. Brown in Mem. Wern. Soc. I, 75 (1809).

A. scholaris ; R. Brown in Mem. Wern. Soc. I, 75 (1809)	—	—	—	—	—	Q.	—	B.fl.IV,312	M.fr.VI,117				...	As., Afr.
A. verticillosa ; F. v. M., fragm. VI, 116 (1868)	—	—	—	—	—	Q.	N.A.	B.fl.IV,313	M.fr.VI,116.					
A. ophioxyloides ; F. v. M., fragm. I, 57 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,313	M.fr.VI,117.					
A. villosa ; Blume, Bijdr. 1038 (1826)	—	—	—	—	—	Q.	—	B.fl.IV,313	M.fr.VI,117				...	As.
A. linearis ; Bentham, Fl. Austr. IV, 314 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,314						
A. constricta ; F. v. M., fragm. I, 57 (1858)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,314	M.fr.III,163; IV,170; VI,117; VII,131.					

ICHNOCARPUS ; R. Brown in Mem. Wern. Soc. I, 75 (1809).

I. frutescens ; R. Brown in Mem. Wern. Soc. I, 75 (1809)	—	—	—	—	—	Q.	N.A.	B.fl.IV,315	M.fr.VI,118				...	As.

WRIGHTIA ; R. Brown in Mem. Wern. Soc. I, 75 (1809). (Balfouria.)

W. pubescens ; R. Brown, prodr. 467 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,316	...	...	...	...	...	As., P.
W. saligna ; F. v. M. in Benth. Fl. Austr. IV, 316 (1869)	—	—	—	—	—	Q.	N.A.	B.fl.IV,316	M.fr.V,161; VI,118.					
W. Cunninghamii ; Bentham, Fl. Austr. IV, 317 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,317						

PARSONIA ; R. Brown in Mem. Wern. Soc. I, 64 (1809). (Lyonsiae subgenus.)

P. lanceolata ; R. Brown, prodr. 466 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,318	M.fr.VI,126.					
P. velutina ; R. Brown, prodr. 466 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,318	M.fr.VI,130.					
P. Leichhardtii ; F. v. M., fragm. VI, 128 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,319	M.fr.VI,128.					
P. ventricosa ; F. v. M. in Transact. phil. Inst. Vict. II, 71 (1857)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,319	M.fr.VI,130.					

LYONSIA ; R. Brown in Mem. Wern. Soc. I, 66 (1809).

L. lilacina ; F. v. M. in Bentham, Fl. Austr. IV, 321 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,321	M.fr.VI,127.					
L. induplicata ; F. v. M. in Bentham, Fl. Austr. IV, 321 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,321	M.fr.VI,129.					
L. straminea ; R. Brown, prodr. 466 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,321	M.fr.VI,129.					
L. reticulata ; F. v. M., Rep. Burdek. Exped. 16 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,321	M.fr.VI,129.					
L. diaphanophleba ; F. v. M. in Benth. Fl. Austr. IV, 322 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,322	M.fr.II,158; VI,130.					
L. Langiana ; F. v. M., fragm. VI, 128 (1868)	—	—	—	—	—	Q.	—	B.fl.IV,322	M.fr.VI,128.					
L. largiflorens ; F. v. M. in Benth. Fl. Austr. IV, 322 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,322						
L. latifolia ; Bentham, Fl. Austr. IV, 323 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,323						
L. oblongifolia ; Bentham, Fl. Austr. IV, 323 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,323						
L. eucalyptifolia ; F. v. M. in Benth. Fl. Austr. IV, 323 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,323	M.fr.II,159; VI,130.					

ASCLEPIADEAE.

N. J. Jacquin, Misc. Austr. I, 35 (1778).—Genera 15, Species 61.

GYMNANTHERA ; R. Brown in Mem. Wern. Soc. I, 58 (1809).

G. nitida ; R. Brown, prodr. 464 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,326	M.fr.V,160	...	...			P.

SECAMONE ; R. Brown in Mem. Wern. Soc. I, 55 (1809).

S. elliptica ; R. Brown, prodr. 464 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,327	M.fr.V,161	...	...			As.
S. ovata ; R. Brown, prodr. 464 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,328						

VINCETOXICUM ; Dodoens, stirp. hist. pemptades, 704 (1583). (Oxystelma partly, Cynoctonum, Cynanchi subgenus.)

V. ovatum ; Bentham, Fl. Austr. IV, 330 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,330						
V. elegans ; Bentham, Fl. Austr. IV, 330 (1869)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,330	M.fr.V,160.					

V. carnosum ; Bentham, Fl. Austr. IV, 331 (1869) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,331						
V. leptolepis ; Bentham, Fl. Austr. IV, 331 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,331						
CYNANCHUM ; Linné, gen. plant. 63 (1737). (Endotropis.)														
C. erubescens ; R. Brown, prodr. 463 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,332						
C. floribundum ; R. Brown, prodr. 463 (1810) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,332	M.fr.V,160.					
C. pedunculatum ; R. Brown, prodr. 463 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,333	M.fr.V,160.					
C. puberulum ; F. v. M. in Benth. Fl. Austr. IV, 333 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,333						
SARCOSTEMMA ; R. Brown in Mem. Wern. Soc. I, 50 (1809).														
S. Australe ; R. Brown, prodr. 463 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,328	M.fr.V,160; XI,80.					
DAEMIA ; R. Brown in Mem. Wern. Soc. I, 50 (1809). (Pentatropis, Rhyncharrhena.)														
D. linearis ; F. v. M., Cens. 94 (1882) from Decaisne (1833) ...	W.A.	—	—	—	—	—	—	B.fl.IV,329						
D. atropurpurea ; F. v. M., Cens. 94 (1882) ...	—	—	—	—	—	Q.	—	B.fl.IV,329	M.fr.I,128.					
D. quinquepartita ; F. v. M., Cens. 94 (1882)...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,329	M.fr.I,128; XI,80.					
D. Kempeana ; F. v. M. in Wing's S. Sc. Record, Aug. (1882) ...	—	S.A.	—	—	—	—	—	—						
GYMNEMA ; R. Brown in Mem. Wern. Soc. I, 33 (1809). (Bidaria.)														
G. sylvestre ; R. Brown in Mem. Wern. Soc. I, 33 (1809) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,343	...	...	...	As.		
G. pleiadenium ; F. v. M., fragm. XI, 77 (1879) ...	—	—	—	—	—	Q.	—	—	M.fr.XI,77.					
G. Muelleri ; Bentham, Fl. Austr. IV, 343 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,343						
G. brevifolium ; Bentham, Fl. Austr. IV, 343 (1809) ...	—	—	—	—	—	Q.	—	B.fl.IV,343						
G. trinerve ; R. Brown, prodr. 462 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,343						
G. stenophyllum ; A. Gray in Proc. Am. Ac. of Sc. V, 335 (1864) ...	—	—	—	—	—	—	N.A.	B.fl.IV,344	...	...	...	P.		
GONGRONEMA ; Decaisne in De Candolle, prodr. VIII, 624 (1844).														
G. micradenia ; Bentham & J. Hooker, gen. plant. II, 770 (1876)...	—	—	—	—	—	Q.	—	B.fl.IV,344	M.fr.XI,78.					
TYLOPHORA ; R. Brown in Mem. Wern. Soc. I, 28 (1809). (Hybanthera.)														
T. erecta ; F. v. M. in Benth. Fl. Austr. IV, 334 (1869) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,334	M.fr.IX,71; XI,79.					
T. macrophylla ; Bentham, Fl. Austr. IV, 334 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,334						
T. grandiflora ; R. Brown, prodr. 460 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,334	M.fr.IX,71; XI,80.					
T. floribunda ; Bentham, Fl. Austr. IV, 335 (1869) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,335						
T. calcarata ; Bentham, Fl. Austr. IV, 335 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,335						
T. barbata ; R. Brown, prodr. 460 (1810) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,335	M.fr.V,160; XI,79.					
T. Woollsii ; Bentham, Fl. Austr. IV, 335 (1869) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,335						
T. paniculata ; R. Brown, prodr. 460 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,336	M.fr.IX,71.					
T. enervis ; F. v. M., fragm. IX, 70 (1875) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,70.					
T. flexuosa ; R. Brown, prodr. 460 (1810) ...	—	—	—	—	—	—	N.A.	B.fl.IV,336	M.fr.XI,79.					
T. biglandulosa ; A. Gray in Proc. of Am. acad. sc. vol. V (1864)...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.					
MARSDENIA ; R. Brown in Mem. Wern. Soc. I, 28 (1809). (Leichhardtia, Stephanotis.)														
M. cinerascens ; R. Brown, prodr. 461 (1810)...	—	—	—	—	—	Q.	N.A.	B.fl.IV,337						
M. rhyncholepis ; F. v. M., fragm. XI, 78 (1879) ...	—	—	—	—	—	Q.	—	—	M.fr.XI,78.					
M. flavescens ; Cunningham in Bot. Mag. t. 3289 (1833) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,337	M.fr.V,160.					
M. cymulosa ; Bentham, Fl. Austr. IV, 338 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,338						
M. velutina ; R. Brown, prodr. 461 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,338						
M. Hullsii ; F. v. M. in Benth. Fl. Austr. IV, 338 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,338						
M. araujacea ; F. v. M., fragm. VI, 135 (1868) ...	—	—	—	—	—	Q.	—	B.fl.IV,339	M.fr.VI,135.					
M. tubulosa ; F. v. M., fragm. IX, 71 (1875) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,71.					
M. rostrata ; R. Brown, prodr. 461 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,339	M.fr.V,160; VI,135; IX,71.					
M. Fraseri ; Bentham, Fl. Austr. IV, 339 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,339						
M. longiloba ; Bentham, Fl. Austr. IV, 340 (1869) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,340	M.fr.XI,79.					

M. suaveolens ; R. Brown, prodr. 461 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,340							
M. leptophylla ; F. v. M. in Benth. Fl. Austr. IV, 340 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,340							
M. Leichhardtiana ; F. v. M., fragm. V, 160 (1866)...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,341	M.fr.V,160.						
M. viridiflora ; R. Brown, prodr. 461 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,341	M.fr.V,160.						
M. coronata ; Bentham, Fl. Austr. IV, 341 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,341							
M. microlepis ; Bentham, Fl. Austr. IV, 342 (1869)...	—	—	—	—	—	Q.	—	B.fl.IV,342							
THOZETIA ; F. v. M. in Benth. Fl. Austr. IV, 347 (1869).															
T. racemosa ; F. v. M. in Benth. Fl. Austr. IV, 347 (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,347							
HOYA ; R. Brown in Mem. Wern. Soc. I, 26 (1809).															
H. carnosa ; R. Brown in Mem. Wern. Soc. I, 26 (1809) ...	—	—	—	—	—	Q.	—	B.fl.IV,346	...						As.
H. Australis ; R. Brown in Transact. hort. Soc. VII, 28 (1826) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,346	M.fr.V,159; XI,80					...	P.
H. Nicholsoniae ; F. v. M., fragm. V, 159 (1886) ...	—	—	—	—	—	Q.	—	B.fl.IV,347	M.fr.IX,71,190.						
DISCHIDIA ; R. Brown in Mem. Wern. Soc. I, 32 (1809).															
D. nummularia ; R. Brown, prodr. 461 (1810)...	—	—	—	—	—	Q.	—	B.fl.IV,345	M.fr.V,161 ...					...	As., P.
D. Timorensis ; Decaisne in nouv. annal. du Mus. III, 377, t. 17 (1835) ...	—	—	—	—	—	Q.	N.A.	—	...					...	As.
D. Rafflesiana ; Wallich, pl. Asiat. rar. II, 35, t. 142 (1831) ...	—	—	—	—	—	Q.	N.A.	—	...					...	As.
MICROSTEMMA ; R. Brown in Mem. Wern. Soc. I, 25 (1809).															
M. tuberosum ; R. Brown, prodr. 459 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,345	M.fr.I,58.						
M. glabriflorum ; F. v. M., fragm. I, 58 (1858) ...	—	—	—	—	—	—	N.A.	B.fl.IV,345	M.fr.I,58.						
CEROPEGIA ; Linné, gen. plant. 65 (1737).															
C. Cumingiana ; Decaisne in De Candolle, prodr. VIII, 642 (1844) ...	—	—	—	—	—	Q.	—	B.fl.IV,348	M.fr.V,159 ...					...	As.

CONVOLVULACEAE.

A. L. de Jussieu, Gen. plant. 132 (1789), from B. de Jussieu (1759).—Genera 13, Species 70.

ERYCIBE ; Roxburgh, pl. Corom. III, 31, t. 159 (1798).															
E. paniculata ; Roxburgh, pl. Corom. III, 31, t. 159 (1798) ...	—	—	—	—	—	Q.	—	B.fl.IV,411	M.fr.VI,100 ...					...	As.
LEPISTEMON ; Blume, Bijdrag. 722 (1825).															
L. urceolatus ; F. v. M.; L. Fitzalani, F. v. M., fragm. X, 111 (1877) ...	—	—	—	—	—	Q.	—	B.fl.IV,427	M.fr.X,111 ...					...	P.
L. Lucae ; F. v. M. in Vict. Naturalist, Oct. (1885)...	—	—	—	—	—	Q.	—								
IPOMOEA ; Linné, syst. nat. 8 (1735); Linné, gen. plant. 47 (1737). (Pharbitis, Batatas, Calonyction, Aniseia, Skinnera.)															
I. digitata ; Linné, syst. nat. edit. decim. n. 5 (1758) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,414	M.fr.VI,98; X,111					...	As., Afr., Am., P.
I. Davenporti ; F. v. M., fragm. VI, 97 (1868) ...	—	S.A.	—	—	—	—	N.A.	B.fl.IV,415	M.fr.VII,254; VIII,17; X,117.						
I. bona nox ; Linné, spec. plant. edit. secund. 228 (1762) ...	—	—	—	—	N.S.W.	Q.	—	—	M.fr.IX,74; X,73,113.						As., Am., P.
I. palmata ; Forskael, Fl. Aegypt. Arabic. 43 (1775) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,415	M.fr.VI,98; X,111					...	As., Afr., Am., P., [N.Z.
I. quinata ; R. Brown, prodr. 486 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,415	M.fr.VI,98; X,112					...	As., P.
I. diversifolia ; R. Brown, prodr. 487 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,416	M.fr.X,112.						
I. dissecta ; Willdenow, phytogr. 5, t. 2 (1794) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,416	M.fr.VI,98; X,112					...	As., Afr., P.
I. sinuata ; Ortega, decad. nov. pl. hort. Matrit. VII, 84 (1789) ...	—	—	—	—	—	Q.	—	—	M.fr.VIII,17; X,113					...	As., Afr., Am., P.
I. hederacea ; N. J. Jacquin, Collectan. I, 124 (1776) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,416	M.fr.VI,99; X,112					...	As., Afr., Am., P.
I. congesta ; R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	Q.	—	B.fl.IV,417	M.fr.VI,99; X,112					...	P.
I. purpurea ; Roth, Catalect. bot. I, 36 (1797) ...	—	—	—	—	—	Q.	—	B.fl.IV,417	M.fr.VI,99; X,112					...	Am.
I. peltata ; Choisy, Convolvulac. orient. 70 (1834) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,418	...					...	As., Afr., P.
I. Calobra ; Hill & F. v. M., fragm. XI, 73 (1879) ...	—	—	—	—	—	Q.	—	—	M.fr.XI,73,137.						
I. alata ; R. Brown, prodr. 484 (1810) ...	—	—	—	—	—	—	N.A.	B.fl.IV,418	M.fr.X,112 ...					..	Am.
I. Turpethum ; R. Brown, prodr. 485 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,418	M.fr.VI,98 ...					...	As., Afr., Am., P.

I. grandiflora; Lamarck, illustr. des. genr. I, 407 (1791)	—	—	—	—	—	Q.	N.A.	B.fl.IV,418	M.fr.VI,98; IX,74; X,112.		As., Afr., P.
I. costata; F. v. M. in Benth. Fl. Austr. IV, 419 (1869)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,419	M.fr.X,112.		
I. Pes Caprae; Roth. nov. spec. plant. 109 (1821)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,419	M.fr.VI,97; X,112	...	As., Afr., Am., P.
I. carnosa; R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,420	...	...	As., Afr., Am., P.
I. aquatica; Forskael, fl. Aeg. Arab. 44 (1775)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,420	M.fr.VI,97 ...	...	As., Afr.
I. gramines; R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421			
I. velutina; R. Brown, prodr. 485 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421			
I. abrupta; R. Brown, prodr. 485 1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421	M.fr.X,112.		
I. denticulata; Choisy in De Candolle, prodr. IX, 379 (1845)	—	—	—	—	—	Q.	N.A.	B.fl.IV,421	M.fr.I,185; X,112	...	As., P.
I. gracilis; R. Brown, prodr. 484 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,422	M.fr.I,185; X,112.		
I. sepiaria; Koenig in Roxb. Fl. Indic. ed. Carey II, 90 (1824)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,422	...	...	As., P.
I. Muelleri; Bentham, Fl. Austr. IV, 423 (1869)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,423	M.fr.X,112.		
I. cymosa; Roemer & Schultes, syst. veg. IV, 241 (1819)	—	—	—	—	—	Q.	—	B.fl.IV,423	...	...	As., Afr., P.
I. linifolia; Blume, Bijdrag. 721 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,423	...	...	As.
I. chryseides; Ker in Bot. Regist. t. 270 (1818)	—	—	—	—	—	Q.	—	B.fl.IV,423	M.fr.VI,98 ...	...	As., Afr.
I. flava; F. v. M. in Benth. Fl. Austr. IV, 424 (1869)	—	—	—	—	—	Q.	N.A.	B.fl.IV,424			
I. obscura; Ker in Bot. Regist. t. 239 (1817)	—	—	—	—	—	Q.	—	B.fl.IV,424	...	...	As., Afr.
I. incisa; R. Brown, prodr. 486 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,424	M.fr.I,186.		
I. uniflora; Roemer & Schultes, syst. IV, 247 (1819)	—	—	—	—	—	Q.	—	B.fl.IV,425	M.fr.VI,99 ...	...	As., Afr., P.
I. angustiflora; N. J. Jacquin, Collect. II, 367 (1788)	—	—	—	—	—	Q.	N.A.	B.fl.IV,425	M.fr.VI,97; X,112	...	As., Afr., P.
I. plebeja; R. Brown, prodr. 484 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,426	M.fr.VI,99; X,112.		
I. eriocarpa; R. Brown, prodr. 484 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,426	M.fr.VI,98; X,112	...	As., Afr., P.
I. heterophylla; R. Brown, prodr. 487 (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,426	M.fr.VI,98; X,112.		
I. erecta; R. Brown, prodr. 487 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,427	M.fr.VI,98.		
I. cataractae; Endl. prod. fl. ins. Norf. 53 (1833)	—	—	—	—	N.S.W.	Q.	—	—			
ARGYREIA; Loureiro, Fl. Cochinch. I, 134 (1790).											
A. speciosa; Sweet, hort. Brit. sec. ed. 373 (1830)	—	—	—	—	—	Q.	—	—	...	...	As.
CONVOLVULUS; Tournefort, inst. 82, t. 17 (1700), from W. Turner (1538). (Calystegia.)											
C. erubescens; Sims, Bot. Magaz. t. 1067 (1808)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,429	M.fr.VI,99; X,113; XI,73.		
C. parviflorus; Vahl, symbol. bot. III, 29 (1794)	—	—	—	—	—	Q.	N.A.	B.fl.IV,430	M.fr.VI,99; X,113	...	As., P.
C. marginatus; Poiret, Encycl. méth. suppl. II (1811)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,430	M.fr.VI,99; X,113	...	N.Z.
C. sepium; Linné, spec. plant. 153 (1753)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,430	M.fr.VI,99; X,113.		E., As., Afr., Am., N.Z.
POLYMERIA; R. Brown, prodr. 488 (1810).											
P. marginata; Bentham, Fl. Austr. IV, 432 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,432			
P. longifolia; Lindley in Mitch. Trop. Austr. 398 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,432	M.fr.X,114.		
P. angusta; F. v. M., fragm. VI, 100 (1868)	—	S.A.	—	—	—	—	N.A.	B.fl.IV,432	M.fr.VI,100.		
P. distigma; Bentham, Fl. Austr. IV, 433 (1869)	—	—	—	—	—	—	N.A	B.fl.IV,433			
P. occidentalis; F. v. M., inedit.	W.A.	—	—	—	—	—	—	B.fl.IV,433			
P. calycina; R. Brown, prodr. 488 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,433	M.fr.VI,100.		
P. ambigua; R. Brown, prodr. 488 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,433			
PORANA; Burmann, Fl. Ind. 51, t. 21 (1768). (Duperreya.)											
P. sericea; F. v. M., fragm. VI, 100 (1868)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,434	M.fr.II,22.		
BREWERIA; R. Brown, prodr. 487 (1810).											
B. linearis; R. Brown, prodr. 488 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,435			
B. media; R. Brown, prodr. 488 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,436	M.fr.X,113.		
B. brevifolia; Bentham, Fl. Austr. IV, 436 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,436			
B. rosea; F. v. M., fragm. I, 233 (1858)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,436	M.fr.VI,100; X,113.		
B. pannosa; R. Brown, prodr. 488 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,436	M.fr.I,233; VI,100; X,113.		

EVOLVULUS ; Linné, spec. plant. ed. sec. 391 (1762).

E. linifolius ; Linné, spec. plant. ed. sec. 392 (1762)	W.A.	S.A.	—	—	N.S.W.	Q. N.A.	B.fl.IV,437	M.fr.VI,100; X,113.	As., Afr., Am., P.

DICHONDRA ; R. & G. Forster, char. gen. 39, t. 20 (1776).

D. repens ; R. &. G. Forster, char. gen. 39, t. 20 (1776)	W.A.	S.A.	T.	V.	N.S.W.	Q. N.A.	B.fl.IV,438	M.fr.VI,100; X,113.	E., As., Afr., Am.

CRESSA ; Linné, Amoen. Acad. I, 121 (1747).

C. Cretica ; Linné, spec. plant. 223 (1753)	W.A.	S.A.	—	V.	N.S.W	Q. N.A.	B.fl.IV,437	M.fr.VI,100	E., As., Afr., Am.

WILSONIA ; R. Brown, prodr. 490 (1810).

W. humilis ; R. Brown, prodr. 490 (1810)	W.A.	S.A.	T.	V.	N.S.W.	— —	B.fl.IV,439	M.fr.VI,101; X,113.	
W. rotundifolia ; Hooker, icon. plant. t. 410 (1842)	W.A.	S.A.	—	V.	N.S.W.	— —	B.fl.IV,440	M.fr.VI,101.	
W. Backhousi ; J. Hooker in Lond. Journ. VI, 275 (1847)	W.A.	S A.	T.	V.	N.S.W.	— —	B.fl.IV,440	M.fr.VI,101; VIII,17.	

CUSCUTA ; Tournefort, inst. app. 652, t. 422 (1700), from C. Banhin (1623), from Bock (1539).

C. Chinensis ; Lamarck, Encycl. méth. II, 229 (1786)	—	—	—	—	—	Q. —	B.fl.IV,441	M.fr.X,113	As., Afr.
C. Australis ; R. Brown, prodr. 491 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q. —	B.fl.IV,441	M.fr.I,100; X,113	As., Afr., Am.
C. Tasmanica ; Engelmann in Transact. Acad. St. Louis I, 512 (1860)	—	S.A.	T.	V.	—	— —	B.fl.IV,441		

HYDROPHYLLEAE.

R. Brown in Bot. Regist. t. 242 (1817).—Genus 1, Species 2.

HYDROLEA ; Linné, spec. plant. ed. sec. 328 (1762).

H. Zeilanica ; Vahl, sym. bot. II, 46 (1791)	—	—	—	—	—	Q. N.A.	B.fl.IV,382		As., Afr., Am.
H. spinosa ; Linné, spec. plant. ed. sec. 328 (1762)	—	—	—	—	—	Q. —	B.fl.IV,383		Am.

SOLANACEAE.

Haller, enum. stirp. Helv. praef. 34 (1742).—Genera 9, Species 79.

PHYSALIS ; Linné, syst. nat. 8 (1735); gen. plant. 51 (1737).

P. minima ; Linné, spec. plant. 183 (1753)	—	—	—	—	N.S.W.	Q. N.A.	B.fl.IV,466	M.fr.VI,144	As., Afr., Am., P.

SOLANUM ; Tournefort, inst. 148, t. 62 (1700) from Celsus. [N.Z.

S. nigrum ; Linné, spec. plant. 186 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q. N.A.	B.fl.IV,446	M.fr.VI,145	E., As., Afr., Am., P.,
S. vescum ; F. v. M. in Transact. Vict. Inst. 69 (1855)	—	—	T.	V.	N.S.W.	— —	--	M.fr.I,123	N.Z.
S. aviculare ; G. Forster, prodr. 18 (1786)	—	S.A.	T.	V.	N.S.W.	Q. —	B.fl.IV,447	M.fr.I,123; VI,144	N.Z.
S. simile ; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q —	B.fl.IV,448	M.fr.I,123; VI,144.	
S. fasciculatum ; F. v. M., fragm. 1, 123 (1859)	W.A.	S.A.	—	—	—	— —	—	M.fr.VI,144.	
S. Bauerianum ; Endlicher, prodr. fl. Norfolk, 54 (1833)	—	—	—	—	N.S.W.	— —	—	M.fr.IX,74.	
S. viride ; Solander in G. Forster, prodr. 89 (1786)	—	—	—	—	—	Q. —	B.fl.IV,449	M.fr.VI,145; IX,75	P.
S. tetrandrum ; R. Brown, prodr. 445 (1810)	—	—	—	—	—	Q. N.A.	B.fl.IV,449		
S. Shanesii ; F. v. M., fragm. VI, 144 (1868)	—	—	—	—	—	Q. —	B.fl.IV,448	M.fr.VI,254.	
S. verbascifolium ; Linné, spec. plant. 184 (1753)	—	—	—	—	N.S.W.	Q. —	B.fl.IV,449	M.fr.VI,145	As., Am., P.
S. discolor ; R. Brown, prodr. 445 (1810)	—	—	—	—	N.S.W.	Q. N.A.	B.fl.IV,450	M.fr.II,166; VI,147.	
S. stelligerum ; Smith, Exot. Bot. II, 57, t. 88 (1805)	—	—	—	—	N.S.W.	Q. —	B.fl.IV,450	M.fr.VI,147.	
S. parvifolium ; R. Brown, prodr. 446 (1810)	—	—	—	—	N.S.W.	Q. —	B.fl.IV,451	M.fr.II,164; VI,147.	
S. ferocissimum ; Lindley in Mitch. Three Exped. II, 58 (1838)	—	S.A.	—	—	N.S.W.	Q. —	B.fl.IV,451		
S. sporadotrichum ; F. v. M. in Melb. Chemist, Oct. (1882)	—	—	—	—	N.S.W.	Q. —	—		
S. defensum ; F. v. M., fragm. V, 193 (1866)	—	—	—	—	—	Q. —	B.fl.IV,451	M.fr.V,193.	
S. violaceum ; R. Brown, prodr. 445 (1810)	—	—	—	—	N.S.W.	Q. —	B.fl.IV,452		
S. tethrathecum ; F. v. M., fragm. II, 165 (1861)	—	—	—	—	N.S.W.	Q. —	B.fl.IV,453	M.fr.II,165.	
S. elachophyllum ; F. v. M., fragm. II, 164 (1861)	—	—	—	—	—	Q. —	B.fl.IV,453	M.fr.II,164.	
S. orbiculatum ; Dunal in Poir. Encycl. méth. XIII, 762 (1817)	W.A.	S.A.	—	—	—	— —	B.fl.IV,453	M.fr.VI,145.	

S. oligacanthum; F. v. M. in Transact. phil. Soc. Vict. I. 19 (1854)	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,454	M.fr.VI,145.		
S. esuriale; Lindley in Mitch. Three Exped. II, 43 (1838)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,454			
S. chenopodinum; F. v. M., fragm. II, 165 (1861)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,454	M.fr.VI,146.		
S. Sturtianum; F. v. M. in Transact. phil. Soc. Vict. I, 19 (1854)	—	S.A.	—	—	N.S.W.	—	N.A.	B.fl.IV,454	M.fr.VI,146.		
S. furfuraceum; R. Brown, prodr. 446 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,455			
S. dianthophorum; Dunal, hist. des Solan. 183 (1813)	—	—	—	—	—	Q.	—	B.fl.IV,455	M.fr.VI,145.		
S. Dallachyi; Bentham, Fl. Austr. IV, 456 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,456	M.fr.VI,145.		
S. densivestitum; F. v. M. in Benth. Fl. Austr. IV, 456 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,456			
S. nemophilum; F. v. M., fragm. II, 161 (1861)	—	—	—	—	—	Q.	—	B.fl.IV,456	M.fr.II,161.		
S. Oldfieldii; F. v. M., fragm. II, 161 (1861)	W.A.	—	—	—	—	—	—	B.fl.IV,457	M.fr.VI,145.		
S. semiarmatum; F. v. M., fragm. II, 163 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,457	M.fr.VI,146.		
S. xanthocarpum; Schrader in Wendl. sert. Hannov. 8, t. 2 (1795)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,458	...	...	As., P.
S. Hystrix; R. Brown, prodr. 446 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,458			
S. cataphractum; Cunningham in Benth. Fl. Austr. IV, 459 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,459			
S. pungetium; R. Brown, prodr. 446 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,459	M.fr.VI,146.		
S. eremophilum; F. v. M. in Schlecht. Linnaea XXV, 432 (1852)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.IV,459	M.fr.VI,146.		
S. campanulatum; R. Brown, prodr. 446 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460	M.fr.VI,146.		
S. adenophorum; F. v. M., fragm. II, 162 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460	M.fr.II,162.		
S. cinereum; R. Brown, prodr. 446 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,460			
S. lacunarium; F. v. M. in Transact. phil. Soc. Vict. I, 18 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,461	M.fr.VI,146.		
S. petrophilum; F. v. M. in Linnaea XXV, 433 (1852)	—	S.A.	—	—	N.S.W.	—	—	B.fl.IV,461	M.fr.VI,146.		
S. diversiflorum; F. v. M., fragm. VI, 146 (1868)	—	—	—	—	—	—	N.A.	B.fl.IV,461	M.fr.VI,146.		
S. carduiforme; F. v. M., fragm. II, 163 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.IV,462	M.fr.II,163.		
S. melanospermum; F. v. M., fragm. II, 163 (1861)	—	—	—	—	—	—	N.A.	B.fl.IV,462	M.fr.II,163.		
S. horridum; Dunal, Solan. synops. 28 (1816)	—	—	—	—	—	—	N.A.	B.fl.IV,463			
S. echinatum; R. Brown, prodr. 447 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,463	M.fr.VI,145.		
S. lasiophyllum; Dunal in Poir. Encycl. méth. XIII, 764 (1817)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,463	M.fr.VI,145.		
S. ellipticum; R. Brown, prodr. 446 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,464	M.fr.VI,145.		
S. quadriloculatum; F. v. M., fragm. II, 161 (1861)	—	—	—	—	—	Q.	N.A.	B.fl.IV,464	M.fr.II,161.		
S. phlomoides; Cunningham in Benth. Fl. Austr. IV, 464 (1869)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,464			
S. Cunninghamii; Bentham, Fl. Austr. IV, 465 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,465			
LYCIUM; Linné, gen. plant. 57 (1737), from Celsus.											
L. Australe; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,467	M.fr.I,83,243; II,179.		
DATURA; Linné, gen. plant. 48 (1737).											
D. Leichhardtii; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,468	M.fr.VI,144.		
NICOTIANA; Tournefort, inst. 117, t. 41 (1700), from C. Bauhin (1623).											
N. suaveolens; Lehmann, gener. Nicot. hist. 43 (1818)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,469	M.fr.VI,144.		
ANTHOTROCHE; Endlicher, nov. stirp. mus. Vindob. dec. 6 (1839).											
A. pannosa; Endlicher, nov. stirp. mus. Vind. dec. 7 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,467	M.fr.VI,142.		
A. Blackii; F. v. M., fragm. VIII, 232 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,232.		
A. Walcottii; F. v. M., fragm. I, 123 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,468	M.fr.I,123.		
ANTHOCERCIS; Labillardière, Nov. Holl. pl. spec. II, 19 (1806). (Cyphanthera, Eadesia.)											
A. viscosa; R. Brown, prodr. 448 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,475	M.fr.VI,143.		
A. litorea; Labillardière, Nov. Holl. pl. spec. II, 19, t. 158 (1806)	W.A.	—	—	—	—	—	—	B.fl.IV,476	M.fr.VI,143; XI,123.		
A. gracilis; Bentham in De Candolle, prodr. X, 192 (1846)	W.A.	—	—	—	—	—	—	B.fl.IV,476			
A. genistoides; Miers, Illustr. of plants II, App. 27, t. 83 (1857)	W.A.	—	—	—	—	—	—	B.fl.IV,476	M.fr.I,122; VI,143.		
A. anisantha; Endlicher in Ann. Wien. Mus. II, 201 (1838)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,477	M.fr.I,122; VI,143.		

A. intricata; F. v. M., fragm. I, 211 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,477	M.fr.I,211.			
A. arborea; F. v. M., fragm. I, 212 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,477	M.fr.I,212; VI,143.			
A. angustifolia; F. v. M. in Transact. phil. Soc. Vict. I, 21 (1854)	—	S.A.	—	—	—	—	—	B.fl.IV,478	M.fr.II,139.			
A. fasciculata; F. v. M., fragm. I, 122 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,478	M.fr.I,122; VI,143.			
A. microphylla; F. v. M., fragm. I, 179 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,478	M.fr.I,179; VI,143.			
A. myosotidea; F. v. M. in Transact. phil. Soc. Vict. I, 20 (1854)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,478	M.fr.I,179; VI,143.			
A. scabrella; Bentham in De Candolle, prodr. X, 192 (1846)	—	—	—	—	N.S.W.	—	—	B.fl.IV,479				
A. albicans; Cunningham in Field's N.S.W. 335, t. 2 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,479	M.fr.VI,143.			
A. Odgersii; F. v. M., fragm. X, 19 (1876)	—	—	—	—	—	—	N.A.	—	M.fr.X,19.			
A. Tasmanica; J. Hooker, Fl. Tasman. I, 289, t. 92 (1860)	—	—	T.	—	—	—	—	B.fl.IV,479	M.fr.VI,143.			
A. Eadesii; F. v. M., fragm. II, 139 (1861)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,480	M.fr.VI,142.			
A. racemosa; F. v. M., fragm. I, 211 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,480	M.fr.I,211.			
ISANDRA; F. v. M. in Wing's S. Sc. Rec. III, 2 (1883).												
I. Bancroftii; F. v. M. in Wing's S. Sc. Rec. III, 2 (1883)	W.A.	—	—	—	—	—	—	—				
DUBOISIA; R. Brown, prodr. fl. Nov. Holl. 448 (1810).												
D. myoporoides; R. Brown, prodr. 448 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,474	M.fr.VI,144	...	P.	
D. Hopwoodi; F. v. M., fragm. VIII, 232 (1874)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,480	M.fr.II,138; VI,143; X,20,121.			
D. Leichhardti; F. v. M. in Wing's S. Sc. Rec. II, 222 (1882)	—	—	—	—	N.S.W.	—	—	B.fl.IV,480	M.fr.VI,142.			

SCROPHULARINAE.

MIRBEL, Elém. de Botan. II, 879 (1815).—Genera 22, Species 80.

MIMULUS; Linné in Act. Upsal. 82 (1741). (Uvedalia.)												
M. Uvedaliae; Bentham in De Candolle, prodr. X, 369 (1846)	—	—	—	—	—	Q.	N.A.	B.fl.IV,482	M.fr.VI,103; IX,167.			
M. debilis; F. v. M. in Transact. phil. Inst. Vict. I, 62 (1857)	—	—	—	—	—	—	N.A.	—	M.fr.VI,103.			
M. gracilis; R. Brown, prodr. 439 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,482	M.fr.VI,103; IX,167...		As., Afr.	
M. repens; R. Brown, prodr. 439 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,482	M.fr.VI,103; IX,166...		N.Z.	
M. prostratus; Bentham in De Candolle, prodr. X, 373 (1846)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,483	M.fr.IX,167.			
MAZUS; Loureiro, Fl. Cochinch. II, 385 (1790).												
M. Pumilio; R. Brown, prodr. 439 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,484	M.fr.VI,104; IX,167...		N.Z.	
ADENOSMA; R. Brown, prodr. fl. Nov. Holl. 442 (1810). (Pterostigma.)												
A. coerulea; R. Brown, prodr. 443 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,484	...	...	...	As.
A. Muelleri; Bentham, Fl. Austr. IV, 485 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,485				
STEMODIA; Linné, syst. ed. dec. 1118 (1759). (Morgania, Limnophila.)												
S. lythrifolia; F. v. M. in Benth. Fl. Austr. IV, 486 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,486				
S. grossa; Bentham, Fl. Austr. IV, 486 (1869)	W.A.	—	—	—	—	—	N.A.	B.fl.IV,486				
S. viscosa; Roxburgh, pl. of Coromand. II, 33, t. 163 (1798)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,486	...	...	...	As.
S. linophylla; F. v. M., fragm. X, 88 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,88.			
S. debilis; Bentham, Fl. Austr. IV, 487 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,487				
S. pedicellaris; F. v. M., fragm. VIII, 231 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,231.			
S. Morgania; F. v. M., fragm. X, 89 (1876)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,488	M.fr.VI,104.			
S. sessiliflora; F. v. M., Cens. 97 (1882) from Blume (1826)	—	—	—	—	—	—	N.A.	—	...	...	...	As.
S. gratioloides; F. v. M., fragm. X, 89 (1876) from R. Brown (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,489	M.fr.X,89 ...		...	As., Afr.
S. punctata; F. v. M., Cens. 97 (1882) from Blume (1826)...	—	—	—	—	—	Q.	—	B.fl.IV,490	...	...	...	As.
S. hirsuta; Heyne in Wallich, numer. list. 3930 (1828)	—	—	—	—	—	—	N.A.	B.fl.IV,490	...	...	...	As.
S. tenuiflora; Bentham, Scrophul. Indic. 23 (1835)	—	—	—	—	—	—	N.A.	B.fl.IV,490	...	...	...	As., P.
S. Kingii; F. v. M. in Transact. R. S. of Vict. XXIII, 57 (1886)	W.A.	—	—	—	—	—	N.A.	—				

BRAMIA ; Lamarck, Encycl. méthod. I, 459 (1783). (Herpestis.)

B. floribunda ; F. v. M., fragm. IX, 167 (1875) from R. Brown (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,491	M.fr.VI,104	...	As., Afr.
B. Indica ; Lamarck, Encycl. méth. I, 459 (1783) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,491	M.fr.IX,167	...	As., Afr., Am.

GRATIOLA ; Ruppius, Fl. Jenens. 241 (1718), from Dodoens (1556).

G. pedunculata ; R. Brown, prodr. 435 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,492	M.fr.VI,108; IX,167.		
G. Peruviana ; Linné, spec. plant. 17 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,493	M.fr.VI,103; IX,167...		Am., N.Z.
G. nana ; Bentham in De Candolle, prodr. X, 404 (1846) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,493	M.fr.IX,167	...	N.Z.

DOPATRIUM ; Hamilton in Benth. Scroph. Ind. 4 (1835).

D. junceum ; Hamilton in Benth. Scroph. Ind. 4 (1835) ...	—	—	—	—	—	Q.	—	B.fl.IV,494	...	...	...	As.

ARTANEMA ; D. Don in Sweet, Fl. gard. t. 234 (1829).

A. fimbriatum ; D. Don in Sweet, Fl. Gard. t. 234 (1829) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,495	M.fr.VI,104; IX,167.		

LINDERNIA ; Allioni, stirp. agr. Nicaeens enum. 178, t. 5 (1757). (Pyxidaria, Vandellia, Bonnaya, Ilysanthes, Torenia partly, Tittmannia partly, Ilyogeton.)

L. crustacea ; F. v. M., Cens. 97 (1882) from Linné (1767) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,496	...	...	...	As., Afr., Am.
L. pubescens ; F. v. M., Cens. 97 (1882) from Bentham (1846) ...	—	—	—	—	—	—	N.A.	B.fl.IV,496	M.fr.VI,102.		
L. pyxidaria ; Linné, mant. alter. 252 (1771) ...	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,496	...	...	...	E., As., P.
L. scapigera ; R. Brown, prodr. 441 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,497	M.fr.VI,102.		
L. subulata ; R. Brown, prodr. 441 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,497	M.fr.VI,102.		
L. lobelioides ; F. v. M., Cens. 97 (1882) ...	—	—	—	—	—	—	N.A.	B.fl.IV,497	M.fr.VI,102.		
L. clausa ; F. v. M., fragm. VI, 102 (1868) ...	—	—	—	—	—	—	N.A.	B.fl.IV,498			
L. veronicifolia ; F. v. M., fragm. VI, 101 (1868) ...	—	—	—	—	—	Q.	—	B.fl.IV,498	M.fr.VI,101	...	As., P.
L. serrata ; F. v. M., Cens. 97 (1882) from Roxburgh (1820) ...	—	—	—	—	—	Q.	—	—	...	...	...	As.

HEMIARRHENA ; Bentham, Fl. Austr. IV, 518 (1869).

H. plantaginea ; Bentham in Hook. icon. pl. t. 1059 (1869) ...	—	—	—	—	—	—	N.A.	B.fl.IV,518	M.fr.VI,102.		

PEPLIDIUM ; Delile, Fl. Aegypt. 148 (1813).

P. humifusum ; Delile, Fl. Aegypt. 148 (1813) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,500	M.fr.VI,105	...	As., Afr.
P. Muelleri ; Bentham, Fl. Austr. IV, 500 (1869) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,500			

MICROCARPAEA ; R. Brown, prodr. 435 (1810).

M. muscosa ; R. Brown, prodr. 435 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,501	M.fr.VI,104	...	As.

GLOSSOSTIGMA ; Arnott in nov. act. acad. Caes. Leop. Carol. XVIII, 355 (1836). (Tricholoma.)

G. spathulatum ; Arnott in nov. act. Leop. XVIII, 355 (1836) ...	—	—	—	—	—	Q.	—	B.fl.IV,501	...	..	...	As., Afr.
G. Drummondii ; Bentham in De Candolle, prodr. X, 426 (1846) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,502	M.fr.VI,104.		
G. elatinoides ; Bentham in J. Hook. Fl. New Zeal. I, 189 (1853) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,502	...	...	...	N.Z.

LIMOSELLA ; Lindern, opusc. plant. Argentorat. 156, t. 5 (1728).

L. aquatica ; Linné, spec. plant. 631 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,502	M.fr.VI,105; IX,165...		E., As., Afr., Am., [N.Z.
L. Curdieana ; F. v. M., fragm. IX, 166 (1875) ...	—	S.A.	—	V.	N.S.W.	Q.	—	—	M.fr.IX,166; XI,135.		

SCOPARIA ; Linné, syst. nat. ed. séxt. 87 (1748).

S. dulcis ; Linné, spec. plant. 116 (1753) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,504	M.fr.VI,104	...	Am.

OURISIA ; Commerçon in A. L. de Jussieu, gen. pl. 100 (1789).

O. integrifolia ; R. Brown, prodr. 439 (1810) ...	—	—	T.	—	—	—	—	B.fl.IV,512	M.fr.VI,105; IX,168.		

VERONICA ; Tournefort, inst. 143, t. 60 (1700) from Fuchs (1543). (Pygmaea.)

V. densifolia ; F. v. M., fragm. II, 137, t. 63 (1861) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,505	M.fr.VI,103.		
V. formosa ; R. Brown, prodr. 434 (1810) ...	—	—	T.	—	—	—	—	B.fl.IV,506	M.fr.VI,102; IX,167.		
V. decorosa ; F. v. M. in Schlechtend. Linnaea XXV, 430 (1852) ...	—	S.A.	—	—	—	—	—	B.fl.IV,506			
V. perfoliata ; R. Brown, prodr. 434 (1810) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,507	M.fr.VI,102; IX,167		

V. Derwentia ; Littlejohn in Andr. Bot. Reposit. t. 531 (1808)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,507	M.fr.VI,102; IX,168.		
V. arenaria ; Cunningham in De Candolle, prodr. X, 463 (1846)	—	—	—	—	N.S.W.	—	—	B.fl.IV,507	M.fr.VI,102; IX,167.		
V. nivea ; Lindley, Bot. Regist. Misc. 42 (1842)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,508	M.fr.VI,102; IX,167.		
V. gracilis ; R. Brown, prodr. 435 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,508	M.fr.VI,103; IX,168.		
V. distans ; R. Brown, prodr. 435 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,509	M.fr.VI,102.		
V. calycina ; R. Brown, prodr. 435 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,509	M.fr.VI,103.		
V. plebeja ; R. Brown, prodr. 435 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,510	M.fr.VI,102.	...	N.Z.
V. notabilis ; F. v. M. in Benth. Fl. Austr. IV, 511 (1869)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,511	M.fr.IX,168.		
V. serpillifolia ; Linné, spec. plant. 12 (1753)...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,511	M.fr.VI,103	...	E., As., Afr., Am.
V. peregrina ; Linné, spec. plant. 14 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,511	M.fr.VI,103; IX,168.		Am.
RAMPHICARPA ; Bentham in Hook. Comp. to the Bot. Mag. I, 368 (1835).											
R. longiflora ; Bentham in Hook. Comp. Bot. Mag. I, 368 (1835)	—	—	—	—	—	Q.	N.A.	B.fl.IV,518	M.fr.VI,104; IX,168.		As.
CENTRANTHERA ; R. Brown, prodr. fl. Nov. Holl. 438 (1810).											
C. hispida ; R. Brown, prodr. 438 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,513	M.fr.VI,104; IX,168.		As.
SOPUBIA ; Hamilton in D. Don, prodrom. flor. Nepalens. 88 (1825).											
S. trifida ; Hamilton in D. Don, prodr. fl. Nep. 88 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,513			
BUECHNERA ; Linné, hort. Cliffort. 501 (1737). (Buchnera, Striga, Campuleia.)											
B. tetragona ; R. Brown, prodr. 437 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,514	M.fr.IX,168.		
B. urticifolia ; R. Brown, prodr. 437 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,514	M.fr.VI,104.		
B. linearis ; R. Brown, prodr. 437 (1810)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,515			
B. tenella ; R. Brown, prodr. 437 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,515			
B. gracilis ; R. Brown, prodr. 437 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,515	M.fr.VI,104.		
B. ramosissima ; R. Brown, prodr. 438 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,515			
B. coccinea ; Bentham, Scrophul. Ind. 40 (1835)	—	—	—	—	—	Q.	—	B.fl.IV,516			
B. parviflora ; R. Brown, prodr. 438 (1810)	W.A.	—	—	—	—	Q.	N.A.	B.fl.IV,517	M.fr.VI,104	...	As., Afr., P.
B. multiflora ; F. v. M., Cens. 98 (1882) from Bentham (1836)	—	—	—	—	—	—	N.A.	B.fl.IV,517	... ...	...	As.
B. curviflora ; R. Brown, prodr. 438 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,517	M.fr.VI,104.		
EUPHRASIA ; Tournefort, inst. 174, t. 78 (1700) from Matthaeus (1478).											
E. Brownii ; F. v. M., fragm. V, 88 (1865)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,521	M.fr.IX,168	...	P.
E. scabra ; R. Brown, prodr. 437 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,521	M.fr.V,89; IX,168.		
E. arguta ; R. Brown, prodr. 437 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,522			
E. cuspidata ; J. Hooker, Fl. Tasm. I, 298 (1860)	—	—	T.	—	—	—	—	B.fl.IV,522	M.fr.V,90.		
E. antarctica ; Bentham in De Candolle, prodr. X, 555 (1846)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,522	M.fr.V,90 ...	...	Am., N.Z.

OROBANCHEAE.

A. L. de Jussieu in Ann. Mus. V, 25 (1804).—Genus 1, Species 1.

OROBANCHE ; Tournefort, inst. 175, t. 81 (1700), from l'Ecluse (1583).

O. Australiana ; F. v. M., Cens. 98 (1882)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,533			

LENTIBULARINAE.

L. C. Richard in Bulliard, Fl. Paris. ed. sec. I, 26 (1796).—Genera 2, Species 25.

UTRICULARIA ; Linné, syst. nat. 8 (1735); Linné, gen. plant. 5 (1737).

U. stellaris ; Linné, fil. suppl. plant. 86 (1781)	—	—	—	—	—	Q.	—	B.fl.IV,525	M.fr.VI,161; IX,48...		As., Afr.
U. flexuosa ; Vahl, enum. plant. I, 198 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.IV,525	.. ...	...	As.
U. exoleta ; R. Brown, prodr. 430 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,526	M.fr.VI,161	...	As.

U. tubulata; F. v. M., fragm. IX, 48 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,48.						
U. albiflora; R. Brown, prodr. 431 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,526							
U. pygmaea; R. Brown, prodr. 432 (1810)	—	—	—	—	—	Q.	B.fl.IV,526								
U. fulva; F. v. M. in Transact. phil. Inst. Vict. III, 63 (1858)	—	—	—	—	—	—	N.A.	B.fl.IV,527	M.fr.VI,160.						
U. chrysantha; R. Brown, prodr. 432 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,527	M.fr.VI,161; IX,48.						
U. bifida; Linné, spec. plant. 18 (1753)	—	—	—	—	—	Q.	—	B.fl.IV,527	...	...	...	As.			
U. cyanea; R. Brown, prodr. 431 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,527	M.fr.VI,161.	..	...	As.			
U. lateriflora; R. Brown, prodr. 431 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,528	M.fr.VI,162; IX,48.						
U. simplex; R. Brown, prodr. 431 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,528	M.fr.VI,162.						
U. monantha; J. Hooker, Fl. Tasm. I, 299 (1860)	—	—	T.	—	—	—	—	B.fl.IV,528							
U. dichotoma; Labillardière, Nov. Holl. pl. spec. I, 11, t. 8 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,529	M.fr.VI,160.						
U. volubilis; R. Brown, prodr. 430 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,529	M.fr.VI,161.						
U. Hookeri; Lehmann, nov. stirp. pugill. VIII, 47 (1844)	W.A.	—	—	—	—	—	—	B.fl.IV,530	M.fr.VI,162.						
U. violacea; R. Brown, prodr. 431 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,530	M.fr.VI,160.						
U. Menziesii; R. Brown, prodr. 431 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,530	M.fr.VI,160.						
U. biloba; R. Brown, prodr. 432 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,531	...	...	...	As.			
U. leptoplectra; F. v. M. in Austr. Chem. VIII, 50 (1885)	—	—	—	—	—	—	N.A.	—							
U. limosa; R. Brown, prodr. 432 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,531							
U. Baueri; R. Brown, prodr. 431 (1810)	—	—	—	—	—	Q.	—	B.fl.IV,531	...	...	...	As.			
U. lasiocaulis; F. v. M. in Austr. Chem. VIII, 55 (1885)	—	—	—	—	—	—	N.A.	—							
POLYPOMPHOLYX; Lehmann in der Bot. Zeitung, Halle, 109 (1844). (Tetralobus.)															
P. multifida; F. v. M., fragm. VI, 162 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,532	M.fr.VI,162.						
P. tenella; Lehmann, nov. stirp. pugill. VIII, 50 (1844)	W.A.	S.A.	T.	V.	—	—	—	B.fl.IV,532	M.fr.VI,162.						

PODOSTEMONEAE.

L. Cl. Richard in Humb. Bonpl. & Kunth, Nov. gen. I, 246 (1815).—Genus 1, Species 1.

Genus and species not yet determined	—	—	—	—	—	Q.	—	—	

GESNERIACEAE.

Humboldt, Bonpland & Kunth, nov. gen. et spec. pl. II, 392 (1817).—Genera 4, Species 4.

FIELDIA; Cunningham in Field's New South Wales 363, t. 2 (1825).

F. Australis; Cunningham in Field's N.S.W. 364, t. 2 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,535	M.fr.IV,147; IX,191.

NEGRIA; F. v. M., fragm. VII, 151 (1871). (Rhabdothamnus partly.)

N. rhabdothamnoides; F. v. M., fragm. VII, 152 (1871)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,152; VIII,281; XI,133.

BAEA; Commerçon in Lamarck, Encycl. méth. I, 401 (1783). (Streptocarpus, Boea, Bea.)

B. hygroscopica; F. v. M., fragm. IV, 146 (1864)	—	—	—	—	—	Q.	—	B.fl.IV,535	M.fr.IV,146.

ROETTLERA; Vahl. enum. plant. I, 87 (1805). (Didymocarpus.)

R. Kinnearii; F. v. M. in Vict. Naturalist III, 159 (1887)	—	—	—	—	—	Q.	—	—	

BIGNONIACEAE.

Ventenat; Tabl. du. règn. végét. II, 402 (1799).—Genera 4, Species 7.

TECOMA; A. L. de Jussieu, gen. plant. 139 (1789).

T. Australis; R. Brown, prodr. 471 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,537	
T. Hillii; F. v. M., fragm. X, 101 (1877)	—	—	—	—	—	Q.	—	—	M.fr.X,101; XI,136.
T. jasminoides; Lindley, Bot. Regist. t. 2002 (1838)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,537	

DOLICHANDRONE ; Fenzl in Ann. nat. hist. third ser. X, 31 (1862). (Spathodea partly, Dolichandra partly.)

D. heterophylla ; F. v. M., fragm. IV, 149 (1864) ... ... — — — — — Q. N.A. B.fl.IV,539 M.fr.IV,149.
D. filiformis ; Seemann, Jour. of Bot. I, 226 (1863) ... ... — — — — — — N.A. B.fl.IV,539

HAUSSMANNIA ; F. v. M., fragm. IV, 148 (1864).

H. jucunda ; F. v. M., fragm. IV, 148 (1864)... ... ... — — — — — Q. — B.fl.IV,540 M.fr.IV,148.

DEPLANCHEA ; Vieillard in Bullet. de la Soc. Linn. de Norm. VII, 11 (1862). (Diplanthera, Bulweria.)

D. tetraphylla ; F. v. M., from R. Brown (1810) ... ... — — — — — Q. — B.fl.IV,556 M.fr.IV,147;VI,72.

PEDALINAE.

R. Brown, prodr. Fl. Nov. Holl. 519 (1810).—Genus 1, Species 3.

JOSEPHINA ; Ventenat, Jardin de la Malmaison 103, t. 67 (1804).

J. grandiflora ; R. Brown, prodr. 520 (1810) ... ... — — — — — Q. — B.fl.IV,540 ... ... P.
J. Imperatricis ; Ventenat, Jard. de la Malm. t. 67 (1804)... .. .. — — — — — — N.A. B.fl.IV,557 M.fr.VI,163... ... As.
J. Eugeniae ; F. v. M. in Hook. Kew Misc. IX, 370, t. 11 (1857)... ... ... W.A. S.A. — — — Q. N.A. B.fl.IV,557 M.fr.VI,163.

ACANTHACEAE.

A. L. de Jussieu, gen. plant. 102 (1789), from B. de Jussieu (1759).—Genera 13, Species 30.

THUNBERGIA ; Retzius, Physiogr. Saellsk. Handl. I, 163 (1776).

T. Arnhemica ; F. v. M., fragm. IX, 73 (1875) ... ... — — — — — — N.A. — M.fr.IX,73 ... ... As.
T. Powelli ; F. v. M. in Wing's South. Sc. Rec. II, 34 (1882) ... ... — — — — — — N.A. —
T. fragrans ; J. Koenig in Roxb. pl. Coromand. I & IV (1795) ... ... — — — — — — N.A. —

NELSONIA ; R. Brown, prodr. fl. Nov. Holl. 485 (1810).

N. campestris ; R. Brown, prodr. 481 (1810) ... . — — — — — Q. N.A. B.fl.IV,543 M.fr.VI,88; XI,18 ... As., Afr., Am.

EBERMAYERA ; Nees in Wallich, pl. Asiat. rar. III, 75 (1832).

E. glauca ; Nees in De Candolle, prodr. XI, 73 (1847) ... ... — — — — — Q. N.A. B.fl.IV,544 M.fr.VI,88; XI,18 ... As.

HYGROPHILA ; R. Brown, prodr. fl. Nov. Holl. 479 (1810).

H. angustifolia ; R. Brown, prodr. 479 (1810) ... ... — — — — — Q. N.A. B.fl.IV,544 M.fr.VI,88 ... ... As.

STROBILANTHES ; Blume, Bijdr. tot de fl. van Nederl. Ind. 796 (1826).

S. Tatei ; F. v. M. in Transact. Roy. Soc. S. Austr. V, Aug. (1882) ... ... — — — — — — N.A. —

RUELLIA ; Plumier, nov. gen. 12, t. 2 (1703). (Dipteracanthus.)

R. bracteata ; R. Brown, prodr. 479 (1810) .. .. — — — — — Q. N.A. B.fl.IV,546 M.fr.VI,91 ... ... As.
R. primulacea ; F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ... ... — S.A. — — — Q. — B.fl.IV,546 M.fr.XI,18.
R. corynotheca ; F. v. M. in Benth. Fl. Austr. IV, 546 (1869) ... ... — — — — N.S.W. Q. — B.fl.IV,546 M.fr.XI,18.
R. Australis ; R. Brown, prodr. 479 (1810) ... ... — S.A. — — N.S.W. Q. — B.fl.IV,547 M.fr.VI,91.
R. spiciflora ; F. v. M. in Benth. Fl. Austr. IV, 547 (1869) ... ... — — — — — Q. — B.fl.IV,547
R. acaulis ; R. Brown, prodr. 479 (1810) ... ... — — — — — Q. — B.fl.IV,547

ACANTHUS ; Tournefort, inst. 176, t. 81 (1700) from Theophrastos. (Dilivaria.)

A. ilicifolius ; Linné, spec. plant. 939 (1753) ... ... — — — — — Q. N.A. B.fl.IV,548 M.fr.X,18 ... ... As., P.

JUSTICIA ; Houston in Linné, gen. plant. 4 (1737). (Rostellaria, Rostellularia.)

J. procumbens ; Linné, spec. plant. 15 (1753) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.IV,549 M.fr.VI,91; XI,18; XII,24 As., Afr., P.
J. Bonneyana ; F. v. M. in Wing's S. Sc. Record II, 74 (1882) ... ... — S.A. — — N.S.W. — — — M.fr.X,73.
J. hygrophiloides ; F. v. M., fragm. VI, 89 (1868) ... ... — — — — — Q. — B.fl.IV,550 M.fr.VI,89.
J. Kempeana ; F. v. M., fragm. XI, 101 (1881) ... ... W.A. S.A. — — — — N.A. — M.fr.XI,101.

J. cavernarum ; F. v. M., fragm. VI, 91 (1868)						—	—	—	—	—	Q.	—	B.fl.IV,550	M.fr.VI,91.		
J. eranthemoides ; F. v. M., fragm. VI, 90 (1868)						—	—	—	—	N.S.W.	Q.	—	B.fl.IV,551	M.fr.VI,90.		
GRAPTOPHYLLUM ; Nees in Wallich, pl. Asiat. rar. III, 102 (1832). (Earlia, Thyrsacanthus.)																
G. Earlii ; F. v. M., fragm. VI, 87 (1868)						—	—	—	—	—	Q.	—	B.fl.IV,551	M.fr.III,160.		
G. ilicifolium ; F. v. M. in Benth. Fl. Austr. IV, 552 (1869)						—	—	—	—	—	Q.	—	B.fl.IV,552			
G. spinigerum ; F. v. M., fragm. XI, 17 (1878)						—	—	—	—	—	Q.	—	—	M.fr.XI,17.		
DICLIPTERA ; A. L. de Jussieu in Ann. du Mus. IX, 267 (1807). (Brochosiphon.)																
D. glabra ; Decaisne in Nouv. Annal. du Mus. III, 388 (1834)						—	—	—	—	—	—	N.A.	B.fl.IV,552	M.fr.VI,88; XI,18	...	As.
D. spicata ; Decaisne in Nouv. Annal. du Mus. III, 389 (1834)						—	—	—	—	—	Q.	N.A.	B.fl.IV,553	M.fr.VI,89	...	As., P.
D. Burmanni ; Nees in Wallich, pl. Asiat. rar. III, 112 (1832)						—	—	—	—	—	Q.	—	—	M.fr.VII,62	...	As.
HYPOESTES ; Solander in R. Brown, prodr. 474 (1810).																
H. floribunda ; R. Brown, prodr. 474 (1810)						—	—	—	—	N.S.W.	Q.	N.A.	B.fl.IV,553	M.fr.XI,18.		
ERANTHEMUM ; Linné, Fl. Zeyl. 6 (1747).																
E. variabile ; R. Brown, prodr. 477 (1810)						—	—	—	—	N.S.W.	Q.	—	B.fl.IV,555	M.fr.VI,91; XI,18	...	P.
E. tenellum ; Bentham, Fl. Austr. IV, 555 (1869)						—	—	—	—	—	Q.	—	B.fl.IV,555			
DICLADANTHERA ; F. v. M., fragm. XII, 23 (1882).																
D. Forrestii ; F. v. M., fragm. XII, 23 (1882)					W.A.	—	—	—	—	—	—	—	—	M.fr.XII,23		

LABIATAE.

ADANSON, Famill. des plant. II, 180 (1763), from B. de Jussieu (1759).—Genera 20, Species 125.

OCIMUM ; Tournefort, inst. 203, t. 96 (1700), from Theophrastos and Dioscorides. (Ocymum.)																
O. sanctum ; Linné, mantiss. plant. 85 (1767)						—	—	—	—	—	Q.	N.A.	B.fl.V,74	M.fr.IV,46	...	As., Afr., P.
MOSCHOSMA ; Reichenbach, conspect. regn. vegetab. 115 (1828). (Basilicum 1802.)																
M. polystachya ; Bentham in Wall. pl. As. rar. II, 13 (1831)						—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,75	M.fr.VI,108	...	As., Afr., P.
ORTHOSIPHON ; Bentham in Wall. pl. As. rar. II, 15 (1831).																
O. stamineus ; Bentham in Wall. pl. As. rar. II, 15 (1831)						—	—	—	—	—	Q.	N.A.	B.fl.V,76	M.fr.IV,47; VI,108; IX,161.		As., P.
COLEUS ; Loureiro, Fl. Cochinch. II, 372 (1790).																
C. scutellarioides ; Bentham in Wall. pl. Asiat. rar. II, 16 (1831)						—	—	—	—	—	Q.	N.A.	B.fl.V,79	M.fr.IV,47; VI,108	...	As., P.
PLECTRANTHUS ; L'Héritier, stirp. nov. I, 85 (1785).																
P. foetidus ; Bentham, labiat. 35 (1832)						—	—	—	—	—	Q.	—	—	M.fr.IX,161; XI,135.		
P. congestus ; R. Brown, prodr. 506 (1810)						—	—	—	—	—	Q.	—	B.fl.V,78	M.fr.VI,108.		
P. parviflorus ; Henckel, adumbrat. pl. hort. Halens. (1806)					—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,78	M.fr.IV,47; VI,108; IX,161.		P.	
P. longicornis ; F. v. M., fragm. V, 51 (1865)						—	—	—	—	—	Q.	—	B.fl.V,77	M.fr.V,51.		
POGOSTEMON ; Desfontaines in Mém. du Mus. II, 154, t. 6 (1815). (Dysophylla.)																
P. verticillatus ; Hasskarl, catal. hort. Bogor. 131 (1844)						—	—	—	—	—	Q.	N.A.	B.fl.V,81	M.fr.V,200; IX,161	...	As.
MENTHA ; Tournefort, inst. 188, t. 89 (1700), from Hippocrates and Theophrastos.																
M. laxiflora ; Bentham in De Candolle, prodr. XII, 174 (1848)						—	—	—	V.	N.S.W.	—	—	B.fl.V,82	M.fr.VI,109; IX,162.		
M. grandiflora ; Bentham in Mitch. Trop. Austr. 362 (1848)						—	—	—	—	N.S.W.	Q.	—	B.fl.V,83			
M. Australis ; R. Brown, prodr. 505 (1810)						—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.V,83	M.fr.VI,109; IX,162.		
M. gracilis ; R. Brown, prodr. 505 (1810)						—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,83	M.fr.VI,109.		
M. saturejoides ; R. Brown, prodr. 505 (1810)					W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,84	M.fr.VI,109.		
LYCOPUS ; Tournefort, inst. 180, t. 83 (1700), from Plinius.																
L. Australis ; R. Brown, prodr. 500 (1810)						—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,85	M.fr.VI,109.		

SALVIA ; Linné, gen. plant. 6 (1737), from Plinius.

S. plebeja ; R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,85	M.fr.VI,109; IX,161	...	As., P.

BRUNELLA ; L'Ecluse, rar. stirp. hist. II, 42, 43 (1576), from Brunfels (1536). (Prunella.)

B. vulgaris ; A. De Candolle, prodr. XII, 410 (1848)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,87	M.fr.VI,109	...	E., As., Afr. Am.

SCUTTELLARIA ; Hermann, hort. Lugd. bot. cat. (1687), from Cortusi (1591).

S. mollis ; R. Brown, prodr. 507 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.V,88	M.fr.VI,109.		
S. humilis ; R. Brown, prodr. 507 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,88	M.fr.VI,109.		

ANISOMELES ; R. Brown, prodr. 503 (1810).

A. salvifolia ; R. Brown, prodr. 503 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,89	M.fr.VI,109; IX,162	...	P.

LEUCAS ; J. Burmann, thesaur. Zeyl. 140 (1737).

L. flaccida ; R. Brown, prodr. 505 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,90	M.fr.VI,109	...	As.

PROSTANTHERA ; Labillardière, Nov. Holl. pl. spec. 18, t. 157 (1806). (Chilodia, Cryphia, Klanderia.)

P. lasiantha ; Labillardière, Nov. Holl. pl. spec. II, 18, t. 157 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,93	M.fr.VI,105.		
P. prunelloides ; R. Brown, prodr. 508 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,94	M.fr.VI,106.		
P. coerulea ; R. Brown, prodr. 508 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,94	M.fr.VI,106.		
P. ovalifolia ; R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,95	M.fr.VI,106.		
P. melissifolia ; F. v. M., fragm. I, 19 (1858)	—	—	—	V.	—	—	—	B.fl.V,95	M.fr.I,19,242.		
P. incisa ; R. Brown, prodr. 509 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,95	M.fr.VI,107; IX,162.		
P. rotundifolia ; R. Brown, prodr. 509 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,96	M.fr.VI,107; IX,162.		
P. Schultzii ; F. v. M., cens. fourth suppl. 4 (1889)	—	S.A.	—	—	—	—	N.A.	—			
P. violacea ; R. Brown, prodr. 509 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.V,96	M.fr.VI,107; IX,162.		
P. incana ; Cunningham in Benth. lab. 455 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.V,97	M.fr.IX,162.		
P. hirtula ; F. v. M., first gen. Report 16 (1853)	—	—	—	V.	N.S.W.	—	—	B.fl.V,97			
P. denticulata ; R. Brown, prodr. 509 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.V,97	M.fr.VI,105.		
P. rugosa ; Cunningham in Benth. lab. 456 (1836)	—	—	—	—	N.S.W.	—	—	B.fl.V,98			
P. marifolia ; R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,98	M.fr.VI,105.		
P. rhombea ; R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,99			
P. spinosa ; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)	—	S.A.	—	V.	—	—	—	B.fl.V,99	M.fr.VI,108; IX,162.		
P. cuneata ; Bentham in De Candolle, prodr. XII, 560 (1848)	—	—	T.	V.	N.S.W.	—	—	B.fl.V,99	M.fr.VI,108.		
P. linearis ; R. Brown, prodr. 509 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.V,100	M.fr.VI,106.		
P. phylicifolia ; F. v. M., fragm. I, 19 (1858)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,100	M.fr.I,19; IX,162.		
P. decussata ; F. v. M., fragm. I, 126 (1858)	—	—	—	V.	—	—	—	B.fl.V,100	M.fr.I,126.		
P. empetrifolia ; Sieber in Spreng. syst. cur. post. 226 (1827)	—	—	—	—	N.S.W.	—	—	B.fl.V,101			
P. lithospermoides ; F. v. M., fragm. VI, 107 (1868)	—	—	—	—	—	Q.	—	B.fl.V,101	M.fr.VI,107.		
P. Behriana ; Schlechtendal, Linnaea XX, 610 (1847)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,102	M.fr.VI,106.		
P. Wilkieana ; F. v. M., fragm. VIII, 230 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,230.		
P. Baxteri ; Cunningham in Benth. lab. 452 (1836)	W.A.	S.A.	—	—	—	—	—	B.fl.V,102	M.fr.VI,106.		
P. Campbelli ; F. v. M. in Wing's S. Sc. Record II, 25 (1882)	W.A.	—	—	—	—	—	—	—			
P. canaliculata ; F. v. M., fragm. VI, 105 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,102	M.fr.VI,105.		
P. nivea ; Cunningham in Benth. lab. 452 (1836)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,103	M.fr.VI,106.		
P. striatiflora ; F. v. M. in Schlechtend. Linnaea XXV, 425 (1852)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,103	M.fr.VI,105; IX,162.		
P. saxicola ; R. Brown, prodr. 509 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,104	M.fr.VI,107; IX,162		
P. debilis ; F. v. M., fragm. VIII, 147 (1874)	—	—	—	V.	—	—	—	—	M.fr.VIII,147.		
P. odoratissima ; Bentham in Mitch. Trop. Austr. 291 (1848)	—	—	—	—	—	Q.	—	B.fl.V,104			
P. Eckersleyana ; F. v. M., fragm. X, 17 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,17.		
P. euphrasioides ; Bentham in Mitch. Trop. Austr. 360 (1848)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,104	M.fr.VI,108.		
P. staurophylla ; F. v. M., fragm. IX, 73 (1875)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,73.		

P. cryptandroides; Cunningham in Benth. lab. 453 (1836)...				—	—	—	—	N.S.W.	—	—	B.fl.V,105	
P. eurybioides; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)				—	S.A.	—	—	—	—	—	B.fl.V,105	M.fr.VI,105.
P. ringens; Bentham in Mitch. Trop. Austr. 363 (1848)				—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,106	
P. Leichhardtii; Bentham, Fl. Austr. V, 106 (1870)				—	—	—	—	—	Q.	—	B.fl.V,106	M.fr.VII,108.
P. Walteri; F. v. M., fragm. VII, 108 (1870)				—	—	—	V.	N.S.W.	—	—	—	M.fr.VII,108.
P. coccinea; F. v. M. in Transact. phil. Soc. Vict. I, 48 (1854)			W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,106	M.fr.VI,108; IX,162.
P. chlorantha; F. v. M. in Benth. Fl. Austr. V, 108 (1870)			—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,108	M.fr.IX,162; X,18.
P. calycina; F. v. M. in Benth. Fl. Austr. V, 107 (1870)			—	S.A.	—	—	—	—	—	B.fl.V,107	
P. Grylloana; F. v. M., fragm. X, 17 (1876)			W.A.	—	—	—	—	—	—	—	M.fr.X,17.
WRIXONIA; F. v. M., fragm. X, 18 (1876).												
W. prostantheroides; F. v. M., fragm. X, 18 (1876)			W.A.	—	—	—	—	—	—	—	M.fr.X,18.
HEMIGENIA; R. Brown, prodr. 502 (1810). (Hemiandra, Colobandra, Atelandra.)												
H. macrantha; F. v. M., fragm. I, 210 (1859)			W.A.	—	—	—	—	—	—	B.fl.V,112	M.fr.I,210; XI,19.
H. rigida; Bentham in De Candolle, prodr. XII, 563 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,112	
H. ramosissima; Bentham in De Candolle, prodr. XII, 565 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,113	
H. microphylla; Bentham in De Candolle, prodr. XII, 565 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,113	M.fr.VI,112; IX,162.
H. incana; Bentham in De Candolle, prodr. XII, 566 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,113	
H. canescens; Bentham in De Candolle, prodr. XII, 566 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,114	
H. podalyrina; F. v. M., fragm. VI, 112 (1868)			W.A.	—	—	—	—	—	—	B.fl.V,114	M.fr.VI,112.
H. platyphylla; Bentham in De Candolle, prodr. XII, 566 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,115	
H. obovata; F. v. M., fragm. XI, 19 (1878)			W.A.	—	—	—	—	—	—	—	B.fl.XI,19.
H. glabrescens; Bentham in De Candolle, prodr. XII, 566 (1848)...			W.A.	—	—	—	—	—	—	B.fl.V,115	
H. obtusa; Bentham in De Candolle, prodr. XII, 567 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,115	
H. sericea; Bentham in Hueg. enum. 80 (1837)			W.A.	—	—	—	—	—	—	B.fl.V,116	M.fr.VI,112.
H. barbata; Bartling in Lehm. pl. Preiss. I, 360 (1845)			W.A.	—	—	—	—	—	—	B.fl.V,116	
H. curvifolia; F. v. M., fragm. I, 210 (1859)			W.A.	—	—	—	—	—	—	B.fl.V,117	M.fr.I,210.
H. scabra; Bentham, Fl. Austr. V, 117 (1870)			W.A.	—	—	—	—	—	—	B.fl.V,117	
H. humilis; Bentham in De Candolle, prodr. XII, 567 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,117	
H. westringioides; Bentham in De Candolle, prodr. XII, 568 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,117	
H. teretiuscula; F. v. M., fragm. VI, 111 (1868)			W.A.	—	—	—	—	—	—	B.fl.V,118	M.fr.VI,111.
H. brachyphylla; F. v. M., fragm. X, 19 (1876)			W.A.	—	—	—	—	—	—	—	M.fr.X,19.
H. purpurea; R. Brown, prodr. 502 (1810)			—	—	—	—	N.S.W.	—	—	B.fl.V,118	M.fr.VI,112.
H. cuneifolia; Bentham, Fl. Austr. V, 118 (1870)			—	—	—	—	N.S.W.	—	—	B.fl.V,118	
H. Drummondii; Bentham, Fl. Austr. V, 119 (1870)			W.A.	—	—	—	—	—	—	B.fl.V,119	
H. pimelifolia; F. v. M., fragm. VI, 112 (1868)			W.A.	—	—	—	—	—	—	B.fl.V,119	M.fr.VI,112.
H. diplanthera; F. v. M., fragm. VI, 111 (1868)			W.A.	—	—	—	—	—	—	B.fl.V,119	M.fr.VI,111.
H. pungens; F. v. M., fragm. XI, 20 (1878)			W.A.	—	—	—	—	—	—	B.fl.V,109	M.fr.XI,20.
H. leiantha; F. v. M., fragm. XI, 20 (1878)			W.A.	—	—	—	—	—	—	B.fl.V,110	M.fr.XI,20.
H. loganiacea; F. v. M., fragm. XI, 19 (1878)			W.A.	—	—	—	—	—	—	B.fl.V,112	M.fr.VI,113.
MICROCORYS; R. Brown, prodr. 502 (1810). (Anisandra.)												
M. longifolia; Bentham in De Candolle, prodr. XII, 568 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,121	
M. longiflora; F. v. M., fragm. VI, 113 (1868)			W.A.	—	—	—	—	—	—	B.fl.V,121	M.fr.VI,113.
M. tenuifolia; Bentham, Fl. Austr. V, 122 (1870)			W.A.	—	—	—	—	—	—	B.fl.V,122	
M. capitata; Bentham in De Candolle, prodr. XII, 568 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,122	
M. pimeloides; F. v. M., fragm. I, 156 (1859)			W.A.	—	—	—	—	—	—	B.fl.V,123	M.fr.I,156.
M. subcanescens; Bentham, Fl. Austr. V, 123 (1870)			W.A.	—	—	—	—	—	—	B.fl.V,123	
M. ericifolia; Bentham in De Candolle, prodr. XII, 569 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,123	
M. glabra; Bentham in De Candolle, prodr. XII, 569 (1848)			W.A.	—	—	—	—	—	—	B.fl.V,124	

M. exserta; Bentham, Fl. Austr. V, 124 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,124				
M. virgata; R. Brown, prodr. 502 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,125	M.fr.VI,112; VIII,231.			
M. Macrediana; F. v. M., fragm. VIII, 231 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,231.			
M. barbata; R. Brown, prodr. 502 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,125				
M. lenticularis; F. v. M., fragm. VI, 113 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,126	M.fr.VI,113.			
M. obovata; Bentham in De Candolle, prodr. XII, 569 (1848)	W.A.	—	—	—	—	—	—	B.fl.V,126	M.fr.XI,19.			
M. purpurea; R. Brown, prodr. 502 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,126	M.fr.VI,112.			
WESTRINGIA; Smith in Kongl. Svenska Vetenskaps Akad. Handlingar 171 (1797).												
W. cephalantha; F. v. M., fragm. VI, 110 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,127	M.fr.VI,110.			
W. rosmariniformis; Smith in Vet. Acad. Handl. 171 (1797)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.V,128	M.fr.VI,110; IX,162.			
W. Dampieri; R. Brown, prodr. 501 (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.V,129	M.fr.VI,110; IX,163.			
W. rigida; R. Brown, prodr. 501 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,129	M.fr.IX,163.			
W. angustifolia; R. Brown, prodr. 501 (1810)	—	—	T.	—	—	—	—	—.	M.fr.VI,110; IX,163.			
W. senifolia; F. v. M. in Transact. phil. Soc. Vict. I, 49 (1854)	—	—	—	V.	—	—	—	B.fl.V,130	M.fr.VI,110; IX,163.			
W. longifolia; R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,131	M.fr.VI,110; IX,163.			
W. glabra; R. Brown, prodr. 501 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.V,131	M.fr.VI,110; IX,163.			
W. rubiifolia; R. Brown, prodr. 501 (1810)	—	—	T.	—	—	—	—	B.fl.V,131	M.fr.VI,110.			
AJUGA; Linné, gen. plant. 167 (1737), from Scribonius (1529).												
A. Australis; R. Brown, prodr. 503 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,136	M.fr.VI,109.			
TEUCRIUM; Tournefort, inst. 207, t. 98 (1700), from Dioscorides.												
T. ajugaceum; F. v. M. & Bailey in syn. Queensland Flora, sec. suppl. 48 (1888)	—	—	—	—	—	Q.	—	—				
T. argutum; R. Brown, prodr. 504 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,135	M.fr.VI,110; IX,162.			
T. sessiliflorum; Bentham in De Candolle, prodr. XI, 580 (1848)...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,134	M.fr.VI,110.			
T. corymbosum; R. Brown, prodr. 504 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,133	M.fr.VI,110.			
T. fililoburn; F. v. M. in Benth. Fl. Austr. V, 134 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,134	M.fr.IX,162.			
T. integrifolium; F. v. M. in Benth. Fl. Austr. V, 133 (1870)	—	S.A.	—	—	—	Q.	N.A.	B.fl.V,133	M.fr.IX,162.			
T. racemosum; R. Brown, prodr. 504 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,133	M.fr.VI,109; IX,162.			

VERBENACEAE.

ADANSON, Familles des plantes II, 195 (1763), from B. de Jussieu (1759).—Genera 22, Species 82.

SPARTOTHAMNUS; Cunningham in Loudon's hort. Brit. 600 (1830).												
S. teucriiflorus; F. v. M. in Wing's S. Sc. Rec. II, 55 (1882)	W.A.	S.A.	—	—	—	—	N.A.	—				
S. junceus; Cunningham in Loud. hort. Brit. 600 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,55	M.fr.VI,153; IX,5	...	N.Z.	
S. puberulus; F. v. M., in Wing's S. Sc. Rec. II, 55 (1882)	—	S.A.	—	—	N.S.W.	Q.	—	—				
ONCINOCALYX; F. v. M. in Wing's S. Sc. Rec. III, 70 (1883).												
O. Betchei; F. v. M. in Wing's S. Sc. Rec. III, 70 (1883)	—	—	—	—	N.S.W.	—	—	—				
LIPPIA; Houston in Linné, syst. nat. 8 (1735); Linné, gen. plant. 347 (1737). (Zapania.)												
L. nodiflora; Cl. Richard in Mich. fl. bor. Amer. II, 15 (1803)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,35	M.fr.VI,151; IX,4	...	As., Afr., Am., P.	
L. geminata; Humboldt, Bonpland et Kunth, nov. gen. et spec. pl. II, 215 (1817)	—	—	—	—	—	Q.	—	B.fl.V,35	M.fr.VI,151	...	Am.	
VERBENA; Tournefort, inst. 200, t. 94 (1700), from l'Ecluse (1576).												
V. officinalis; Linné, spec. plant. 11 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.V,36	M.fr.VI,153; IX,5	...	E., As., Afr.	
V. macrostachya; F. v. M., fragm. I, 60 (1858)	—	S.A.	—	—	—	Q.	N.A.	B.fl.V,36	M.fr.I,60; IX,5.			
LACHNOSTACHYS; Hooker, icon. plant. t. 414 (1842). (Walcottia, Pycnolachne.)												
L. albicans; Hooker, icon. plant. 414 (1842)	W.A.	—	—	—	—	—	—	B.fl.V,38	M.fr.V,159.			
L. Cliftoni; F. v. M., fragm. IX, 3 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,3.			

L. verbascifolia; F. v. M., fragm. VI, 158 (1868)	W.A.	—	—	—	—	—	—					B.fl.V,38	M.fr.VI,158; IX,4.
L. ferruginea; Hooker, icon. plant. 415 (1842)	W.A.	—	—	—	—	—	—					B.fl.V,39	M.fr.VI,159.
L. Walcottii; F. v. M., fragm. II, 140 (1861)	W.A.	—	—	—	—	—	—					B.fl.V,39	M.fr.I,241; VI,159.
NEWCASTLIA; F. v. M. in Hook. Kew Misc. IX, 22 (1857).													
N. cladotricha; F. v. M. in Hook. Kew Misc. IX, 22 (1857)	W.A.	S.A.	—	—	—	—	N.A.					B.fl.V,40	M.fr.I,184; III,21; VI,155; VIII,50.
N. bracteosa; F. v. M., fragm. VIII, 49 (1873)	—	S.A.	—	—	—	—	—					—	M.fr.VIII,230; IX,4.
N. chrysotricha; F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—					—	M.fr.X,15.
N. hexarrhena; F. v. M., fragm. X, 16 (1876)	W.A.	—	—	—	—	—	—					—	M.fr.X,16.
N. spodiotricha; F. v. M., fragm. III, 21, t. 21 (1862)	W.A.	S.A.	—	—	—	—	N.A.					B.fl.V,40	M.fr.VIII,230; IX,4.
N. Dixoni; F. v. M. & Tate in Proceed. R. S. of S.A. X, 81 (1887)	—	S.A.	—	V.	—	—	—					—	
N. cephalantha; F. v. M., fragm. IX, 4 (1875)	—	S.A.	—	—	—	—	—					—	M.fr.IX,4.
PHYSOPSIS; Turczaninow in Bullet. de la Soc. Imper. des Natur. de Mosc. XXII, 34 (1849).													
P. spicata; Turczaninow in Bull. de Mosc. XXII, 35 (1849)	W.A.	—	—	—	—	—	—					B.fl.V,41	M.fr.VI,154.
MALLOPHORA; Endlicher in Annal. des Wien. Mus. II, 206 (1838). (Lachnocephalus.)													
M. globiflora; Endlicher in Ann. des Wien. Mus. II, 206 (1838)	W.A.	—	—	—	—	—	—					B.fl.V,41	M.fr.VI,154; X,15.
DICRASTYLIS; Drummond & Harvey in Hook. Kew Misc. VII, 56 (1855). (Dicrastyles.)													
D. Doranii; F. v. M., fragm. VIII, 230 (1874)	—	S.A.	—	—	—	—	—					—	M.fr.VIII,230.
D. ochrotricha; F. v. M., fragm. IV, 161 (1864)	—	S.A.	—	—	—	—	N.A.					B.fl.V,42	M.fr.I,60; VI,155; VIII,230; IX,4.
D. Gilesii; F. v. M., fragm. VIII, 229 (1874)	—	S.A.	—	—	—	—	—					—	M.fr.X,15.
D. fulva; Drummond & Harvey in Kew Misc. VII, 56 (1855)	W.A.	—	—	—	—	—	—					B.fl.V,43	M.fr.I,236; II,160; VI,155.
D. reticulata; Drummond & Harvey in Hook. Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	—					B.fl.V,43	M.fr.VI,155; IX,4.
D. Nicholasii; F. v. M., fragm. X, 15 (1876)	W.A.	—	—	—	—	—	—					—	M.fr.X,15.
D. parvifolia; F. v. M., fragm. II, 16 (1861)	W.A.	—	—	—	—	—	—					B.fl.V,43	M.fr.VI,155; X,15.
D. Beveridgei; F. v. M., fragm. VIII, 50 (1873)	—	S.A.	—	—	—	—	—					—	M.fr.IX,4.
D. Stoechas; Drummond & Harvey in Kew Misc. VII, 57 (1855)	W.A.	—	—	—	—	—	—					B.fl.V,44	M.fr.VI,155.
D. Lewellini; F. v. M., fragm. XI, 86 (1880)	—	S.A.	—	—	—	—	—					—	M.fr.VIII,50; XI,137.
CHLOANTHES; R. Brown, prodr. Flor. Nov. Holl. 513 (1810). (Pityrodia, Quoya, Dasymalla, Depremesnilia.)													
C. Stoechadis; R. Brown, prodr. 514 (1810)	—	—	—	—	N.S.W.	—	—					B.fl.V,45	M.fr.VI,156.
C. parviflora; Walpers, repertor. bot. syst. IV, 58 (1848)	—	—	—	V.	N.S.W.	Q.	—					B.fl.V,46	M.fr.IX,5.
C. coccinea; Bartling in Lehm. pl. Preiss. I, 352 (1845)	W.A.	—	—	—	—	—	—					B.fl.V,48	M.fr.VI,156; IX,5.
C. salvifolia; F. v. M., Pityrodia, R. Brown, prodr. 513 (1810)	—	—	—	—	—	Q.	—					B.fl.V,48	M.fr.VI,155,255; IX,5.
C. hemigenioides; F. v. M., fragm. VI, 156 (1868)	W.A.	—	—	—	—	—	N.A.					B.fl.V,48	M.fr.VI,156.
C. uncinata; Turczaninow in Bull. de Mosc. XXXVI, 194 (1863)	W.A.	—	—	—	—	—	—					B.fl.V,48	M.fr.VI,156,
C. Bartlingii; Lehmann, Ind. sem. hort. Hamb. (1844)	W.A.	—	—	—	—	—	—					B.fl.V,49	M.fr.VI,156.
C. verbascina; F. v. M., fragm. I, 233 (1859)	W.A.	—	—	—	—	—	—					B.fl.V,50	M.fr.IV,80; VI,157; IX,5.
C. stachyodes; F. v. M., fragm. V, 50 (1855)	W.A.	—	—	—	—	—	—					B.fl,V,50	M.fr.VI,158.
C. loxocarpa; F. v. M., fragm. II, 22 (1861)	W.A.	—	—	—	—	—	N.A.					B.fl.V,51	M.fr.IV,80; VI,156.
C. dilatata; F. v. M., fragm. VI, 157 (1868)	W.A.	—	—	—	—	—	—					B.fl.V,51	M.fr.VI,157.
C. cuneata; F. v. M., Cens. 103 (1882)	W.A.	—	—	—	—	—	—					B.fl.V,51	
C. Oldfieldii; F. v. M., fragm. I, 234 (1859)	W.A.	—	—	—	—	—	—					B.fl.V,52	M.fr.II,182; IV,80.
C. atriplicina; F. v. M., fragm. I, 235 (1859)	W.A.	—	—	—	—	—	N.A.					B.fl.V,52	M.fr.VI,157; IX,5.
C. paniculata; F. v. M., fragm. IV, 80 (1864)	W.A.	—	—	—	—	—	N.A.					B.fl.V.53	M.fr.IV,80.
C. lepidota; F. v. M. in Wing's S. Sc. Rec. III, 3 (1883)	W.A.	—	—	—	—	—	—					—	
C. loricata; F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—					—	M.fr.X,14.
C. halganiacea; F. v. M., fragm. X, 14 (1876)	W.A.	—	—	—	—	—	—					—	M.fr.X,14.
C. Depremesnilii; Depremesnilia chrysocalyx, F. v. M. X, 59 (1876)	W.A.	—	—	—	—	—	—					—	M.fr.X,59.
NESOGENES; A. de Candolle, prodr. XI, 703 (1847).													
N. euphrasioides; A. de Candolle, prodr. XI, 703 (1847)	—	—	—	—	—	Q.	—					—	

HEMIPHORA ; F. v. M., fragm. X, 13 (1876).

H. Elderi ; F. v. M., fragm. X, 13 (1876)	W.A.	—	—	—	—	—	—	—					M.fr.X,13.		

DENISONIA ; F. v. M. in Journ. Linn. Soc. II, 159 (1858).

D. ternifolia ; F. v. M., fragm. I, 124, t. 2 (1859)	—	—	—	—	—	—	N.A.	B.fl.V,54					M.fr.I,124.		

CYANOSTEGIA ; Turczaninow in Bull. Soc. Mosc. XXII, 35 (1849). (Bunnya.)

C. Turczaninowii ; F. v. M., fragm. VI, 154 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,54					M.fr.IX,5.		
C. Bunnyana ; F. v. M., fragm. V, 36 (1865)	—	—	—	—	—	—	N.A.	B.fl.V,54					M.fr.VI,153.		

CALLICARPA ; Linné in act. litter. et scien. Upsal. 80 (1741).

C. cana ; Linné, mantisa. plant. alt. 198 (1771)	—	—	—	—	—	Q.	N.A.	B.fl.V,56			...	...	Aa., P.		
C. pedunculata ; R. Brown, prodr. 513 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,57							
C. longifolia ; Lamarck, Encycl. méth. I, 562 (1783)	—	—	—	—	—	Q.	—	B.fl.V,57			...	...	As.		

TATEA ; F. v. M., in Transact. Roy. Soc. of S. Austr. VI, 33 (1883).

T. acaulis ; F. v. M. in Transact. R. Soc. of S. Austr. VI, 34 (1883)	—	—	—	—	—	—	N.A.	—							

PREMNA ; Linné, mantiss. plant. alter. 154 (1771).

P. obtusifolia ; R. Brown, prodr. 512 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,58	M.fr.III,36.						
P. integrifolia ; Linné, mant. plant. alt. 154 (1771)	—	—	—	—	—	Q.	N.A.	B.fl.V,59	M.fr.VI,153		...	As., P.			
P. limbata ; Bentham, Fl. Austr. V, 59 (1870)	—	—	—	—	—	Q.	—	B.fl.V,59							
P. Dallachyana ; Bentham, Fl. Austr. V, 59 (1870)	—	—	—	—	—	Q.	—	B.fl.V,59							
P. acuminata ; R. Brown, prodr. 512 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,60	M.fr.III,36.						

CLERODENDRUM ; Burmann, thesaur. Zeylan. 66, t. 29 (1737). (Clerodendron, Glossocarya.)

C. hemiderma ; F. v. M. in Benth. Fl. Austr. V, 61 (1870)	—	—	—	—	—	Q.	—	B.fl.V,61	M.fr.VI,151; IX,5.						
C. inerme ; Gaertner, de fruct. I, 271, t. 57 (1788)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,61	M.fr.VI,152		...	As., P.			
C. Tracyanum ; F. v. M. in Benth. Fl. Austr. V, 62 (1870)	—	—	—	—	—	Q.	—	B.fl.V,62	M.fr.V,61.						
C. tomentosum ; R. Brown, prodr. 510 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,62	M.fr.VI,152.						
C. lanceolatum ; F. v. M., fragm. III, 145 (1863)	W.A.	—	—	—	—	—	N.A.	B.fl.V,63	M.fr.III,145.						
C. floribundum ; R. Brown, prodr. 511 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,63	M.fr.III,145; IX,5.			P.			
C. Cunninghamii ; Bentham, Fl. Austr. V, 64 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,64							
C. costatum ; R. Brown, prodr. 511 (1810)	—	—	—	—	—	Q.	—	B.fl.V,64							

GMELINA ; Linné, gen. plant. ed. sec. 526 (1742).

G. macrophylla ; Bentham, Fl. Austr. V, 65 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.V,65	M.fr.IV,128.						
G. fasciculiflora ; Bentham, Fl. Austr. V, 65 (1870)	—	—	—	—	—	Q.	—	B.fl.V,65	M.fr.V,35.						
G. Leichhardtii ; F. v. M. in Benth. Fl. Austr. V, 66 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,66	M.fr.III,58,158; VI,153.						

VITEX ; Tournefort, inst. 603, t. 373 (1700), from Plinius.

V. trifolia ; Linné, spec. plant. 638 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,66	M.fr.III,59; VI,152; IX,5.			Aa., P.			
V. Lignum vitae ; Cunningham in De Candolle, prodr. XI, 692 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,67	M.fr.III,58; IX,5.						
V. acuminata ; R. Brown, prodr. 512 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.V,67	M.fr.V,34; VI,152; IX,5.						
V. glabrata ; R. Brown, prodr. 512 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.V,68	M.fr.III,59; V,35; VI,153.						

FARADAYA ; F. v. M., fragm. phytogr. Austr. V, 21 (1865).

F. splendida ; F. v. M., fragm. V, 21 (1865)	—	—	—	—	—	Q.	—	B.fl.V,69	M.fr.VI,153.						

AVICENNIA ; Linné, syst. nat. 8 (1735); Linné, gen. plant. 27 (1737).

A. officinalis ; Linné, spec. plant. 110 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,69			...	...	As., Afr., Am., P., N.Z.		

MYOPORINAE.

R. Brown, prodr. 514 (1810).—Genera 2, Species 76.

EREMOPHILA; R. Brown, prodr. 518 (1810). (Pholidia, Stenochilus, Eremodendron, Pseudopholidia, Sentis, Duttonia, Pholidiposis.)

E. Mackinlayi; F. v. M., fragm. IV, 80 (1864)	W.A.	—	—	—	—	—	—	B.fl.V,17	M.fr.V,109.
E. Bowmani; F. v. M., fragm. II, 139 (1863)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,18	M.fr.II,139; III,166.
E. strongylophylla; F. v. M., fragm. X, 87 (1876)	W.A.	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.X,87.
E. leucophylla; Bentham, Fl. Austr. V, 18 (1870)	W.A.	S.A.	—	—	—	—	—	B.fl.V,18	M.fr.X,87.
E. Forrestii; F. v. M., fragm. VII, 49 (1869)	W.A.	—	—	—	—	—	—	B.fl.V,18	M.fr.VII,49.
E. eriocalyx; F. v. M., fragm. I, 236 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,19	M.fr.I,236.
E. Maitlandi; F. v. M. in Benth. Fl. Austr. V, 19 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,19	
E. rotundifolia; F. v. M., fragm. I, 207 (1859)	—	S.A.	—	—	—	—	—	B.fl.V,19	M.fr.I,207.
E. oppositifolia; R. Brown, prodr 518 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,20	
E. Mitchelli; Bentham in Mitch. Trop. Austr. 31 (1848)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,21	M.fr.VI,148.
E. Paisleyi; F. v. M., Rep. Babb. Exped. 17 (1858)	—	S.A.	—	—	—	—	—	B.fl.V,20	
E. Sturtii; R. Brown, App. to Sturt's Exped. 22 (1849)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.V,21	M.fr.VI,148.
E. exilifolia; F. v. M., fragm. X, 88 (1876)	W.A.	S.A.	—	—	—	—	—	—	M.fr.X,88.
E. Dempsteri; F. v. M., fragm. X, 60 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,60.
E. Gibsoni; F. v. M., fragm. VIII, 227 (1874)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,227.
E. Berryi; F. v. M., fragm. VIII, 228 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,228; X,88.
E. Clarkei; F. v. M., fragm. I, 208 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,21	M.fr.I,208.
E. Gilesii; F. v. M., fragm. VIII, 49 (1873)	—	S.A.	—	—	—	—	—	—	M.fr.VIII,49.
E. Latrobei; F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,22	M.fr.I,125.
E. graciliflora; F. v. M., fragm. I, 208 (1858)	W.A.	—	—	—	—	—	—	B.fl.V,23	M.fr.II,181.
E. longifolia; F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,23	M.fr.VI,148.
E. Laanii; F. v. M. in Melb. Chem. Jan. (1885)	W.A.	—	—	—	—	—	—	—	
E. Drummondii; F. v. M., fragm. VI, 147 (1868)	W.A.	—	—	—	—	—	—	B.fl.V,24	M.fr.VI,147.
E. Hughesii; F. v. M., fragm. VIII, 228 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,228.
E. polyclada; F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,24	
E. bignoniflora; F. v. M. in Proceed. Roy. Soc. Tasm. III, 294 (1858)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,25	M.fr.VI,148.
E. Freelingii; F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.V,25	M.fr.VI,148.
E. Fraseri; F. v. M., fragm. XI, 51 (1878)	W.A.	—	—	—	—	—	N.A.	—	M.fr.XI,51.
E. Macdonnelli; F. v. M., Rep. Babb. Exped. 18 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,22	M.fr.V,23; VI,150.
E. Goodwinii; F. v. M., Rep. Babb. Exped. 17 (1858)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.V,25	M.fr.VI,148.
E. Elderi; F. v. M., fragm. VIII, 228 (1874)	W.A.	S.A.	—	—	—	—	—	—	M.fr.VIII,228.
E. Willsii; F. v. M., fragm. III, 21, t. 20 (1862)	W.A.	S.A.	—	—	—	Q.	—	B.fl.V,26	M.fr.III,21.
E. platycalyx; F. v. M., fragm. V, 109 (1866)	W.A.	—	—	—	—	—	—	B.fl.V,26	M.fr.V,109.
E. viscida; Endlicher, nov. stirp. dec. 51 (1839)	W.A.	—	—	—	—	—	—	B.fl.V,26	
E. Brownii; F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,27	M.fr.VI,148.
E. subfloccosa; Bentham, Fl. Austr. V, 28 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,28	
E. Oldfieldii; F. v. M., fragm. I, 208 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,28	M.fr.I,208.
E. Duttonii; F. v. M., Rep. Babb. Exped. 16 (1858)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,28	M.fr.VI,148.
E. maculata; F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A,	B.fl.V,29	M.fr.I,125.
E. denticulata; F. v. M., fragm. I, 125 (1859)	W.A.	S.A.	—	—	—	—	—	B.fl.V,29	M.fr.IV,170.
E. latifolia; F. v. M. in Schlechtend. Linnaea XXV, 428 (1852)	W.A.	S.A.	—	—	N.S.W.	—	—	B.fl.V,30	M.fr.V,109; VI,148.
E. alternifolia; R. Brown, prodr. 518 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,30	M.fr.V,109; VI,148.
E. Dalyana; F. v. M., fragm. V, 22 (1865)	—	S.A.	—	—	N.S.W.	—	—	B.fl.V,10	M.fr.V,22.
E. Pantoni; F. v. M. in Wing's S. Sc. Rec. II, 25 (1882)	—	—	—	—	—	—	N.A.	—	
E. scoparia; F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,10	M.fr.V,22; VI,148.
E. Delisserii; F. v. M., fragm. V, 108, t. 42 (1865)	W.A.	S.A.	—	—	—	—	—	B.fl.V,11	M.fr.V,189.

E. crassifolia; F. v. M. in Proceed. Roy. Soc. Tasm. III, 297 (1858)	—	S.A.	—	—	—	—	—	B.fl.V,11	M.fr.VI,148.				
E. resinosa; F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)	W.A.	—	—	—	—	—	—	B.fl.V,11					
E. Behrii; F. v. M. in Proceed. Roy. Soc. Tasm. III, 296 (1858)...	—	S.A.	—	—	—	—	—	B.fl.V,12					
E. Woollsiana; F. v. M., fragm. I, 125, t. 7 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,12	M.fr.VI,151.				
E. Christophori; F. v. M., fragm. IX, 120 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,198.				
E. brevifolia; F. v. M., fragm. I, 126 (1859)	W.A.	—	—	—	—	—	—	B.fl.V,12	M.fr.VI,150.				
E. Weldii; F. v. M., fragm. VII, 109 (1870)	W.A.	S.A.	—	—	—	—	—	—	M.fr.IX,120.				
E. densifolia; F. v. M., fragm. II, 160 (1861)	W.A.	—	—	—	—	—	—	B.fl.V,13	M.fr.II,160.				
E. gibbosifolia; F. v. M., Rep. Babb. Exped. 18 (1858)	—	S.A.	—	V.	—	—	—	B.fl.V,13	M.fr.VI,150.				
E. divaricata; F. v. M. in Proceed. Roy. Soc. Tasm. III, 293 (1858)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,14	M.fr.IV,48;VI,150.				
E. microtheca; F. v. M. in Benth. Fl. Austr. V, 14 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,14	M.fr.II,160.				
E. adenotricha; F. v. M. in Benth. Fl. Austr. V, 15 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,14					
E. santalina; F. v. M. in Proceed. Roy. Soc. Tasm. III, 295 (1858)	—	S.A.	—	—	—	—	—	B.fl.V,15					
E. Youngii; F. v. M., fragm. X, 16 (1876)	W.A.	—	—	—	—	—	—	—	M.fr.X,16.				
MYOPORUM; Banks & Solander in G. Forster, prodr. 44 (1786). (Bertolonia, Andrewsia, Pogonia, Disoon.)													
M. tenuifolium; G. Forster, fl. ins. Austr. prodr. 44 (1786)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.V,4	M.fr.VI,149	...	P.		
M. obscurum; Endlicher, prodr. fl. Norf. 54 (1833)...	—	—	—	—	N.S.W.	—	—	—					
M. Dampieri; Cunningham in De Candolle, prodr. XI, 708 (1847)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.V,4	M.fr.VI,149.				
M. deserti; Cunningham in Hueg. enum. 78 (1837)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.V,5	M.fr.VI,148; VII,110.				
M. laxiflorum; Bentham, Fl. Austr. V, 6 (1870)	—	—	—	—	—	Q.	—	B.fl.V,6	M.fr.V,23.				
M. insulare; R. Brown, prodr. 515 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,6	M.fr.VI,149; VII,110.				
M. viscosum; R. Brown, prodr. 516 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.V,5	M.fr.VII,109.				
M. serratum; R. Brown, prodr. 516 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,4	M.fr.VI,149; VII,110.				
M. oppositifolium; R. Brown, prodr. 516 (1810)	W.A.	—	—	—	—	—	—	B.fl.V,7	M.fr.VI,149.				
M. brevipes; Bentham, Fl. Austr. V, 6 (1870)	—	S.A.	—	—	—	—	—	B.fl.V,6					
M. humile; R. Brown, prodr. 516 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.V,6	M.fr.VI,150.				
M. debile; R. Brown, prodr. 516 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.V,6	M.fr.VI,149.				
M. platycarpum; R. Brown, prodr. 516 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.V,7	M.fr.VI,150.				
M. floribundum; Cunningham in Hueg. enum. 78 (1837)	—	—	—	V.	N.S.W.	—	—	B.fl.V,8	M.fr.I,126; VI,150.				
M. Beckeri; F. v. M. in Benth. Fl. Austr. V, 7 (1870)	W.A.	—	—	—	—	—	—	B.fl.V,7	M.fr.I,156,244; VI,150.				
M. salsoloides; Turczaninow in Bull. de Mosc. XXXVI, 226 (1863)	W.A.	—	—	—	—	—	—	B.fl.V,8	M.fr.I,126; VI,150.				

ASPERIFOLIAE.

HALLER, enum. stirp. Helv. Praef. 34 (1742).—Genera 12, Species 52.

CYNOGLOSSUM; Tournefort, inst. 139, t. 57 (1700) from Dioscorides and Nicandros.													
C. latifolium; R. Brown, prodr. 495 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,408	M.fr.VI,115; IX.122; XI,125.				
C. suaveolens; R. Brown, prodr. 495 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,409	M.fr.VI,115; IX,122.				
C. Australe; R. Brown, prodr. 495 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,409	M.fr.VI,115; IX,122.				
C. Drummondii; Bentham, Fl. Austr. IV, 409 (1869)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,409	M.fr.IX,122.				
ROCHELIA; Reichenbach in Fl. Regensb. bot. Zeit. I, 243 (1824). (Maccoya.)													
R. Maccoya; F. v. M. in Benth. Fl. Austr. IV, 408 (1869)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,408	M.fr.I,127; IX,121.				
LAPPULA; Rivinus in Rupp. Fl. Jenens. (1718), from Dalechamp. (Echinospermum.)													
L. concava; F. v. M., Cens. 100 (1882)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,407	M.fr.VI,116; IX,121.				
ERITRICHUM; Schrader in commentat. Goetting. IV, 186 (1820). (Eritrichium, Lappulae subgenus.)													
E. Australasicum; A. de Candolle, prodr. X, 134 (1846)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.IV,406	M.fr.XI,125.				

MYOSOTIS; Ruppius, Fl. Jenens. 9 (1718), from Dioscorides and Plinius. (Exarrhena.)

M. Australis; R. Brown, prodr. 495 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,405	M.fr.VI,116; IX,122	...	N.Z.
M. exarrhena; F. v. M., inedit.	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,406	M.fr.VI,115; IX,122.		

POLLICHIA; Medikus, Bot. Beobach. 247 (1783). (Trichodesma.)

P. Zeylanica; F. v. M., Cens. 100 (1882) from N. Burmann (1768)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.IV,404	M.fr.VI,115; IX,122; XI,125.		As., Afr., P.
P. latisepalea; F. v. M., Cens. 100 (1882)	—	—	—	—	—	Q.	N.A.	—	M.fr.X,102,121.		

HALGANIA; Gaudichaud in Freyc. voy. Bot. 448 (1826).

H. solanacea; F. v. M. in Hook. Kew Misc. IX, 21 (1857)	—	—	—	—	—	—	N.A.	B.fl.IV,401	M.fr.VI,115.		
H. litoralis; Gaudichaud in Freyc. voy. Bot. 449, t. 59 (1826)	W.A.	—	—	—	—	—	—	B.fl.IV,401	M.fr.VI,115.		
H. corymbosa; Lindley, Bot. Regist. XXV, App. 40 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,402			
H. sericiflora; Bentham, Fl. Austr. IV, 402 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,402			
H. cyanea; Lindley, Bot. Regist. XXV, App. 40 (1839)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,402	M.fr.VI,115; IX,122.		
H. lavandulacea; Endlicher in Ann. Wien. Mus. II, 205 (1838)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,402	M.fr.I,209; VI,115; XI,125.		
H. integerrima; Endlicher in Ann. Wien. Mus. II, 205 (1838)	W.A.	—	—	—	—	—	—	B.fl.IV,402			

HELIOTROPIUM; Tournefort, inst. 138, t. 57 (1700) from Theophrastos, Dioscorides & Plinius. (Tiaridium, Heliophytum, Schleidenia restorable.)

H. Indicum; Linné, spec. plant. 130 (1753)	—	—	—	—	—	Q.	N.A.	—	M.fr.XI,124	...	As., Afr., Am.
H. pleiopterum; F. v. M., fragm. IX, 121 (1875)	—	S.A.	—	—	—	—	—	—	M.fr.IX,121.		
H. Curassavicum; Linné, spec. plant. 130 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,393	M.fr.VI,116; IX,121	...	Afr., Am., P.
H. Europaeum; Linné, spec. plant. 130 (1753)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,394	M.fr.VI,116	...	E., As., Afr.
H. undulatum; Vahl, symbol. bot. I, 13 (1795)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.IV,394	M.fr.VI,116; IX,121; XI,124.		As., Afr.
H. asperrimum; R. Brown, prodr. 493 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.IV,394	M.fr.VI,116; IX,121.		
H. crispatum; F. v. M. in Benth. Fl. Austr. IV, 395 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,395			
H. fasciculatum; R. Brown, prodr. 494 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,395			
H. vestitum; Bentham, Fl. Austr. IV, 395 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,395			
H. brachygyne; Bentham, Fl. Austr. IV, 396 (1869)	—	—	—	—	—	Q.	—	B.fl.IV,396			
H. epacrideum; F. v. M. in Benth. Fl. Austr. IV, 396 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,396			
H. ovalifolium; Forskael, Fl. Aegypt. Arab. 47 (1775)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,396	M.fr.VI,115	...	As., Afr.
H. strigosum; Willdenow, sp. pl. I, 743 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.IV,397	...	...	As., Afr.
H. prostratum; R. Brown, prodr. 494 (1810)...	—	—	—	—	—	—	N.A.	B.fl.IV,397			
H. bracteatum; R. Brown, prodr. 493 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,397			
H. pauciflorum; R. Brown, prodr. 493 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,398			
H. conocarpum; F. v. M. in Benth. Fl. Austr. 398 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,398			
H. filaginoides; Bentham, Fl. Austr. IV, 398 (1869)	—	S.A.	—	—	—	—	—	B.fl.IV,398			
H. ventricosum; R. Brown, prodr. 494 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,399			
H. paniculatum; R. Brown, prodr. 494 (1810)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.IV,399			
H. Cunninghamii; Bentham, Fl. Austr. IV, 400 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,400			
H. diversifolium; F. v. M. in Benth. Fl. Austr. IV, 400 (1869)	—	—	—	—	—	—	N.A.	B.fl.IV,400			

COLDENIA; Linné, amoen. acad. ed. prim. 119 (1747). (Lobophyllum.)

C. procumbens; Linné, spec. plant. 125 (1753)	—	S.A.	—	—	—	Q.	N.A.	B.fl.IV,391	M.fr.VI,115	...	As., Afr.

TOURNEFORTIA; Linné, syst. nat. 8 (1735); Linné, gen. plant. 55 (1737).

T. argentea; Linné, fil. suppl. plant. 133 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.IV,389	M.fr.IV,95; VI,116	...	As., Afr., P.
T. mollis; F. v. M., fragm. I, 59 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.IV,390	M.fr.VI,116.		
T. sarmentosa; Lamarck, Illustr. des genr. I, 416 (1791)	—	—	—	—	—	Q.	—	B.fl.IV,390	M.fr.IV,95; VI,116	...	P.

EHRETIA; P. Browne, Civ. and Nat. Hist. of Jamaic. 168, t. 16 (1756).

E. acuminata; R. Brown, prodr. 497 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,387	M.fr.V,21; IX,122; XI,124.		As.
E. pilosula; F. v. M., fragm. V, 20 (1865)	—	—	—	—	—	Q.	—	B.fl.IV,388	M.fr.V,20.		
E. saligna; R. Brown, prodr. 497 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.IV,388	M.fr.V,21; IX,122.		

E. membranifolia; R. Brown, prodr. 497 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,388	M.fr.V,21; VI,115; IX,122.						
E. laevis; Roxburgh, Pl. Corom. I, 42, t. 56 (1795) ...	—	—	—	—	—	Q.	—	B.fl.IV,389	..	..	...	...	As.		
CORDIA; Plumier, gen. plant. 13, t. 14 (1703).															
C. subcordata; Lamarck, Illustr. des genr. I, 421 (1791) ...	—	—	—	—	—	Q.	N.A.	B.fl.IV,385	M.fr.VI,114; IX,122	..	As., Afr., P.				
C. aspera; G. Forster, florul. ins. Austr. prodr. 18 (1786) ...	—	—	—	—	—	Q.	—	B.fl.IV,386	M.fr.V,193; VI,114; IX,122.	P.					
C. Myxa; Linné, spec. plant. 190 (1753) ...	—	—	—	—	—	Q.	—	B.fl.IV,386	M.fr.I,59; VI,114...	...	As., Afr., P.				

ERICACEAE.

A. L. de Jussieu, Gen. plant. 159 (1789)—Genera 5, Species 7.

RHODODENDRON; Linné, spec. plant. I, 392 (1753).															
R. Lochae; F. v. M. in Vict. Naturalist III, 157 (1887) ...	—	—	—	—	—	Q.	—	—							
PERNETTYA; Gaudichaud in Ann. des sc. nat. V, 102 (1825).															
P. Tasmanica; J. Hooker in Lond. Journ. of Bot. VI, 268 (1847) ...	—	—	T.	—	—	—	—	B.fl.IV,140	...	...	...	N.Z.			
GAULTIERA; Kalm in Linné, amoen. acad. III, 14, fig. 6 (1751). (Gaultheria.)															
G. hispida; R. Brown, prodr. 559 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,141							
G. lanceolata; J. Hooker in Lond. Journ. of Bot. VI, 267 (1847)...	—	—	T.	—	—	—	—	B.fl.IV,141							
G. antipoda; G. Forster, florul. ins. Austr. pr. 34 (1786) ...	—	—	T.	—	—	—	—	B.fl.IV,142	...	...	...	N.Z.			
AGAPETES; D. Don in G. Don, gen. syst. of dich. plants III, 862 (1834). (Vaccinii subgenus.)															
A. Meiniana; F. v. M. in Vict. Naturalist III, 158 (1887)...	—	—	—	—	—	Q.	—	—							
WITTSTEINIA; F. v. M., fragm. II, 136 (1861).															
W. vacciniacea; F. v. M., fragm. II, 136 (1861) ...	—	—	—	V.	—	—	—	B.fl.IV,139	M.fr.III,166.						

EPACRIDEAE.

R. Brown, prodr. 535 (1810).—Genera 18, Species 275.

STYPHELIA; Solander in G. Forster, fl. ins. Austr. prodr. 13 (1786). (Epacris, Forst. (1776) partly, Stiphelia, Ardisia, Peroa, Perojoa, Ventenatia, Cyathodes, Stenanthera, Astroloma, Leucopogon, Melichrus, Acrotriche, Monotoca, Soleniscia, Stomarrhena, Pentataphrus, Mesotriche, Phanerandra, Froebelia, Pentaptelion, Androstoma, Cyathopsis, Lissanthe partly.)

S. adscendens; R. Brown, prodr. 537 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,146	M.fr.VI,36; XI,122.						
S. longifolia; R. Brown, prodr. 537 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,147							
S. laeta; R. Brown, prodr. 537 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,147							
S. triflora; Andrews, Bot. Reposit. t. 72 (1799) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,147	M.fr.XI,122.						
S. viridis; Andrews, Bot. Reposit. t. 312 (1806) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,148							
S. Hainesii; F. v. M., fragm. IV, 96, t. 28 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,148	M.fr.VIII,55.						
S. tubiflora; Smith, Specim. of the Bot. of New Holl. 45, t. 14 (1793) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,148							
S. tenuiflora; Lindley, Bot. Regist. XXV, App. 25 (1839)...	W.A.	—	—	—	—	—	—	B.fl.IV,148	M.fr.IV,97; VI,46; VIII,54.						
S. melaleucoides; F. v. M., fragm. IV, 97 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,149	M.fr.VI,30.						
S. pusilliflora; F. v. M., fragm. IV, 105 (1864) ...	—	S.A.	—	—	—	—	—	B.fl.IV,149	M.fr.I,178; VI,31.						
S. Leucopogon; F. v. M., fragm. IV, 97 (1864) ...	W.A.	—	—	—	—	—	—	B.fl.IV,149	M.fr.III,143; VI,31.						
S. lasionema; F. v. M., fragm. VI, 40 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VIII,54.						
S. macrocalyx; F. v. M., fragm. VI, 37 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VI,37.						
S. xerophylla; F. v. M., fragm. VI, 38 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VI,38.						
S. pentapogonea; F. v. M., fragm. VI, 36 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,153	M.fr.VI,36.						
S. prostrata; F. v. M., Cens. 105 (1882) from R. Brown (1810) ...	W.A.	—	—	—	—	—	—	B.fl.IV,154							
S. tecta; Sprengel, syst. I, 657 (1825)...	W.A.	—	—	—	—	—	—	B.fl.IV,154	M.fr.VI,37.						

S. Candolleana; F. v. M., fragm. VI, 38 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,154	M.fr.VIII,54.
S. microdonta; F. v. M., Cens. 105 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,155	
S. pallida; Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,155	M.fr.VI,37.
S. compacta; Sprengel, syst. I, 657 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,155	M.fr.VI,38.
S. humifusa; Persoon, synops. plant. I, 174 (1805)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,156	M.fr.VI,37; VIII,54.
S. Epacridis; F. v. M., fragm. VI, 38 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,156	M.fr.VI,38.
S. Drummondii; F. v. M., fragm. VI, 37 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,157	M.fr.VI,37.
S. microcalyx; F. v. M., fragm. VI, 37 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,157	M.fr.VI,37.
S. Baxteri; F. v. M., fragm. VI, 37 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,158	M.fr.IV,97.
S. Sonderi; F. v. M., fragm. VI, 36 (1867)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,158	M.fr.VIII,54; IX,48; XI,122.
S. longiflora; F. v. M., fragm. VIII, 54 (1873)	W.A.	—	—	—	—	—	—	B.fl.IV,158	M.fr.VIII,54.
S. pinifolia; Sprengel, syst. veg. I, 659 (1825)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,159	M.fr.IV,99; VI,36.
S. procumbens; Persoon, synops. I, 174 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,162	M.fr.VI,38; VIII,55.
S. urceolata; F. v. M., fragm. VI, 38 (1867)	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,162	M.fr.VIII,54.
S. Billardieri; F. v. M., fragm. VI, 43 (1867)	—	—	T.	—	—	—	—	B.fl.IV,169	M.fr.VIII,54.
S. Milligani; F. v. M., Pentschondra verticillata, J. Hook. Fl. Tas. I, 256 (1860)	—	—	T.	—	—	—	—	B.fl.IV,164	
S. straminea; Sprengel, syst. I, 656 (1825)	—	—	T.	—	—	—	—	B.fl.IV,169	M.fr.VI,43.
S. Hookeri; F. v. M., fragm. VI, 44 (1867)	—	—	T.	—	—	—	—	B.fl.IV,169	M.fr.VI,44; VIII,54.
S. dealbata; Sprengel, syst. I, 659 (1825)	—	—	T.	—	—	—	—	B.fl.IV,170	M.fr.VI,43; VIII,54.
S. abietina; Labillardière, Nov. Holl. pl. spec. I, 48, t. 68 (1804)	—	—	T.	—	—	—	—	B.fl.IV,170	M.fr.VI,43.
S. Oxycedrus; Labillardière, Nov. Holl. pl. spec. I, 49, t. 69 (1804)	—	—	T.	V.	—	—	—	B.fl.IV,170	M.fr.IV,100; VI,43 ... N.Z.
S. parvifolia; F. v. M. in papers of R. S. Tasm. 86 (1874)	—	—	T.	—	—	—	—	B.fl.IV,171	
S. sapida; F. v. M., fragm. VI, 42 (1867)	—	—	—	—	N.S.W.	—	—	B.fl.IV,175	M.fr.VI,42; VIII,54.
S. strigosa; Smith, Specim. of the Bot. of New Holl. 49 (1793)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,176	M.fr.VI,42; IX,48; XI,122.
S. verticillata; Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,184	M.fr.IV,122; VI,43; VIII,55.
S. interrupta; Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,184	
S. amplexicaulis; Rudge in Transact. Linn. Soc. VIII, 292, t. 8 (1805)	—	—	—	—	N.S.W.	—	—	B.fl.IV,185	M.fr.VI,44.
S. alternifolia; Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,185	
S. lanceolata; Smith, Specim. of the Bot. of New Holl. 49 (1793)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,185	M.fr.IV,124; VI,43.
S. Richei; Labillardière, Nov. Holl. pl. spec. I, 44, t. 60 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,186	M.fr.IV,123; VI,42; VIII,54. N.Z.
S. Australis; F. v. M., fragm. VI, 43 (1867)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,186	M.fr.IV,123.
S. capitellata; F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,187	M.fr.VI,31.
S. revoluta; Sprengel, syst. I, 657 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,187	M.fr.VI,31.
S. grandiuscula; F. v. M., fragm. VI, 47 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,188	M.fr.VI,47.
S. reflexa; Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,188	M.fr.VI,32.
S. corifolia; F. v. M., Cens. 105 (1882) from Endlicher (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,188	
S. distans; Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,189	M.fr.VI,32.
S. gibbosa; F. v. M., Cens. 106 (1882) from Stschegleew (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,189	
S. thymifolia; F. v. M., Cens. 106 (1882) from Lindley (1869)	—	—	—	V.	—	—	—	B.fl.IV,189	
S. cordata; F. v. M., Cens. 106 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,190	
S. Bossiaea; F. v. M., fragm. VI, 47 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,190	M.fr.VI,47.
S. hirsuta; F. v. M., fragm. VI, 31 (1867)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,191	M.fr.VI,31.
S. collina; Labillardière, Nov. Holl. pl. spec. I, 47, t. 65 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,191	M.fr.VI,45; VIII,54.
S. glacialis; F. v. M., Cens. 106 (1882) from Lindley (1838)	—	—	—	V.	—	—	—	B.fl.IV,191	
S. compacta; F. v. M., Cens. 106 (1882) from Stschegl. (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,192	
S. squarrosa; F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,192	M.fr.VI,31.
S. microphylla; Sprengel, syst. I, 656 (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,192	M.fr.VI,31.
S. tetragona; F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,193	M.fr.VI,31.
S. phyllostachys; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,193	M.fr.VI,32.
S. glabella; Sprengel, syst. I, 655 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,194	M.fr.VI,32.
S. semiopposita; F. v. M., fragm. VI, 49 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,194	M.fr.VI,49.

S. florulenta; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,194	
S. striata; Sprengel, syst. I, 656 (1825)	W.A.	S.A.	—	—	—	—	—	B.fl.IV,194	
S. costata; F. v. M. in Wing's S. Sc. Rec. I, new ser. (1885)	—	S.A.	—	V.	—	—	—	—	
S. lasiostachya; F. v. M., Cens. 106 (1882) from Stschegl. (1859)...	W.A.	—	—	—	—	—	—	B.fl.IV,195	
S. carinata; Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,195	
S. opponens; F. v. M., fragm. VI, 48 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,196	M.fr.VI,48.
S. oppositifolia; F. v. M., fragm. VI, 32 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,196	M.fr.VI,32.
S. tamariscina; Sprengel, syst. I, 656 (1825)...	W.A.	—	—	—	—	—	—	B.fl.IV,197	
S. bracteolaris; F. v. M., Cens. 106 (1882) from Bentham (1869)...	W.A.	—	—	—	—	—	—	B.fl.IV,197	
S. blepharophylla; F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,197	M.fr.VI,34.
S. tenuis; F. v. M., Cens. 106 (1882) from De Candolle (1833)	W.A.	—	—	—	—	—	—	B.fl.IV,197	
S. gnaphalioides; F. v. M., Cens. 106 (1882) from Stschegl. (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,198	
S. concurva; F. v. M., fragm. VI, 36 (1867)...	—	S.A.	—	—	—	—	—	B.fl.IV,198	M.fr.III,144.
S. Gilbertii; F. v. M., Cens. 106 (1882) from Stschegl. (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,198	
S. gracilis; Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,199	
S. acicularis; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,199	
S. cryptantha; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,199	
S. gracillima; F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,200	M.fr.VI,34.
S. cymbiformis; F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,200	M.fr.VI,34.
S. apiculata; Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,201	M.fr.III,144; IV,105; VI,31.
S. polystachya; Sprengel, syst. I, 659 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,201	M.fr.VI,31.
S. virgata; Labillardière, Nov. Holl. pl. spec. I, 46, t. 64 (1804)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,201	M.fr.VI,42.
S. pulchella; F. v. M., fragm. VI, 34 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,202	M.fr.VI,34.
S. polymorpha; F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,202	M.fr.VI,31.
S. assimilis; F. v. M., Cens. 106 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,202	M.fr.IV,106; VI,32.
S. Oldfieldii; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,203	
S. cucullata; Sprengel, syst. I, 656 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,203	M.fr.VI,32.
S. sprengelioides; F. v. M., Cens. 106 (1882) from Sonder (1845)...	W A.	—	—	—	—	—	—	B.fl.IV,204	M.fr.VI,34.
S. obtusata; F. v. M., Cens. 106 (1882) from Sonder (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,204	
S. fimbriata; F. v. M., Cens. 106 (1882) from Stschegl. (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,204	
S. ozothamnoides; F. v. M., Cens. 106 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,204	M.fr.VI,155.
S. plumiflora; F. v. M., fragm. VI, 29 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,205	M.fr.XI,122.
S. unilateralis; F. v. M., Cens. 106 (1882) from Stschegl. (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,205	
S. montana; F. v. M., fragm. VI, 45 (1867)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,206	M.fr.VI,45.
S. Macraei; F. v. M., fragm. VI, 46 (1867)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,206	M.fr.VI,46; VIII,54.
S. pleurandroides; F. v. M., fragm. VI, 32 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,206	M.fr.III,143.
S. linifolia; F. v. M., fragm. VI, 36 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207	M.fr.VI,36.
S. pluriloculata; F. v. M., fragm. VI, 32 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207	M.fr.I,37.
S. pleiosperma; F. v. M., fragm. VI, 41 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,207	M.fr.VI,41.
S. rubicunda; F. v. M., fragm. VI, 31 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,208	M.fr.IV,99.
S. attenuata; F. v. M., Cens. 106 (1882) from Cunningham (1825)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,208	
S. conferta; F. v. M., Cens. 106 (1882) from Bentham (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,208	
S. mutica; F. v. M., fragm. VI, 45 (1867)	—	—	—	—	N.S.W.	—	—	B.fl.IV,209	M.fr.VI,45.
S. ericoides; Smith, Specim. of the Bot. of New Holl. 48 (1793)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,209	M.fr.VI,45.
S. brevicuspis; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,210	
S. propinqua; Sprengel, syst. I, 658 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,210	M.fr.IV,103; VI,34.
S. subulifolia; F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,211	M.fr.IV,103.
S. Allittii; F. v. M., fragm. VI, 34 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,211	M.fr.IV,103.
S. racemulosa; F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,211	M.fr.IV,103.
S. pendula; Sprengel, syst. I, 657 (1825)	W.A.	—	—	—	—	—	—	B.fl.IV,212	M.fr.VI,33.
S. concinna; F. v. M., Cens. 106 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,212	

S. margarodes ; Sprengel, syst. I, 657 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,213	M.fr.VI,36.		
S. flavescens ; F. v. M., fragm. VI, 33 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,213	M.fr.IV,100.		
S. blepharolepis ; F. v. M., fragm. VI, 48 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,213	M.fr.VI,48.		
S. esquamata ; Sprengel, syst. I, 658 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,214			
S. rotundifolia ; Sprengel, syst. I, 655 (1825)...	W.A.	—	—	—	—	—	—	B.fl.IV,214			
S. cordifolia ; F. v. M., fragm. VIII, 54 (1873)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,214	M.fr.IV,105; VI,45.		
S. megacarpa ; F. v. M., fragm. VI, 32 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,215	M.fr.IV,102.		
S. ruscifolia ; Sprengel, syst. I, 656 (1825) ...	—	—	—	—	—	Q.	—	B.fl.IV,215			
S. imbricata ; Sprengel, syst. I, 656 (1825) ...	—	—	—	—	—	Q.	—	B.fl.IV,215			
S. cuspidata ; Sprengel, syst. I, 657 (1825) ...	—	—	—	—	—	Q.	—	B.fl.IV,216			
S. leptospermoides ; Sprengel, syst. I, 659 (1825)	—	—	—	—	—	Q.	—	B.fl.IV,216			
S. acuminata ; Sprengel, syst. I, 659 (1825) ...	—	—	—	—	—	—	N.A.	B.fl.IV,216			
S. flexifolia ; Sprengel, syst. I, 659 (1825) ...	—	—	—	—	—	Q.	—	B.fl.IV,217			
S. biflora ; Sprengel, syst. I, 659 (1825) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,217	M.fr.IV,104.		
S. setigera ; Sprengel, syst. I, 659 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,217	M.fr.VI,45.		
S. exolasia ; F. v. M., fragm. VI, 34 (1867) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,217	M.fr.VI,34.		
S. Fraseri ; F. v. M., Cens. 107 (1882) from Cunningham (1840) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,218	M.fr.IV,105; VI,46	...	N.Z.
S. hirtella ; F. v. M., Cens. 107 (1882)...	—	S.A.	—	—	—	—	—	B.fl.IV,218			
S. Oldfieldii ; F. v. M., Cens. 107 (1882) ...	W.A.	—	—	—	—	—	—	B.fl.IV,218			
S. erubescens ; F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,219	M.fr.IV,102.		
S. lissanthoides ; F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,219	M.fr.IV,101.		
S. stricta ; F. v. M., Cens. 107 (1882) from Bentham (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,219			
S. Mitchellii ; F. v. M., Cens. 107 (1882) from Bentham (1869) ...	—	—	—	—	—	Q.	—	B.fl.IV,220			
S. juniperina ; Sprengel, syst. I, 658 (1825) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.IV,220	M.fr.IV,104; VI,46.		
S. rufa ; F. v. M., fragm. VI, 46 (1867) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,221	M.fr.VIII,54; IX,48.		
S. conostephioides ; F. v. M., fragm. VI, 34 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,221	M.fr.IV,98.		
S. deformis ; Sprengel, syst. I, 658 (1825) ...	—	—	—	—	N.S.W.	—	—	B.fl.IV,221			
S. pogonocalyx ; F. v. M., Cens. 107 (1882) ...	W.A.	—	—	—	—	—	—	B.fl.IV,222			
S. breviflora ; F. v. M., Cens. 107 (1882) ...	W.A.	—	—	—	—	—	—	B.fl.IV,222	M.fr.IV,102.		
S. dura ; F. v. M., Cens. 107 (1882) from Bentham (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,222			
S. multiflora ; Sprengel, syst. I, 658 (1825) ...	W.A.	—	—	—	—	—	—	B.fl.IV,222	M.fr.VI,32.		
S. appressa ; Sprengel, syst. I, 658 (1825) ...	—	—	—	V.	N.S.W.	—	—	B.fl.IV,223			
S. Neo-Anglica ; F. v. M., Cens. 107 (1882) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,223			
S. obtecta ; F. v. M., Cens. 107 (1882) from Bentham (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,223			
S. crassifolia ; F. v. M., fragm. VI, 40 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,224	M.fr.VI,40.		
S. strongylophylla ; F. v. M., fragm. VI, 33 (1867) ...	W.A.	—	—	—	—	—	—	B.fl.IV,224	M.fr.IV,101.		
S. crassiflora ; F. v. M., fragm. VI, 33 (1867)...	W.A.	—	—	—	—	—	—	B.fl.IV,224	M.fr.VI,33.		
S. corynocarpa ; F. v. M., Cens. 107 (1882) from Sonder (1845) ...	W.A.	—	—	—	—	—	—	B.fl.IV,224			
S. Woodsii ; F. v. M., Cens. 107 (1882) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,225	M.fr.VI,33.		
S. leptantha ; F. v. M., Cens. 107 (1882) from Bentham (1869) ...	W.A.	—	—	—	—	—	—	B.fl.IV,225			
S. ericifolia ; F. v. M.; Pentachondra ; J. Hook. in Lond. Journ. VI, 271 (1847)	—	—	T.	—	—	—	—	B.fl.IV,164			
S. divaricata ; Sprengel, syst. I, 658 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,226	M.fr.VI,44.		
S. aggregata ; Sprengel, syst. I, 657 (1825) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,227	M.fr.VI,44; VIII,55.		
S. serrulata ; Labillardière, Nov. Holl. pl. spec. I, 45, t. 62 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,227	M.fr.VI,44.		
S. patula ; Sprengel, syst. I, 657 (1825) ...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,227	M.fr.VI,44; XI,122.		
S. ovalifolia ; Sprengel, syst. I, 656 (1825) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.IV,228	M.fr.VI,44.		
S. ramiflora ; Sprengel, syst. I, 659 (1825) ...	W.A.	—	—	—	—	—	—	B.fl.IV,228	M.fr.		
S. depressa ; Sprengel, syst. I, 655 (1825) ...	W.A.	S.A.	—	—	—	—	—	B.fl.IV,228	M.fr.VI,44.		
S. fasciculiflora ; F. v. M., fragm. VIII, 55 (1873) ...	—	S.A.	—	—	—	—	—	B.fl.IV,229	M.fr.VI,44.		
S. elliptica ; Smith, Specim. of the Bot. of New Holl. 49 (1793) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,230	M.fr.VI,58.		
S. glauca ; Labillardière, Nov. Holl. pl. spec. I, 45, t. 61 (1804) ...	—	—	T.	—	—	—	—	B.fl.IV,230			

S. scoparia; Smith, Specim. of the Bot. of New Holl. 49 (1793)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.IV,231	M.fr.VI,58; XI,122.				
S. ledifolia; F. v. M., Cens. 107 (1882) from Cunningham (1839)	—	—	—	—	N.S.W.	—	—	B.fl.IV,231					
S. empetrifolia; F. v. M. in Papers of R. S. Tasm. 86 (1874)	—	—	T.	—	—	—	—	B.fl.IV,231					
S. oligarrhenoides; F. v. M., fragm. IX, 47 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,47.				
S. minutiflora; F. v. M., fragm. XI, 122 (1881)	W.A.	—	—	—	—	—	—	B.fl.IV,231	M.fr.XI,122.				
OLIGARRHENA; R. Brown, prodr. Fl. Nov. Holl. 549 (1810).													
O. micrantha; R. Brown, prodr. 459 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,232					
NEEDHAMIA; R. Brown, prodr. Fl. Nov. Holl. 549 (1810).													
N. pumilio; R. Brown, prodr. 549 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,174					
BRACHYLOMA; Sonder in Lehm. pl. Preiss. I, 304 (1845). (Lobopogon, Lissanthe partly.)													
B. daphnoides; Bentham, Fl. Austr. IV, 173 (1869)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,173	M.fr.VI,42; VIII,55.				
B. ciliatum; Bentham, Fl. Austr. IV, 173 (1869)	—	S.A.	T.	V.	—	—	—	B.fl.IV,173	M.fr.VI,42.				
B. depressum; Bentham, Fl. Austr. IV, 173 (1869)	—	—	T.	V.	—	—	—	B.fl.IV,173	M.fr.I,36; VI,42.				
B. Scortechinii; F. v. M., fragm. XI, 121 (1881)	—	—	—	—	—	Q.	—	—	M.fr.XI,121.				
B. Preissii; Sonder in Lehm. pl. Preiss. I, 305 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,172	M.fr.VI,39; VIII,55.				
B. concolor; F. v. M., fragm. VI, 39 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,172	M.fr.IV,98.				
B. ericoides; Sonder in Schlecht. Linnaea XXVI, 247 (1853)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.IV,172	M.fr.IV,98; VI,39; XI,46.				
CONOSTEPHIUM; Bentham in Hueg. enum. 76 (1837). (Conostephiopsis.)													
C. pendulum; Bentham in Hueg. enum. 76 (1837)	W.A.	—	—	—	—	—	—	B.fl.IV,160	M.fr.VI,40.				
C. minus; Lindley, Bot. Regist. XXV, App. 25 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,160	M.fr.VI,40.				
C. Roei; Bentham, Fl. Austr. IV, 160 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,160	M.fr.VIII,55.				
C. Preissii; Sonder in Lehm. pl. Preiss. I, 304 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,161	M.fr.VI,40.				
C. planifolium; F. v. M., fragm. VI, 30 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,161	M.fr.VI,30.				
COLEANTHERA; Stschegleew in Bull. Soc. Mosc. XXXII, 4 (1859). (Michiea.)													
C. coelophylla; Bentham, Fl. Austr. IV, 150 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,150					
C. myrtoides; Stschegleew in Bull. de Mosc. XXXII, 4 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,150	M.fr.IV,96; VI,80.				
C. virgata; Stschegleew in Bull. de Mosc. XXXII, 5 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,151					
TROCHOCARPA; R. Brown, prodr. 548 (1810). (Decaspora, Pentachondra.)													
T. laurina; R. Brown, prodr. 548 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,166	M.fr.VI,57; IX,48.				
T. disticha; Sprengel, syst. I, 660 (1825)	—	—	T.	—	—	—	—	B.fl.IV,166	M.fr.VI,57; VIII,55.				
T. thymifolia; Sprengel, syst. I, 660 (1825)	—	—	T.	—	—	—	—	B.fl.IV,166	M.fr.VI,57; VIII,55.				
T. Clarkei; F. v. M., fragm. VI, 57 (1867)	—	—	—	V.	—	—	—	B.fl.IV,167	M.fr.VI,57.				
T. Gunnii; Bentham, Fl. Austr. IV, 167 (1869)	—	—	T.	—	—	—	—	B.fl.IV,167	M.fr.VIII,55.				
T. parviflora; Bentham, Fl. Austr. IV, 167 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,167.					
T. involucrata; F. v. M., fragm. VI, 57 (1867)	—	—	T.	—	—	—	—	B.fl.IV,164	M.fr.VIII,55.				
T. pumila; F. v. M., fragm. VI, 57 (1867)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,164	M.fr.VI,57	N.Z.			
EPACRIS; Cavanilles, icon. et descr. pl. IV, 25, t. 344 et 345 (1797). (Archeria.)													
E. micranthera; F. v. M., fragm. VI, 72 (1867)	—	—	T.	—	—	—	—	B.fl.IV,246	M.fr.VIII,56.				
E. hirtella; J. Hooker in Lond. Journ. of Bot. VI, 271 (1847)	—	—	T.	—	—	—	—	B.fl.IV,246	M.fr.VI,71.				
E. longiflora; Cavanilles, icon. et descr. pl. IV, 25, t. 344 (1797)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,234	M.fr.VI,70; VIII,56.				
E. reclinata; Cunningham in Benth. Fl. Austr. IV, 234 (1869)	—	—	—	—	N.S.W.	—	—	B.fl.IV,234	M.fr.VIII,56.				
E. impressa; Labillardière, Nov. Holl. pl. spec. I, 43, t. 58 (1804)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,235	M.fr.VI,70; VIII,56; XI,122.				
E. sparsa; R. Brown, prodr. 551 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,235	M.fr.VIII,53.				
E. petrophila; J. Hooker, Fl. Tasman. I, 261 (1860)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,236					
E. rigida; Sieber in Spreng. cur. poster. 64 (1827)	—	—	—	—	N.S.W.	—	—	B.fl.IV,236					
E. coriacea; Cunningham in De Candolle, prodr. VII, 763 (1839)	—	—	—	—	N.S.W.	—	—	B.fl.IV,236					
E. crassifolia; R. Brown, prodr. 551 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,237	M.fr.IV,127.				

E. robusta; Bentham, Fl. Austr. IV, 237 (1869)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,237	
E. obtusifolia; Smith, Exot. Bot. I, 77, t. 40 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,237	M.fr.VI,71; VIII,56.
E. myrtifolia; Labillardière, Nov. Holl. pl. spec. I, 41, t. 55 (1804)	—	—	T.	—	—	—	—	B.fl.IV,238	
E. exserta; R. Brown, prodr. 551 (1810)	—	—	T.	—	—	—	—	B.fl.IV,238	
E. mucronulata; R. Brown, prodr. 552 (1810)	—	—	T.	—	—	—	—	B.fl.IV,238	
E. lanuginosa; Labillardière, Nov. Holl. pl. spec. I, 42, t. 57 (1804)	—	S.A.	T.	V.	—	—	—	B.fl.IV,238	M.fr.VIII,56; XI,122.
E. paludosa; R. Brown, prodr. 551 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.IV,239	M.fr.VIII,56.
E. Calvertiana; F. v. M., fragm. VIII, 53 (1874)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,53.
E. heteronema; Labillardière, Nov. Holl. pl. spec. I, 42, t. 56 (1804)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,239	M.fr.IV,127; VI,71; VIII,56. N.Z.
E. serpillifolia; R. Brown, prodr. 240 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.IV,240	
E. microphylla; R. Brown, prodr. 550 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.IV,240	
E. acuminata; Bentham, Fl. Austr. IV, 240 (1869)	—	—	T.	—	—	—	—	B.fl.IV,240	
E. apiculata; Cunningham in Field, N. S. Wales, 340 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.IV,241	
E. pulchella; Cavanilles, icon. IV, 26, t. 345 (1797)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,241	M.fr.VI,71; XI,122.
E. purpurascens; R. Brown, prodr. 550 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,241	M.fr.IV,127; VI,254; VIII,56.
WOOLLSIA; F. v. M., fragm. VIII, 52 (1872). (Lysinema partly.)									
W. pungens; F. v. M., fragm. VIII, 52 (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,243	M.fr.IV,126; VI,70,88; VIII,55; IX,48; XI,122.
LYSINEMA; R. Brown, prodr. 552 (1810).									
L. lasianthum; R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,243	M.fr.III,142; VI,70; VIII,55.
L. conspicuum; R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,243	M.fr.VI,70; VIII,55; XI,122.
L. ciliatum; R. Brown, prodr. 552 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,243	M.fr.VI,69,70; VIII,55.
L. fimbriatum; F. v. M., fragm. IV, 125 (1864)	W.A.	—	—	—	—	—	—	B.fl.IV,244	M.fr.VI,70.
L. elegans; Sonder in Lehm. pl. Preiss. I, 327 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,244	M.fr.VI,70.
PRIONOTES; R. Brown, prodr. fl. Nov. Holl. 553 (1810).									
P. cerinthoides; R. Brown, prodr. 553 (1810)	—	—	T.	—	—	—	—	B.fl.IV,246	M.fr.VI,69; XI,122.
COSMELIA; R. Brown, prodr. fl. Nov. Holl. 553 (1810).									
C. rubra; R. Brown, prodr. 553 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,247	M.fr.VI,64.
PONCELETIA; R. Brown, prodr. fl. Nov. Holl. 554 (1810).									
P. sprengelioides; R. Brown, prodr. 554 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.IV,248	M.fr.I,39; VI,60.
P. monticola; Cunningham in De Candolle, prodr. VII, 768 (1843)	—	—	—	—	N.S.W.	—	—	B.fl.IV,249	M.fr.VI,60.
SPRENGELIA; Smith in Svensk. Vetens. Acad. Handling. 260 (1794). (Poiretia.)									
S. incarnata; Smith in Vetens. Ac. Handl. 260, t. 8 (1794)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.IV,249	M.fr.I,39; VI,54; IX,48; VIII,56; XI,122.
ANDERSONIA; R. Brown, prodr. 553 (1810). (Atherocephala, Homalostoma, Sphincterostoma.)									
A. colossea; F. v. M., fragm. VI, 63 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,251	M.fr.VI,63.
A. patricia; F. v. M., fragm. VI, 79 (1868)	W.A.	—	—	—	—	—	—	B.fl.IV,251	M.fr.VI,79.
A. grandiflora; Stschegleew in Bull. de Mosc. XXXII, 21 (1859)	W.A.	—	—	—	—	—	—	B.fl.IV,252	M.fr.VI,62.
A. setifolia; Bentham, Fl. Austr. IV, 252 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,252	M.fr.VIII,51.
A. involucrata; Sonder in Lehm. pl. Preiss. I, 331 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,252	M.fr.VI,62.
A. homalostoma; Bentham, Fl. Austr. IV, 253 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,253	M.fr.VIII,63.
A. sprengelioides; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,253	M.fr.VI,64.
A. latiflora; F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,253	M.fr.VI,61.
A. gracilis; De Candolle, prodr. VII, 767 (1843)	W.A.	—	—	—	—	—	—	B.fl.IV,254	
A. aristata; Lindley, Bot. Regist. XXV, App. 25 (1839)	W.A.	—	—	—	—	—	—	B.fl.IV,254	M.fr.VI,56.
A. macronema; F. v. M., fragm. VIII, 51 (1873)	W.A.	—	—	—	—	—	—	—	M.fr.VIII,51.
A. parvifolia; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,254	M.fr.IV,125; VI,62.
A. depressa; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,255	
A. coerulea; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,255	M.fr.VI,64.

A. subulata; Bentham, Fl. Austr. IV, 256 (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,256					
A. heterophylla; Sonder in Lehm. pl. Preiss. I, 333 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,256	M.fr.IV,124.				
A. brachyanthera; F. v. M., fragm. VI, 61 (1867)	W.A.	—	—	—	—	—	—	B.fl.IV,256	M.fr.VI,61.				
A. brevifolia; Sonder in Lehm. pl. Preiss. I, 332 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,256	M.fr.VI,61.				
A. variegata; Sonder in Lehm. pl. Preiss. I, 334 (1845)	W.A.	—	—	—	—	—	—	B.fl.IV,257					
A. micrantha; R. Brown, prodr. 554 (1810)	W.A.	—	—	—	—	—	—	B.fl.IV,257	M.fr.VI,61.				
RICHEA; R. Brown, prodr. 555 (1810). (Cystanthe, Pilitis.)													
R. sprengelioides; F. v. M., fragm. VI, 68 (1867)	—	—	T.	—	—	—	—	B.fl.IV,258	M.fr.I,38; VI,68.				
R. procera; F. v. M., fragm. VI, 68 (1867)	—	—	T.	—	—	—	—	B.fl.IV,259	M.fr.I,38; VI,68.				
R. acerosa; F. v. M., fragm. VI, 69 (1867)	—	—	T.	—	—	—	—	B.fl.IV,259	M.fr.I,38; VI,69.				
R. Milligani; F. v. M., fragm. VI, 69 (1867)..	—	—	T.	—	—	—	—	B.fl.IV,259	M.fr.I,38; VIII,56.				
R. Gunnii; J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T.	V.	—	—	—	B.fl.IV,260	M.fr.VI,67; XI,122.				
R. scoparia; J. Hooker in Lond. Journ. VI, 273 (1847)	—	—	T.	—	—	—	—	B.fl.IV,260	M.fr.VI,68.				
R. dracophylla; R. Brown, prodr. 555 (1810)	—	—	T.	—	—	—	—	B.fl.IV,260	M.fr.VI,67.				
R. pandanifolia; J. Hooker, Fl. Antarct. I, 50 (1844)	—	—	T.	—	—	—	—	B.fl.IV,261	M.fr.VI,67; IX,48.				
DRACOPHYLLUM; Labillardière, Voy. II, 211, t. 40 (1798). (Epacris, Forst. (1776) partly.)													
D. Milligani; Hooker, icon. t. 845 (1852)	—	—	T.	—	—	—	—	B.fl.IV,262	M.fr.VI,65.				
D. Fitzgeraldi; F. v. M., fragm. VII, 27 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,280; X,119.				
D. Sayeri; F. v. M. in Aust. Journ. of Pharm. new ser. II, 85 (1887)	—	—	—	—	—	Q.	—	—					
D. secundum; R. Brown, prodr. 556 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.IV,262	M.fr.VI,65; VIII,56.				
D. minimum; F. v. M., fragm. I, 39 (1858)	—	—	T.	—	—	—	—	B.fl.IV,265	M.fr.VI,65; VII,56	...	N.Z.		
SPHENOTOMA; Sweet, Flora Australasica, t. 44 (1828), from R. Brown (1810).													
S. squarrosum; G. Don, gen. syst. III, 785 (1834)	W.A.	—	—	—	—	—	—	B.fl.IV,263	M.fr.VI,66.				
S. Drummondii; F. v. M., Cens. 109 (1882) from Bentham (1869)	W.A.	—	—	—	—	—	—	B.fl.IV,263					
S. dracophylloides; Sonder in Lehm. pl. Preiss. I, 33 (1844)	W.A.	—	—	—	—	—	—	B.fl.IV,264	M.fr.VI,65.				
S. capitatum; Lindley, Bot. Regist. t. 1515 (1832)	W.A.	—	—	—	—	—	—	B.fl.IV,264					
S. gracile; Swest, Fl. Austral. t. 44 (1828)	W.A.	—	—	—	—	—	—	B.fl.IV,264	M.fr.VI,66.				
S. parviflorum; F. v. M., Cens. of Aust. pl. 109 (1882)	W.A.	—	—	—	—	—	—	B.fl.IV,265	M.fr.VI,66.				

APETALEAE GYMNOSPERMAE.

F. v. M. in Woolls's plants of the neighb. of Sydney, 40 (1880).

CONIFERAE.

Haller, enum. stirp. Helv. I, 145 (1742).—Genera 10, Species 29.

DAMMARA; Rumphius, herb. Amboin. II, 174, t. 54 (1741) from Sherard (1704). (Agathis.)													
D. robusta; C. Moore in Transact. pharm. Soc. Vict. II, 174 (1860)	—	—	—	—	—	Q.	—	B.fl.VI,244	M.fr.XI,104.				
ARAUCARIA; A. L. de Jussieu, gen. plant. 413 (1789). (Colymbea, Columbea, Eutassa, Eutacta, Altingia.)													
A. Cunninghamii; Aiton in Sweet. hort. Brit. 475 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,243	...	...	..	P.	
A. Bidwilli; Hooker, Lond. Journ. of Bot. II, 503, t. 18 (1850)	—	—	—	—	—	Q.	—	B.fl.VI,243					
A. excelsa; R. Brown in Ait. hort. Kew, sec. ed. V, 412 (1813)	—	—	—	—	N.S.W.	—	—	—	M.fr.IX,169.				
ATHROTAXIS; D. Don in Transact. Linn. Soc. XVIII, 171 (1839). (Arthrotaxis, Sequoia.)													
A. cupressoides; D. Don in Trans. Linn. Soc. XVIII, 173, t. 13 (1839)	—	—	T.	—	—	—	—	B.fl.VI,242					
A. laxifolia; Hooker, icon. plant. t. 573 (1843)	—	—	T.	—	—	—	—	B.fl.VI,242					
A. selaginoides; D. Don in Transact Linn. Soc. XVIII, 172, t. 14 (1839) ...	—	—	T.	—	—	—	—	B.fl.VI,242	M.fr.XI,104.				

CALLITRIS ; Ventenat, decas generum novorum 10 (1808). (Frenela, Actinostrobus, Leichhardtia, Octoclinis.)

C. Macleayana ; F. v. M. in Rep. Burdek. Exped. 17 (1860)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,235	M.fr.XI,104.
C. Parlatorei ; F. v. M. in Seemann, Journ. of Bot. 267 (1866)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,235	M.fr.V,186; XI,104.
C. Roei ; Bentham & J. Hooker, gen. plant. III, 424 (1880) implied	W.A.	—	—	—	—	—	—	B.fl.VI,236	
C. Drummondii ; Bentham & J. Hooker, gen. plant. III, 424 (1880)	W.A.	—	—	—	—	—	—	B.fl.VI,236	
C. verrucosa ; R. Brown in Mém. du Mus. Par. XIII, 74 (1826)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,236	M.fr.V,237.
C. columellaris ; F. v. M., fragm. V, 198 (1866)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.V,198.
C. Muelleri ; Bentham & J. Hooker, gen. plant. III, 424 (1880)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,237	
C. cupressiformis ; Ventenat, decas. gen. nov. 10 (1808)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,237	
C. calcarata ; R. Brown in Mém. du Mus. Par. XIII, 74 (1826)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,238	
C. oblonga ; L. Cl. Richard, comment. de conif. 49, t. 18 (1826)	—	—	T.	—	—	—	—	B.fl.VI,238	
C. Actinostrobus ; F. v. M., Rep. Burdek. Exped. 19 (1860)	W.A.	—	—	—	—	—	—	B.fl.VI,240	
C. acuminata ; F. v. M., Cens. 109 (1882) from Parlatore (1862)	W.A.	—	—	—	—	—	—	B.fl.VI,240	M.fr.XI,104.

FITZROYA ; J. Hooker in Journ. of the hort. Soc. VI, 264 (1851). (Diselma.)

F. Archeri ; Bentham & J. Hooker, gen. pl. III, 426 (1880)	—	—	T.	—	—	—	—	B.fl.VI,240	

PHEROSPHAERA ; Archer in Hooker's Kew Misc. II, 52 (1850). (Dacrydium partly.)

P. Hookeriana ; Archer in Hooker, Kew Misc. II, 52 (1850)	—	—	T.	—	—	—	—	B.fl.VI,245	M.fr.XI,103.
P. Fitzgeraldi ; F. v. M. in J. Hooker, icon. pl. XIV, 64, t. 1383 (1882)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,102,138.

MICROCACHRYS ; J. Hooker in Lond. Journ. of Bot. IV, 149 (1845). (Dacrydium partly.)

M. tetragona ; J. Hooker, Fl. Tasman. I, 358, t. 100 (1860)	—	—	T.	—	—	—	—	B.fl.VI,241	

DACRYDIUM ; Solander in G. Forster, pl. escul. insul. ocean. austr. 80 (1786).

D. Franklinii ; J. Hooker in Lond. Journ. of Bot. IV, 152, t. 6 (1845)	—	—	T.	—	—	—	—	B.fl.VI,245	

NAGEIA ; Gaertner, de fructib. t. 39 (1788). (Podocarpus.)

N. elata ; F. v. M., Cens. 109 (1882) from R. Brown (1826)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,247	M.fr.IV,87.
N. spinulosa ; F. v. M., Cens. 109 (1882) from R. Brown (1826)	—	—	—	—	N.S.W.	—	—	B.fl.VI,247	
N. Drouyniana ; F. v. M., Cens. of Aust. pl. 109 (1882)	W.A.	—	—	—	—	—	—	B.fl.VI,247	M.fr.IV,86; VI,251; XI,104.
N. alpina ; F. v. M. in papers Roy. Soc. of Tasm. 23 (1879)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,248	M.fr.IV,87; XI,104.

THALAMIA ; Sprengel, Anleit. zur Kenntn. der Gew. zweite Ausg. II, 218 (1817). (Phyllocladus.)

T. asplenifolia ; Sprengel, Anl. zur Kenntn. der. Gew. II, 218 (1817)	—	—	T.	—	—	—	—	B.fl.VI,246	M.fr.XI,104.

CYCADEAE.

L. C. Richard in Persoon, synops. 630 (1807).—Genera 3, Species 14.

CYCAS ; Linné, hort. Cliffort. 482 (1737).

C. media ; R. Brown, prodr. 348 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,249	M.fr.VIII,171.
C. Normanbyana ; F. v. M., fragm. VIII, 169 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,169,283.
C. Kennedyana ; F. v. M. in Melb. Chemist, March (1882)	—	—	—	—	—	Q.	—	—	
C. Cairnsiana ; F. v. M., fragm. X, 63 (1876)	—	—	—	—	—	Q.	—	—	M.fr.X,63,121.

ENCEPHALARTOS ; Lehmann, pugill. VI, 9, t. I, et 3 (1834). (Macrozamia, Lepidozamia, Catakidozamia, Zamia partly.)

E. Pauli Guilielmi ; F. v. M. in Trans. Pharm. Soc. Vict. II, 91 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,251	M.fr.I,86,243; II,179; IX,124.
E. tridentatus ; Lehmann, pugill. VI, 13 (1834)	—	—	—	—	N.S.W.	—	—	B.fl.VI,253	M.fr.III,38; XI,125.
E. spiralis ; Lehmann, pugill. VI, 13 (1834)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,251	M.fr.I,41; II,179; VIII,173.
E. Douglasii ; F. v. M., Cens. 110 (1882)	—	—	—	—	—	Q.	—	—	[126.
E. Fraseri ; Miqnel in Versl. K. Akad. Wet. Amst. XV, 368 (1863)	W.A.	—	—	—	—	—	—	B.fl.VI,252	M.fr.I,41,243; II,179; III,167; VIII,173; XI,
E. Moorei ; F. v. M., fragm. XI, 125 (1881)	—	—	—	—	—	Q.	—	—	M.fr.I,243; XI,125.
E. Macdonnelli ; F. v. M. in Versl. K. Akad. Wet. Amst. XV, 376 (1863)	—	S.A.	—	—	—	—	—	—	M.fr.IX,124; XI,126.
E. Dyeri ; F. v. M. in Melb. Chemist, June (1885)	W.A.	—	—	—	—	—	—	—	
E. Denisonii ; F. v. M. in Journ. Pharm. Soc. Vict. II, 90 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,253	M.fr.I,41; VIII,173; IX,124.

BOWENIA ; Hooker, Bot. Magaz. t. 5398 (1863).

B. spectabilis ; Hooker, Bot. Mag. t. 5398 (1863)	—	—	—	—	—	Q.	—	B.fl.VI,254	M.fr.V,171; VIII,279.

MONOCOTYLEDONEAE.

Ray, meth. plant. nov. (1682).

EUCALYCEAE PERIGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 41 (1880).

SCITAMINEAE.

Linne, philos. botan. 27 (1751).—Genera 7, Species 11.

MUSA; l'Ecluse, éxoticorum libri decem 229 (1605).

M. Banksii; F. v. M., fragm. IV, 132 (1864)...	—	—	—	—	—	Q.	—	B.fl.VI,261	M.fr.IX,190.		
M. Fitzalani; F. v. M., fragm. IX, 188 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,188.		
M. Hillii; F. v. M., fragm. IX, 169 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,169.		

CURCUMA; Linné, gen. plant. 332 (1737) from Hermann (1687).

C. Australasica; J. Hooker, Bot. Mag. t. 5620 (1867)	—	—	—	—	—	Q.	—	B.fl.VI,263	M.fr.VIII,26.		

AMOMUM; Linné, gen. plant. 330 (1737).

A. Dallachyi; F. v. M., fragm. VIII, 25 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,263	M.fr.VIII,25.		

ELETTARIA; White & Matton in Transact. Linn. Soc. X, 249 (1809) from Rheede (1692).

E. Scottiana; F. v. M., fragm. VIII, 24 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,264	M.fr.VIII,24.		

ALPINIA; Linné, gen. plant. 332 (1737). (Hellenia.)

A. racemigera; F. v. M., fragm. VIII, 27 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,265	M.fr.VIII,27.		
A. coerulea; Bentham, Fl. Austr. VI, 265 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,265	M.fr.VIII,26.		
A. arctiflora; F. v. M., fragm. VIII, 25 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,266	M.fr.VIII,25.		

COSTUS; C. Bauhin, pinax 36 (1623).

C. Potierae; F. v. M., fragm. IV, 164 (1864)...	—	—	—	—	—	Q.	—	B.fl.VI,266	M.fr.VIII,27.		

TAPEINOCHEILOS; Miquel in annal. mus. bot. Lugdun. IV, 101, t. 4 (1869).

T. pungens; Miquel in ann. mus. Lugd. IV, 101, t. 4 (1869)	—	—	—	—	—	Q.	—	B.fl.VI,267	M.fr.VIII,26	...	As.

ORCHIDEAE.

Haller, enum. stirp. Helv. praefat. 33 (1742).—Genera 49, Species 271.

PHOLIDOTA; Lindley in Hooker, Exotic Flora II, t. 138 (1825).

P. imbricata; Lindley in Hook. Exot. Fl. II, t. 138 (1825)	—	—	—	—	—	Q.	—	B.fl.VI,290	M.fr.VI,163; V,105	...	As.

STURMIA; Reichenbach in Moessler's Handb. II, 1552 (1828). (Liparis, Richard 1818 not of Zoologists 1738.)

S. reflexa; F. v. M., fragm. II, 72 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,273	M.fr.III,165; V,102.		
S. angustilabris; F. v. M., fragm. IV, 164 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,273	M.fr.IV,164.		
S. coelogynoides; F. v. M., fragm. II, 71 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,273	M.fr.IX,50.		
S. habenarina; F. v. M., fragm. IV, 131 (1864)	—	—	—	—	—	Q.	—	B.fl.VI,273	M.fr.V,102; XI,53.		

OBERONIA; Lindley, gen. and spec. of orchid. pl. 15 (1830). (Titania.)

O. iridifolia; Lindley, gen. and spec. of orchid. pl. 15 (1830)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,274	M.fr.II,24.		
O. Titania; Lindley, fol. orchidac. VIII, 8 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,274	M.fr.II,24; VII,30; XI,53,88.		

PHREATIA; Lindley, gen. and spec. of orchid. pl. 63 (1830). (Plexaure.)

P. Limenophylax; G. Reichenbach in Seem. Bonpl. 54 (1857)	—	—	—	—	—	Q.	—	B.fl.VI,290	M.fr.X,64.					

DENDROBIUM; Swartz in nov. act. Upsal. VI, 82 (1799). (Thelychiton partly, Pedilonum, Coelandria.)

D. bigibbum; Lindley in Paxt. Flower Gard. III, 25 (1853)	—	—	—	—	—	Q.	N.A.	B.fl.VI,277	M.fr.VI,119; XI,87.					
D. dicuphum; F. v. M., fragm. VIII, 28 (1871)	—	—	—	—	—	—	N.A.	B.fl.VI,277	M.fr.VI,119.					
D. Sumneri; F. v. M., fragm. VI, 94 (1868)	—	—	—	—	—	Q.	—	B.fl.VI,278	M.fr.VI,94.					
D. superbiens; G. Reichenbach in Garden. Chron. 516 (1876)	—	—	—	—	—	Q.	—	—	M.fr.XI,22.					
D. undulatum; R. Brown, prodr. 332 (1810)...	—	—	—	—	—	Q.	—	B.fl.VI,278	M.fr.I,87; III,39; XI,87.	As.				
D. Johannis; G. Reichenbach in Garden. Chron. 890 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,279						
D. speciosum; Smith, Exot. Botany I, 17, t. 10 (1804)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,279	M.fr.I,180; VIII,249.					
D. falconirostre; Fitzgerald, Austr. Orch. I, pt. V (1879)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,87.					
D. tetragonum; Cunningham in Bot. Regist. XXV, Misc. 33 (1839)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,279	M.fr.I,87; VI,119.					
D. aemulum; R. Brown, prodr. 333 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,280	M.fr.I,213; XI,87.					
D. Adae; Bailey in Proceed. Roy. Soc. of Q. I, pt. 3 (1884)	—	—	—	—	—	Q.	—	—						
D. Eriae; D. eriaeoides, Bailey, papers on the Queensl. Flora. II, Sept. (1886)...	—	—	—	—	—	Q.	—	—						
D. Fitzgeraldi; F. v. M. in Melb. Chemist, June (1884)	—	—	—	—	—	Q.	—	—						
D. macropus; Bentham & J. Hooker, gen. pl. III, 500 (1883) implied	—	—	—	—	N.S.W.	—	—	—						
D. Bairdianum; Bailey, suppl. to syn. of Q. Fl. Nov. (1884)	—	—	—	—	—	Q.	—	—						
D. Stuartii; Bailey in Proc. Roy. Soc. of Q. I, pt. 2 (1884)	—	—	—	—	—	Q.	—	—						
D. Kingianum; Bidwill in Bot. Regist. XXV, Misc. 11 (1839)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,280	M.fr.III,60.					
D. brachypus; G. Reichenbach in Garcke, Linnaea XLI, 42 (1876)	—	—	—	—	N.S.W.	—	—	—						
D. gracilicaule; F. v. M., fragm. I, 179 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,281	M.fr.III,164; IV,171.					
D. Moorei; F. v. M., fragm. VII, 29 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,281	M.fr.XI,54.					
D. antennatum; Lindley in Hook. Lond. Jour. of Bot. II, 236 (1843)	—	—	—	—	—	Q.	—	—						
D. agrostophyllum; F. v. M., fragm. VIII, 28 (1871)	—	—	—	—	—	Q.	—	B.fl.VI,282	M.fr.VIII,28.					
D. Baileyi; F. v. M., fragm. VIII, 173 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,173.					
D. Smilliae; F. v. M., fragm. VI, 94 (1868)	—	—	—	—	—	Q.	—	B.fl.VI,282	M.fr.VII,63; XI,87.					
D. ophioglossum; G. Reichenbach in Journ. L. Soc. XV, 113 (1876)	—	—	—	—	—	Q.	—	—	M.fr.XI,54.					
D. canaliculatum; R. Brown, prodr. 333 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,282	M.fr.III,126.					
D. Foelschei; F. v. M. in Wing's S. Sc. Record, Oct. (1882)	—	—	—	—	—	—	N.A.	—						
D. monophyllum; F. v. M., fragm. I, 189 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,283	M.fr.III,164; VII,64.					
D. hispidum; A. Richard, sert. Astrol. 13, t. 5 (1833)	—	—	—	—	—	Q.	—	B.fl.VI,283	M.fr.VII,30; VIII,150.					
D. cucumerinum; MacLeay in Bot. Reg., new ser. V, Misc. 58 (1842)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,283	M.fr.III,59; VIII,151.					
D. pugioniforme; Cunningham in Bot. Reg. XXV, Misc. 33 (1839)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,284	M.fr.I,189; III,59.					
D. rigidum; R. Brown, prodr. 333 (1810)	—	—	—	—	—	Q.	—	B.fl.VI,284	M.fr.VIII,248; XI,87.					
D. linguiforme; Swartz in Svensk. Vetensk. Ac. Handl. 247 (1800)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,284	M.fr.I,214.					
D. teretifolium; R. Brown, prodr. 333 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,285	M.fr.I,89; III,60.					
D. Fairfaxii; Fitzgerald & F. v. M. in Sydney Mail 360, Nov. (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,285	M.fr.X,122.					
D. striolatum; G. Reichenbach in Hamburg Gartenz. 313 (1857)...	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,285	M.fr.I,88; III,39; VII,30.					
D. Mortii; F. v. M., fragm. I, 214 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,286	M.fr.II,93; III,60; IX,50.					
D. Beckleri; F. v. M., fragm. V, 95 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,296	M.fr.VII,59.					

BULBOPHYLLUM; Petit-Thouars, orchid. Afric. t. 95 et 108 (1822). (Bolbophyllum, Thelychiton partly, Cirrhopetalum.)

B. Lichenastrum; F. v. M., fragm. VII, 60 (1869)	—	—	—	—	—	Q.	—	B.fl.VI,287	M.fr.X,119.					
B. Baileyi; F. v. M., fragm. IX, 5 (1875)	—	—	—	—	—	Q.	—	—	M.fr.IX,5.					
B. nematopodum; F. v. M., fragm. VIII, 30 (1873)...	—	—	—	—	—	Q.	—	B.fl.VI,287	M.fr.VIII,130.					
B. Shepherdi; F. v. M., fragm. III, 40 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,288	M.fr.I,190.					
B. aurantiacum; F. v. M., fragm. III, 39 (1862)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,288	M.fr.VII,98.					
B. argyropus; G. Reichenbach in Garcke, Linnaea XLI, 42 (1876)	—	—	—	—	N.S.W.	—	—	—						
B. exiguum; F. v. M., fragm. II, 72 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,288	M.fr.II,175; XI,22.					

B. minutissimum; F. v. M., fragm. XI, 53 (1878)	—	—	—	—	N.S.W.	Q.	—	—	M.fr.III,60; V,95; XI,88.							
B. Prenticei; F. v. M. in Wing's South. Sc. Record I, 173 (1881)	—	—	—	—	—	Q.	—	—	B.fl.X,139.							
B. Elisae; F. v. M., fragm. VI, 120 (1868)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,289	M.fr.VI,254; VIII,28.							
B. punctatum; Fitzgerald in Britten's Journ. Bot. XXI, 205 (1883)	—	—	—	—	—	Q.	—	—								
B. clavigerum; Cirrhopetalum; Fitzgerald in Brit. Journ. Bot. XXI,204 (1883)	—	—	—	—	—	Q.	—	—								
B. purpurascens; Bailey in Proc. Roy. Soc. Queensl. I, 88 (1884)	—	—	—	—	—	Q.	—	—								
B. Bowkettae; Bailey in Proc. Roy. Soc. Queensl. I, 88 (1884)	—	—	—	—	—	Q.	—	—								
ERIA; Lindley in Edw. Bot. Regist. t. 904 (1825).																
E. Fitzalani; F. v. M. in Wing's South. Sc. Rec. II, 251 (1882)	—	—	—	—	—	Q.	—	—								
LUISIA; Gaudichaud in Freyc. voy. bot. 426, t. 37 (1826).																
L. teretifolia; Gaudichaud in Freyc. voy. bot. 426, t. 37 (1826)	—	—	—	—	—	Q.	—	—	... ...							P.
SARCOCHILUS; R. Brown, prodr. 332 (1810). (Thrixspermum, Gunnia, Cleisostoma, Saccolabium partly.)																
S. erectus; F. v. M., Cens. 111 (1882) from Fitzgerald (1877)	—	—	—	—	N.S.W.	—	—	—	M.fr.XI,52,139.							
S. tridentatus; G. Reichenbach in Walp. Annal. VI, 500 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,296	M.fr.I,192; II,181; VII,98.							
S. Armitii; F. v. M., fragm. IX, 49 (1875)	—	—	—	—	—	Q.	—	—	M.fr.XI,49.							
S. Beckleri; F. v. M., Cens. of Austr. Pl. 111 (1882)	—	—	—	—	N.S.W.	—	—	B.fl.VI,296	M.fr.VIII,248.							
S. brevilabris; F. v. M., Cens. of Austr. Pl. 111 (1882)	—	—	—	—	—	Q.	—	—	M.fr.VIII,248; XI,139.							
S. Macphersoni; F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	—	Q.	—	B.fl.VI,297	M.fr.VII,96.							
S. Hartmanni; F. v. M., fragm. VIII, 248 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,248.							
S. divitiflorus; F. v. M. in Benth. Fl. Austr. VI, 292 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,292	M.fr.IX,50; XI,22.							
S. falcatus; R. Brown, prodr. 332 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.I,89,243; VII,87; X,50.							
S. Fitzgeraldi; F. v. M., fragm. VII, 115 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.VII,115.							
S. olivaceus; Lindley, Bot. Regist. Misc. XXV, 32 (1839)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,293	M.fr.I,90,191; VII,97.							
S. parviflorus; Lindley, Bot. Regist. Misc. XXIV, 34 (1838)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,294	M.fr.I,89,243; VII,98.							
S. Ceciliae; F. v. M., fragm. V, 42, t. 42 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,294	M.fr.VII,98; XI,22.							
S. Hillii; F. v. M., fragm. II, 94 (1861)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,295	M.fr.I,88,243; III,39; VII,98.							
S. phyllorrhizus; F. v. M., fragm. V, 201 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,295	M.fr.VII,30.							
S. Keffordii; F. v. M., Cens. sec. suppl. (1885) from Bailey (1884)	—	—	—	—	—	Q.	—	—								
S. platystachys; Bailey, syn. Q. Fl., first suppl. 56 (1886)...	—	—	—	—	—	Q.	—	—								
TAENIOPHYLLUM; Blume, Bijdr. 355 (1825).																
T. Muelleri; Lindley in Benth. Fl. Austr. VI, 291 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,291								
ORNITHOCHILUS; Wallich in Lindley, gen and spec. of orchid. plants 242 (1833). (Saccolabium partly.)																
O. Hillii; Bentham in Journ. Linn. Soc. XVIII, 334 (1880)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,298	M.fr.VII,97.							
GEODORUM; Jackson in Andrews, Bot. Reposit. t. 626 (1810).																
G. pictum; Lindley, gen. and spec. of orchid. pl. 175 (1833)	—	—	—	—	—	Q.	N.A.	B.fl.VI,299	M.fr.III,24; V,102; VIII,31.							As., P.
EULOPHIA; R. Brown in Edwards, Bot. Regist. 686 (1822). (Eulophus 1822.)																
E. venosa; G. Reichenbach in Benth. Fl. Austr. VI, 300 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.VI,300	M.fr.I,61.							
E. Fitzalani; F. v. M., fragm. VIII, 30 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,300	M.fr.VII,30.							
E. Holtzei; F. v. M. in Vict. Nat. VI, 55 (1889)	—	—	—	—	—	—	N.A.	—								
DIPODIUM; R. Brown, prodr. 336 (1810). (Wailesia.)																
D. punctatum; R. Brown, prodr. 331 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VI,301	M.fr.X,64.							
D. ensifolium; F. v. M., fragm. V, 42 (1865)	—	—	—	—	—	Q.	—	B.fl.VI,301	M.fr.V,42.							
CYMBIDIUM; Swartz in nov. act. Upsal. VI, 70 (1799).																
C. Hillii; F. v. M. in Regel's Gartenflora, 138 (1879)	—	—	—	—	—	Q.	—	—	M.fr.XI,88.							
C. canaliculatum; R. Brown, prodr. 331 (1810)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VI,202	M.fr.XI,88.							

C. albuciflorum; F. v. M., fragm. I, 188 (1859)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,303	M.fr.I,188							
C. suave; R. Brown, prodr. 331 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,303	M.fr.I,187.							
SPATHOGLOTTIS; Blume, Bijdr. 400 (1825).																
S. Paulinae; F. v. M., fragm. VI, 95 (1868)	—	—	—	—	—	Q.	—	B.fl.VI,304	M.fr.X,64.							
PHAJUS; Loureiro, Fl. Cochinch. II, 529 (1790).																
P. grandifolius; Loureiro, Fl. Cochinch. II, 529 (1790)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,304	M.fr.I,42; IV,163; X,6	...	As., P.					
CALANTHE; R. Brown in Edwards, Bot. Regist. 573 (1821).																
C. veratrifolia; R. Brown in Bot. Regist. 573 (1821)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,305	M.fr.VIII,30; IX,50	...	As., P.					
DIDYMOPLEXIS; Griffith in Calcutta Journ. of Nat. Hist. IV, 383 (1844). (Leucorchis.)																
D. pallens; Griffith in Calc. Journ. IV, 383, t. 17 (1844)	—	—	—	—	—	—	N.A.	—	...	...	...	...	As.			
GALEOLA; Loureiro, Fl. Cochinch. II, 520 (1790). (Erythrorchis, Ledgeria.)																
G. cassythoides; G. Reichenbach, Xen. Orchid. II, 77 (1869)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,307	M.fr.II,167; V,102; VIII,31; XI,105.							
G. Ledgeriana; F. v. M., fragm. XI, 127 (1872)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,307	M.fr.II,167; III,40; V,102; VIII,31.							
EPIPOGUM; Gmelin, Fl. Sibir. I, 11, t. 2 (1747).																
E. nutans; Lindley in Journ. Linn. Soc. I, 177 (1856)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,308	M.fr.VIII,30; IX,50; XI,105.	As., Afr.						
GASTRODIA; R. Brown, prodr. 330 (1810).																
G. sesamoides; R. Brown, prodr. 330 (1810)	W.A.	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,309	M.fr.VIII,151; X,64.							
POGONIA; A. L. de Jussieu, gen. plant. 65 (1789).																
P. uniflora; F. v. M., fragm. V, 201 (1866)	—	—	—	—	—	Q.	—	B.fl.VI,310	M.fr.V,201.							
P. holochila; F. v. M., fragm. V, 200 (1866)...	—	—	—	—	—	Q.	—	B.fl.VI,310	M.fr.V,200.							
P. flabelliformis; Lindley in Wallich's numer. list. 7400 (1828)	—	—	—	—	—	—	N.A.	—	...	...	...	As.				
P. Dallachyana; F. v. M. in Benth. Fl. Austr. VI, 310 (1870)	—	—	—	—	—	Q.	—	B.fl.VI,310								
P. pachystomoides; F. v. M., fragm. VIII, 174 (1874)	—	—	—	—	—	Q.	—	—	M.fr.VIII,174.							
CORYMBORCHIS; Petit-Thouars, hist. des plant. Orchid. recueill. sur les isles d'Afriq. t. 37-38 (1822). (Corymbis.)																
C. veratrifolia; Blume, Collect. des Orchidées 125, t. 42 & 43 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,311	M.fr.VIII,30; X,64	...	As., P.					
ETAERIA; Blume, Bijdr. 409 (1825). (Hetaeria, Rhamphidia.)																
E. tenuis; Bentham in Journ. Linn. Soc. XVIII, 345 (1880)	—	—	—	—	—	Q.	—	B.fl.VI,312	...	...	...	As., P.				
MICROSTYLIS; Nuttall, Genera of North American plants II, 196 (1818).																
M. congesta; G. Reichenbach in Walp. Annal. bot. syst. VI, 206 (1861) ...	—	—	—	—	—	Q.	—	—	M.fr.XI,21.	...	..	As.				
GOODYERA; R. Brown in Aiton. hort. Kew sec. ed. V, 197 (1813).																
G. viridiflora; Blume, Coll. des. Orchid. 41, t. 9 (1858)	—	—	—	—	—	Q.	—	B.fl.VI,313	M.fr.VIII,29.	..	..	As.				
G. polygonoides; F. v. M., fragm. VIII, 29 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,313	M.fr.VII,30; VIII,29; X,64.							
SPIRANTHES; L. C. Richard in Mém. du Mus. IV, 40 (1818). (Gyrostachis, 1807.)																
S. Australis; Lindley, Bot. Regist. 823 (1824)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,314	M.fr.X,65	...	...	E., As., N.Z.				
ANOECTOCHILUS; Blume, flor. Jav. praefat. 6 (1828).																
A. species unascertained	—	—	—	—	—	Q.	—	—								
THELYMITRA; R. & G. Forster, char. gen. 97, t. 49 (1776). (Macdonaldia.)																
T. ixioides; Swartz in Kong. Svensk. Acad. Handl. 228, t. 3 (1800)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,317								
T. circumsepta; Fitzgerald, Austral. Orchids I, part 4 (1877)	—	—	—	—	N.S.W.	—	—	—								
T. crinita; Lindley, Bot. Regist. XXV, App. 49 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,319	M.fr.VI,84.							
T. aristata; Lindley, gen. and spec. of orchid. pl. 521 (1840)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,319								
T. epipactoides; F. v. M., fragm. V, 174 (1866)	—	—	—	V.	—	—	—	—	M.fr.V,174.							

T. longifolia; R. & G. Forster, char. gen. 98, t. 49 (1776)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,319	... ...	P., N.Z.
T. villosa; Lindley, Bot. Regist. XXV, App. 49 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,320	M.fr.V,94,214.	
T. tigrina; R. Brown, prodr. 315 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,320		
T. fusco-lutea; R. Brown, prodr. 315 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,321	M.fr.V,94.	
T. stellata; Lindley, Bot. Regist. XXV, App. 49 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,321		
T. carnea; R. Brown, prodr. 314 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,321	M.fr.V,98.	
T. flexuosa; Endlicher, nov. stirp. decad. 23 (1839)...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,322		
T. antennifera; J. Hooker, Fl. Tasman. II, 4, t. 101 (1860)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,322		
T. Macmillani; F. v. M., frsgm. V, 93 (1865)	—	—	—	V.	—	—	—	B.fl.VI,323	M.fr.V,93.	
T. Mackibbinii; F. v. M. in Melb. Chemist 44 (1881)	W.A.	S.A.	—	V.	—	—	—	—	M.fr.XI,139.	
T. variegata; Lindley in Benth. Fl. Austr. VI, 323 (1870)...	W.A.	—	—	—	—	—	—	B.fl.VI,323	M.fr.V,97,98.	
T. venosa; R. Brown, prodr. 314 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,323	... ...	N.Z.
T. cyanea; Lindley in Benth. Fl. Austr. VI, 323 (1870)	—	—	T.	—	—	—	—	B.fl.VI,323		
T. mucida; Fitzgerald in Gard. Chron. new ser. XVII, 495 (1882)	W.A.	—	—	—	—	—	—	—		
T. luteo-ciliata; Fitzgerald in Gard. Chron. new ser. XVII, 433 (1882)	—	S.A.	—	—	—	—	—	—		
EPIBLEMA; R. Brown, prodr. fl. Nov. Holl. 315 (1810).										
E. grandiflorum; R. Brown, prodr. 315 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,324	M.fr.XI,21.	
DIURIS; Smith in Transact. Linn. Soc. IV, 222 (1798).										
D. alba; R. Brown, prodr. 316 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,325	M.fr.V,102.	
D. punctata; Smith, Exot. Bot. I, 13, t. 8 (1804)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,326	M.fr.V,102.	
D. secundiflora; Fitzgerald, Austral. Orchids I, part 4 (1877)	—	—	—	—	N.S.W.	Q.	—	—		
D. aurea; Smith, Exot. Bot. 15, t. 9 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,327	M.fr.V,172.	
D. palustris; Lindley, gen. and spec. of orchid. pl. 507 (1840)	—	S.A.	T.	V.	—	—	—	B.fl.VI,327	M.fr.V,102.	
D. maculata; Smith, Exot. Bot. I, 57, t. 30 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,327	M.fr.V,102.	
D. pedunculata; R. Brown, prodr. 316 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,328	M.fr.V,102,173.	
D. pallens; Benth. Fl. Austr. VI, 329 (1870)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,329		
D. abbreviata; F. v. M. in Benth. Fl. Austr. VI, 329 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,329		
D. setacea; R. Brown, prodr. 316 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,329	M.fr.V,101.	
D. emarginata; R. Brown, prodr. 316 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VI,330		
D. sulphurea; R. Brown, prodr. 316 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,330	M.fr.V,102,173.	
D. aequalis; F. v. M. in Fitzgerald, Austral. Orchids I, part 2 (1876)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,65.	
D. dendrobioides; Fitzgerald, Austral. Orchids I, part 7 (1882)	—	—	—	—	N.S.W.	—	—	—		
D. longifolia; R. Brown, prodr. 316 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,331	M.fr.V,101,172.	
D. pauciflora; R. Brown, prodr. 316 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,331		
ORTHOCERAS; R. Brown, prodr. fl. Nov. Holl. 317 (1810).										
O. strictum; R. Brown, prodr. 317 (1810)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,332	... ...	N.Z.
CALOCHILUS; R. Brown, prodr. fl. Nov. Holl. 320 (1810).										
C. campester; R. Brown, prodr. 320 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,315	M.fr.VIII,151.	
C. Robertsoni; Bentham, Fl. Austr. VI, 315 (1870)..	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,315		
C. paludosus; R. Brown, prodr. 320 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,316		
CRYPTOSTYLIS; R. Brown, prodr. 317 (1810). (Zosterostylis.)										
C. longifolia; R. Brown, prodr. 317 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,333	M.fr.VII,116; XI,21.	
C. ovata; R. Brown, prodr. 317 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,334	M.fr.VII,116.	
C. erecta; R. Brown, prodr. 317 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,334	M.fr.VII,115.	
C. leptochila; F. v. M. in Benth. Fl. Austr. VI, 324 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,334		
PRASOPHYLLUM; R. Brown, prodr. 317 (1810). (Genoplesium, Carunastylis).										
P. Australe; R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,337		
P. flavum; R. Brown, prodr. 318 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,337		

P. elatum; R. Brown, prodr. 318 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,337	M.fr.V,100.		
P. brevilabre; J. Hooker, Fl. Tasman. II, 11, t. 110 (1860)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,338	M.fr.V,101.		
P. hians; G. Reichenbach, Beitr. zur syst. Pflanzenk. 61 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,338			
P. patens; R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,339	M.fr.V,101.		
P. Deaneanum; Fitzgerald in proc. L. S. of N.S.W. sec. ser. II, 142 (1887) ...	—	—	—	—	N.S.W.	—	—	—			
P. fuscum; R. Brown, prodr. 318 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,339	M.fr.V,101.		
P. cyphochilum; Bentham, Fl. Austr. VI, 340 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,340			
P. ovale; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,341			
P. macrostachyum; R. Brown, prodr. 318 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,341			
P. Reichenbachii; F. v. M., Fimbria; G. Reichenbach, Beitr. 60 (1871) ...	W.A.	—	—	V.	—	—	—	B.fl.VI,341			
P. striatum; R. Brown, prodr. 318 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,342	M.fr.V,100.		
P. parvifolium; Lindley, Bot. Regist. XXV, App. 54 (1939)	W.A.	—	—	—	—	—	—	B.fl.VI,342	M.fr.V,101.		
P. gibbosum; R. Brown, prodr. 381 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,342	M.fr.V,100.		
P. cucullatum; G. Reichenbach, Beitr. zur syst. Pflanzenk. 59 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,343			
P. nigricans; R. Brown, prodr. 319 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,343	M.fr.IX,50.		
P. rufum; R. Brown, prodr. 319 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,344	M.fr.V,100	...	N.Z.
P. brachystachyum; Lindley, gen. and spec. of orchid. pl. 513 (1840)	—	—	T.	—	—	—	—	B.fl.VI,344	M.fr.V,100.		
P. despectans; J. Hooker, Fl. Tasman. II, 13, t. 113 (1860)	—	S.A.	T.	V.	—	—	—	B.fl.VI,345	M.fr.V,100.		
P. fimbriatum; R. Brown, prodr. 319 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VI,345	M.fr.V,100.		
P. Archeri; J. Hooker, Fl. Tasman. II, 14, t. 113 (1860)	—	—	T.	V.	—	—	—	B.fl.VI,345	M.fr.V,100.		
P. intricatum; C. Stuart in Benth. Fl. Austr. VI, 346 (1870)	—	—	T.	V.	—	—	—	B.fl.VI,346			
P. Woollsii; F. v. M., fragm. V, 100 (1866)	—	—	—	—	N.S.W.	—	—	B.fl.VI,346	M.fr.V,100.		
P. apostasioides; Carunastylis, Fitzgerald, Austr. Orchids II, part 3 (1888) ...	—	—	—	—	N.S.W.	—	—	—			
MICROTIS; R. Brown, prodr. fl. Nov. Holl. 320 (1810).											
M. porrifolia; R. Brown, prodr. 320 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,347	M.fr.X,65; XI,21	...	N.Z., P.
M. media; R. Brown, prodr. 321 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,348			
M. alba; R. Brown, prodr. 321 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VI,348			
M. pulchella; R. Brown, prodr. 321 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,349			
M. atrata; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,349	M.fr.I,90,244; X,65.		
CORYSANTHES; R. Brown, prodr. 328 (1810). (Corybas, Nematoceras.)											
C. unguiculata; R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,350			
C. pruinosa; R. Cunningham in N.S. Wales Magaz. n. 1 (1833)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	—	M.fr.IX,49.		
C. fimbriata; R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,351	M.fr.X,65.		
C. bicalcarata; R. Brown, prodr. 328 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,351			
PTEROSTYLIS; R. Brown, prodr. fl. Nov. Holl. 326 (1810).											
P. ophioglossa; R. Brown, prodr. 326 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,354			
P. concinna; R. Brown, prodr. 326 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VI,355	M.fr.V,99; XI,105.		
P. curta; R. Brown, prodr. 326 (1810)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,355	M.fr.V,99.		
P. acuminata; R. Brown, prodr. 326 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,355			
P. nutans; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,356	M.fr.V,99.		
P. pedaloglossa; Fitzgerald, Austral. Orchids II, part 3 (1887)	—	—	—	V.	N.S.W.	—	—	—			
P. pedunculata; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,356	M.fr.VIII,249; XI,105.		
P. nana; R. Brown, prodr. 327 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,357	M.fr.V,99.		
P. cucullata; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,357	M.fr.V,99.		
P. grandiflora; R. Brown, prodr. 327 (1810)	—	—	—	V.	N.S.W	Q.	—	B.fl.VI,358	M.fr.V,99;XI,105.		
P. truncata; Fitzgerald, Austral. Orchids I, part 4 (1878)	—	—	—	—	N.S.W.	—	—	—			
P. reflexa; R. Brown, prodr. 327 (1810)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VI,359	M.fr.XI,105.		
P. praecox; Lindley, gen. and spec. of Orchid. pl. 388 (1840)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,359			

P. obtusa; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,360	M.fr.IX,50; XI,53.	
P. recurva; Bentham, Fl. Austr. VI, 360 (1870)	W.A.	—	—	—	—	—	—	B.fl.VI,360		
P. parviflora; R. Brown, prodr. 327 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,361	M.fr.V,100.	
P. barbata; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,362	M.fr.VIII,151; XI,22,105.	N.Z.
P. mutica; R. Brown, prodr. 328 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,362	M.fr.V,100.	
P. rufa; R. Brown, prodr. 327 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,363	M.fr.V,100.	
P. Daintreyana; F. v. M. in Benth. Fl. Austr. VI, 360 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VI,360		
P. longifolia; R. Brown, prodr. 327 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,364	M.fr.V,100.	
P. vittata; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,364	M.fr.XI,126.	
CALEYA; R. Brown in Aiton, Hort. Kew sec. ed. V, 204 (1813). (Caleana 1810).										
C. major; R. Brown, prodr. 329 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,366	M.fr.XI,105.	
C. minor; R. Brown, prodr. 329 (1810)	—	—	T.	—	N.S.W.	—	—	B.fl.VI,366		
C. nigrita; Lindley, Bot. Regist. XXV, App. 54 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,366		
C. Sullivani; F. v. M. in Melb. Chemist 44 (1882)	—	—	—	V.	—	—	—	—		
DRAKAEA; Lindley, Bot. Regist. XXV, App. 55 (1834). (Spiculaea, Arthrochilus.)										
D. ciliata; G. Reichenbach, Beitr. zur. syst. Pflanzenk. 68 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,367		
D. irritabilis; G. Reichenbach, Beitr. zur syst. Pflanzenk. 68 (1871)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,368	M.fr.I,43; VIII,151; XI,53.	
D. elastica; Lindley, Bot. Regist. XXV, App. 55 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,368		
ACIANTHUS; R. Brown, prodr. fl. Nov. Holl. 321 (1810).										
A. caudatus; R. Brown, prodr. 321 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,369		
A. fornicatus; R. Brown, prodr. 321 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,370		
A. exsertus; R. Brown, prodr. 321 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,370	M.fr.XI,53,105.	
A. viridis; J. Hooker, Fl. Tasman. II, 372 (1860)	—	—	T.	—	—	—	—	B.fl.IV,371	M.fr.IX,50.	
CYRTOSTYLIS; R. Brown, prodr. fl. Nov. Holl. 322 (1810).										
C. reniformis; R. Brown, prodr. 322 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,376		
LYPERANTHUS; R. Brown, prodr. 325 (1810). (Burnettia, Fitzgeraldia.)										
L. nigricans; R. Brown, prodr. 325 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,374	M.fr.VII,134; X,65.	
L. Burnettii; F. v. M., fragm. V, 96 (1865)	—	—	T.	V.	—	—	—	B.fl.VI,375	M.fr.V,96; VII,134.	
L. Forrestii; F. v. M. in Wing's S. Sc. Rec. II, 55 (1882)	W.A.	—	—	—	—	—	—	—		
L. ellipticus; R. Brown, prodr. 325 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VI,374	M.fr.VII,133.	
ERIOCHILUS; R. Brown, prodr. fl. Nov. Holl. 323 (1810).										
E. autumnalis; R. Brown, prodr. 323 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,372	M.fr.XI,22,105.	
E. scaber; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,372		
E. tenuis; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,372		
E. dilatatus; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,372		
E. multiflorus; Lindley, Bot. Regist. XXV, App. 53 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,373	M.fr.XI,105.	
E. fimbriatus; F. v. M. in Wing's S. Sc. Record II, 152 (1882)	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,377	M.fr.XI,104.	
CALADENIA; R. Brown, prodr. 325 (1810). (Leptoceras, Adenochilus.)										
C. Menziesii; R. Brown, prodr. 325 (1810)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VI,377	M.fr.V,101; XI,105.	
C. Nortoni; F. v. M., Cens. 113 (1882) from Fitzgerald (1876)	—	—	—	—	N.S.W.	—	—	—	M.fr.X,64.	
C. Cairnsiana; F. v. M., fragm. VII, 31 (1869)	W.A.	S.A.	—	—	—	—	—	B.fl.VI,378	M.fr.VII,31.	
C. multiclavia; G. Reichenbach, Beitr. zur syst. Pflanzenk. 64 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,378		
C. discoidea; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,382		
C. Patersoni; R. Brown, prodr. 324 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,382	M.fr.V,101.	
C. Drummondii; Bentham, Fl. Austr. VI, 383 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,383		
C. hirta; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,383		

C. Roei; Bentham, Fl. Austr. VI, 383 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,383	
C. Barbarossae; G. Reichenbach, Beitr. zur syst. Pflanzenk. 64 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,383	
C. flava; R. Brown, prodr. 324 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,384	M.fr.V,101; VI,83; XI,105.
C. latifolia; R. Brown, prodr. 324 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,384	M.fr.V,101; XI,105.
C. reptans; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,385	
C. suaveolens; G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,385	M.fr.V,93.
C. serrata; G. Reichenbach, Beitr. zur syst. Pflanzenk. 67 (1871)	W.A.	—	—	—	—	—	—	B.fl.VI,386	M.fr.V,99.
C. carnea; R. Brown, prodr. 324 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,386	M.fr.V,101.
C. congesta; R. Brown, prodr. 324 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,387	
C. aphylla; Bentham, Fl. Austr. VI, 387 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,387	
C. coerulea; R. Brown, prodr. 324 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,388	M.fr.V,101; XI,22.
C. deformis; R. Brown, prodr. 324 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,388	M.fr.V,101; XI,105.
C. sericea; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,389	
C. gemmata; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,389	M.fr.V,101.
C. ixioides; Lindley, Bot. Regist. XXV, App. 52 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,389	
C. unita; Fitzgerald in Gardeners' Chronicle 461 (1882)	W.A.	—	—	—	—	—	—	—	
CHILOGLOTTIS; R. Brown. prodr. 322 (1810).									
C. diphylla; R. Brown, prodr. 323 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VI,390	M.fr.X,116.
C. Gunnii; Lindley, gen and spec. orchid. 387 (1840)	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,391	M.fr.VIII,151; IX,50; X,116; XI,22.
C. trilabra; Fitzgerald in Britten's Journ. of Bot. XXI, 304 (1883)	—	—	—	—	—	Q.	—	—	
GLOSSODIA; R. Brown, prodr. 325 (1810).									
G. major; R. Brown, prodr. 326 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,392	
G. minor; R. Brown, prodr. 326 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,392	
G. Brunonis; Endlicher, nov. stirp. dec. 16 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,393	M.fr.VI,83.
G. intermedia; Fitzgerald in Gardeners' Chronicle 462 (1882)	W.A.	—	—	—	—	—	—	—	
G. emarginata; Lindley, gen. and spec. of orchid. pl. 424 (1840)	W.A.	—	—	—	—	—	—	B.fl.VI,393	M.fr.XI,22.
HABENARIA; Willdenow, spec. plant. IV, 44 (1805).									
H. trinervis; Wight, icon. plant. Ind. or. t. 1701 (1852)	—	—	—	—	—	—	N.A.	B.fl.VI,394	As.
H. elongata; R. Brown, prodr. 313 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,394	M.fr.VII,15.
H. graminea; Lindley, gen. and spec. of orchid. pl. 318 (1840)	—	—	—	—	—	Q.	N.A.	B.fl.IV,394	M.fr.VII,16.
H. ochroleuca; R. Brown, prodr. 313 (1810)	—	—	—	—	—	—	N.A.	B.fl.IV,394	M.fr.V,102.
H. xanthantha; F. v. M., fragm. VII, 16 (1869)	—	—	—	—	—	Q.	—	B.fl.VI,395	M.fr.VII,16.

APOSTASIACEAE.

Ach. Richard, in Ch. d'Orbigny Dictionn. univ. d'hist. nat. II, 34 (1842), from R. Brown (1830).

APOSTASIA; Blume, Bijdr. 423 (1825). (Niemeyera.)									
A. stylidioides; G. Reichenbach in Flora Regens. C. B. Z. 278 (1872)	—	—	—	—	—	Q.	—	B.fl.VI,396	M.fr.VI,96; VII,114.

BURMANNIACEAE.

Sprengel, syst. veget. I, 125 (1825).—Genus 1, Species 2.

BURMANNIA; Linné, syst. nat. 8 (1735); gen. plant. 100 (1737).									
B. disticha; Linné, spec. plant. 287 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VI,397	As.
B. juncea; Solander in R. Brown, prodr. 265 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VI,397	As.

IRIDEAE.

VENTENAT, Tableau du règne végétal II, 188 (1799).—Genera 4, Species 24.

DIPLARRHENA ; Labillardière, Relation du voy. à la rech. de La Per. I, 275, t. 15 (1799)

D. Moraea ; Labillardière, Voy. I, 157, t. 15 (1799) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,400	M.fr.VII,94.		

PATERSONIA ; R. Brown in Bot. Magaz. t. 1041 (1807). (Genosiris, 1804.)

P. glauca ; R. Brown, prodr. 304 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VI,402	M.fr.VII,36.		
P. longiscapa ; Sweet, Fl. Austr. t. 39 (1828)...	—	S.A.	T.	V.	—	—	—	B.fl.VI,402	M.fr.VII,36.		
P. occidentalis ; R. Brown, prodr. 304 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VI,403	M.fr.VII.31.		
P. umbrosa ; Endlicher in Lehm. pl. Preiss. II, 31 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.VII,32.		
P. xanthina ; F. v. M., fragm. I, 214 (1859) ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.I,214.		
P. limbata ; Endlicher in Lehm. pl. Preiss. II, 29 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.VII,32.		
P. juncea ; Lindley, Bot. Regist. XXV, App. 58 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,404	M.fr.VII,33.		
P. Maxwelli ; F. v. M. in Benth. Fl. Austr. VI, 405 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,405	M.fr.VII,34.		
P. pygmaea ; Lindley, Bot. Regist. XXV, App. 58 (1839) ...	W.A.	—	—	—	—	—	—	B.fl.VI,405	M.fr.VII,33.		
P. sericea ; R. Brown in Bot. Mag. t. 1041 (1807) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,406	M.fr.VII,35.		
P. lanata ; R. Brown, prodr. 303 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VI,406	M.fr.VII,35.		
P. rudis ; Endlicher in Lehm. pl. Preiss. II, 29 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VI,406	M.fr.VII,35.		
P. macrantha ; Bentham, Fl. Austr. VI, 407 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,407			
P. glabrata ; R. Brown, prodr. 304 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,407	M.fr.VII,35.		
P. Drummondii ; F. v. M. in Benth. Fl. Austr. VI, 407 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,407			
P. inaequalis ; Bentham, Fl. Austr. VI, 408 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,408			
P. graminea ; Bentham, Fl. Austr. VI, 408 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,408	M.fr.XI,23.		
P. babianoides ; Bentham, Fl. Austr. VI, 408 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VI,408			

IRIS ; Tournefort, inst. 358, t. 186, 188 (1700), from Hippocrates, Dioscorides and Plinius. (Moraea partly.)

I. Robinsoniana ; F. v. M., fragm. VII, 153 (1870) ...	—	—	—	—	N.S.W.	—	—	B.fl.VI,409	M.fr.VII,153; VIII,281.		

SISYRINCHIUM ; Linné, gen. plant. 273 (1737). (Libertia, Orthrosanthus, Renealmia, Nematostigma.)

S. paniculatum ; R. Brown, prodr. 305 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VI,413	M.fr.VII,91.		
S. pulchellum ; R. Brown, prodr. 305 (1810) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VI,413	M.fr.VII,92 ...	...	P., N.Z.
S. cyaneum ; Lindley, Bot. Regist. t. 1090 (1827) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.VI,410	M.fr.VII,92.		
S. polystachyum ; F. v. M., Orthrosanthus, Bentham, Fl. Austr. VI, 411 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,411			

HYDROCHARIDEAE.

LAMARCK & DE CANDOLLE, Flore Française, III, 265 (1805).—Genera 7, Species 9.

HYDROCHARIS ; Linné, syst. nat. 8 (1735); gen. plant. 308 (1737).

H. morsus ranae ; Linné, spec. plant. 1036 (1753) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VI,256	M.fr.VI,199 ...	E., As.	

ENHALUS ; L. C. Richard in Mém. de l'Inst. II, 64 (1811).

E. Koenigii ; Richard in Mém. de l'Inst. II, 64 (1811) ...	—	—	—	—	—	Q.	—	—	M.fr.VIII,219 ...	As.	

HALOPHILA ; Du Petit-Thouars, gen. nov. Madag. n. 6 (1806).

H. ovata ; Gaudichaud, Bot. Voy. Freyc. 470, t. 40 (1826)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,182	M.fr.VIII,219 ...	As.	
H. spinulosa ; Ascherson in Neumayer, Anleit. zu wiss. Beob. 368 (1875) ...	—	—	—	—	—	Q.	—	B.fl.VII,183	M.fr.VIII,219,283.		

OTTELIA ; Persoon, synops. plant. I, 400 (1805). (Damasonium partly, Hymenotheca.)

O. alismoides ; Persoon, synops. plant. I, 400 (1805) ...	—	—	—	—	—	Q.	N.A.	B.fl.VI,257	M.fr.I,94; VI,199.	As.	
O. ovalifolia ; L. C. Richard in Mém. de l'Inst. II, 78 (1811) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VI,257			

BLYXA; Noronha in Du Petit-Thouars, gen. nov. Madag. n. 14 (1806).

B. Roxburghii; L. C. Richard, Mém. de l'Inst. II, 77, t. 5 (1811)	—	S.A.	—	—	—	Q.	N.A.	B.fl. VI,258		...	...	As.

VALLISNERIA; Micheli, nov. plant. gen. 12, t. 10 (1729).

V. spiralis; Linné, spec. plant. 1015 (1753)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl. VI,259		...	..	E., As., Afr., Am.

HYDRILLA; L. C. Richard in Mém. de l'Inst. II, 61 (1811). (Elodeae subgenus.)

E. verticillata; F. v. M., Key to the System of Vict. plants I, 423 (1888)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VI,259	M.fr. I,94	...		As., Afr.

TACCACEAE.

Presl, reliq. Haenk. I, 149 (1828).—Genus 1, Species 1.

TACCA, R. et G. Forster, char. gen. 65 (1776).

T. pinnatifida; R. et G. Forster, char. gen. 65 (1776)	—	—	—	—	—	Q.	N.A.	B.fl. VI,453		...	...	As., P.

HAEMODORACEAE.

R. Brown, prodr. Fl. Nov. Holl. 299 (1810).—Genera 5, Species 66.

HAEMODORUM, Smith in Transact. Linn. Soc. 213 (1798).

H. distichophyllum; Hooker, icon. plant. t. 866 (1852)	—	—	T.	—	—	—	—	B.fl. VI,419	M.fr. VII,118; IX,51.			
H. brevicaule; F. v. M., fragm. I, 64 (1858)	—	—	—	—	—	—	N.A.	B.fl. VI,419	M.fr. VII,118.			
H. sparsiflorum; F. v. M., fragm. VII, 117 (1870)	W.A.	—	—	—	—	—	—	B.fl. VI,420	M.fr. VII,117.			
H. spicatum; R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl. VI,420	M.fr. VII,118.			
H. brevisepaleum; Bentham, Fl. Austr. VI, 420 (1873)	W.A.	—	—	—	—	—	—	B.fl. VI,420				
H. paniculatum; Lindley, Bot. Regist. XXV, App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,420				
H. laxum; R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl. VI,421	M.fr. VII,118.			
H. simplex; Lindley, Bot. Regist. XXV, App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,421				
H. simulans; F. v. M., fragm. VII, 117 (1870)	W.A.	—	—	—	—	—	—	B.fl. VI,421	M.fr. VII,117.			
H. planifolium; R. Brown, prodr. 300 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. VI,422	M.fr. VII,118.			
H. teretifolium; R. Brown, prodr. 300 (1810)	—	—	—	—	N.S.W.	—	—	B.fl. VI,422	M.fr. VII,117.			
H. coccineum; R. Brown, prodr. 300 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VI,422	M.fr. VII,117.			
H. subvirens; F. v. M., fragm. I, 63 (1858)	—	—	—	—	—	—	N.A.	B.fl. VI,423	M.fr. I,63.			
H. ensifolium; F. v. M., fragm. I, 64 (1858)	—	—	—	—	—	Q.	N.A.	B.fl. VI,423	M.fr. VII,118.			
H. parviflorum; Bentham, Fl. Austr. VI, 423 (1873)	—	—	—	—	—	—	N.A.	B.fl. VI,423				
H. leptostachyum; Bentham, Fl. Austr. VI, 423 (1873)	—	—	—	—	—	—	N.A.	B.fl. VI,423				
H. tenuifolium; Cunningham in Benth. Fl. Austr. VI, 425 (1873)	—	—	—	—	—	Q.	—	B.fl. VI,423				

PHLEBOCARYA; R. Brown, prodr. 301 (1810).

P. ciliata; R. Brown, prodr. 301 (1810)	W.A.	—	—	—	—	—	—	B.fl. VI,424	M.fr. VIII,23.			
P. pilosissima; F. v. M. in Benth. Fl. Austr. VI, 425 (1873)	W.A.	—	—	—	—	—	—	B.fl. VI,425	M.fr. VIII,23.			
P. filifolia; F. v. M. in Benth. Fl. Austr. VI, 425 (1873)	W.A.	—	—	—	—	—	—	B.fl. VI,425	M.fr. VIII,23.			

TRIBONANTHES; Endlicher, nov. stirp. dec. I, 27 (1839).

T. brachypetala; Lindley, Bot. Regist. XXV, App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,426	M.fr. VIII,23.			
T. uniflora; Lindley, Bot. Regist. XXV, App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,427	M.fr. VIII,23.			
T. Australis; Endlicher, nov. stirp. dec. I, 27 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,427	M.fr. VIII,23.			
T. variabilis; Lindley, Bot. Regist. XXV, App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,427				
T. longipetala; Lindley, Bot. Regist. XXV, App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl. VI,427	M.fr. VIII,23.			

CONOSTYLIS; R. Brown, prodr. 300 (1810). (Blancoa, Androstemma.)

C. breviscapa; R. Brown, prodr. 301 (1810)	W.A.	—	—	—	—	—	—	B.fl. VI,430				
C. vaginata; Endlicher in Lehm. pl. Preiss. II, 23 (1846)	W.A.	—	—	—	—	—	—	B.fl. VI,431	M.fr. VIII,19.			

C. petrophiloides; F. v. M. in Benth. Fl. Austr. VI, 431 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,431	
C. setosa; Lindley, Bot. Regist. XXV, App. 44, t. 6 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,431	M.fr.VIII,19.
C. aurea; Lindley, Bot. Regist. XXV. App. 44 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,19.
C. melanopogon; Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,20.
C. setigera; R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,432	M.fr.VIII,20.
C. Psyllium; Endlicher in Lehm. pl. Preiss. II, 21 (1846)...	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. villosa; Bentham, Fl. Austr. VI, 433 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. Drummondii; Bentham, Fl. Austr. VI, 433 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,433	
C. involucrata; Endlicher in Lehm. pl. Preiss. II, 23 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,434	
C. gladiata; Bentham, Fl. Austr. VI, 434 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,434	
C. seorsiflora; F. v. M., fragm. I, 158 (1859)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,19.
C. teretiuscula; F. v. M., fragm. VIII, 18 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,18.
C. stylidioides; F. v. M., fragm. VIII, 17 (1872)	W.A.	—	—	—	—	—	—	B.fl.VI,435	M.fr.VIII,17.
C. prolifera; Bentham, Fl. Austr. VI, 436 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. racemosa; Bentham, Fl. Austr. VI, 436 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. candicans; Endlicher, nov. stirp. dec. 20 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI.436	M.fr.VI,255.
C. dealbata; Lindley, Bot. Regist. XXV, App. 45 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,436	
C. Preissii; Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,437	M.fr.VIII,20.
C. bracteata; Endlicher in Lehm. pl. Preiss. II, 16 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,437	M.fr.VIII,20.
C. filifolia; F. v. M., fragm. VIII, 18 (1873)...	W.A.	—	—	—	—	—	—	B.fl.VI,438	M.fr.VIII,18.
C. spinuligera; F. v. M. in Benth. Fl. Austr. VI, 438 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,438	
C. bromelioides; Endlicher in Lehm. pl. Preiss. II, 18 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,438	
C. aculeata; R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,438	M.fr.VIII,19.
C. laxiflora; Bentham, Fl. Austr. VI, 439 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,439	
C. cymosa; F. v. M. in Benth. Fl. Austr. VI, 439 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,439	
C. serrulata; R. Brown, prodr. 300 (1810)	W.A.	—	—	—	—	—	—	B.fl.VI,439	M.fr.VIII,19.
C. caricina; Lindley, Bot. Regist. XXV, App. 45 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,440	M.fr.VIII,20.
C. Androstemma; F. v. M., fragm. VIII, 19 (1873)...	W.A.	—	—	—	—	—	—	B.fl.VI,440	M.fr.VIII,19; IX,51.
C. Bealiana; F. v. M., fragm. IX, 50 (1875)	W.A.	—	—	—	—	—	—	—	M.fr.IX,50.
C. canescens; F. v. M., fragm. VIII, 19 (1873)	W.A.	—	—	—	—	—	—	B.fl.VI,441	M.fr.VIII,19; IX,51.

ANIGOZANTHOS; Labillardière, Relat. du voy. à la rech. de La Perouse I, 411 (1798). (Anigosanthos, Macropidia, Macropodia, Schwaegrichenia.)

A. rufa; Labillardière, Voy. I, 411, t. 22 (1798)	W.A.	—	—	—	—	—	—	B.fl.VI,442	M.fr.VIII,21.
A. pulcherrima; Hooker, Bot. Mag. t. 4180 (1845)	W.A.	—	—	—	—	—	—	B.fl.VI,443	M.fr.VIII,21.
A. flavida; Redouté et De Candolle, les Liliacées t. 176 (1807)	W.A.	—	—	—	—	—	—	B.fl.VI,443	M.fr.VIII,21.
A. Preissii; Endlicher in Lehm. pl. Preiss. II, 26 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,444	M.fr.VIII,23.
A. humilis; Lindley, Bot. Regist. XXV, App. 46 (1839)	W.A.	—	—	—	—	—	—	B.fl.VI,444	M.fr.VIII,21.
A. viridis; Endlicher in Lehm. pl. Preiss. II, 25 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,445	M.fr.VIII,22.
A. Manglesii; D. Don in Sweet, Brit. Fl. Gard. sec. ser. t. 265 (1836)	W.A.	—	—	—	—	—	—	B.fl.VI,445	M.fr.VIII,22.
A. bicolor; Endlicher in Lehm. pl. Preiss. II, 26 (1846)	W.A.	—	—	—	—	—	—	B.fl.VI,446	M.fr.VIII,22.
A. fuliginosus; Hooker, Bot. Mag. t. 4291 (1847)	W.A.	—	—	—	—	—	—	B.fl.VI,447	M.fr.VIII,20.

AMARYLLIDEAE.

J. St. Hilaire, Exposit. des famill. natur. des plantes I, 134, t. 21 (1805).—Genera 7, Species 21.

CAMPYNEMA, Labillardière, Nov. Holl. plant. spec. I, 93 (1804).

C. lineare; Labillardière, Nov. Holl. pl. spec. I, 93, t. 121 (1804)	—	—	T.	—	—	—	—	B.fl.VI,415	M.fr.VI,96.

HYPOXIS; Linné, syst. ed. decim. 986 (1759).

H. hygrometrica; Labillardière, Nov. Holl. pl. spec. I, 82, t. 108 (1804) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,449	M.fr.VI,96.
H. glabella; R. Brown, prodr. 289 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VI,450	M.fr.VI,96.

| | | |

LILIACEAE.

HALLER, enum. stirp. Helv. I, 279 (1742).—Genera 45, Species 161.

SMILAX ; Tournefort, inst. 654, t. 421 (1700), from Theophrastos and Dioscorides.

S. glycyphylla ; Smith in White, Voy. to N.S.Wales 230 (1790) ...	...	...	—	—	—	—		N.S.W.	Q.	—	B.fl.VII,7	M.fr.VII,77.					
S. Australis ; R. Brown, prodr. 293 (1810) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,7	M.fr.VII,78.			
S. purpurata ; G. Forster, fl. ins. Austr. prodr. 70 (1786) ...	...	...	...	—	—	—	—		N.S.W.	—	—	—					

RHIPOGONUM ; R. & G. Forster, char. gen. 49, t. 25 (1776). (Ripogonum.)

R. album ; R. Brown, prodr. 293 (1810) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,9	M.fr.I,43,243; VII,79; VIII,280.			
R. discolor ; F. v. M., fragm. VII, 78 (1870)...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,9	M.fr.VII,78.			
R. Fawcettianum ; F. v. M. in Benth. Fl. Austr. VII, 9 (1878) ...	...	...	...	—	—	—	—		N.S.W.	Q.	—	B.fl.VII,9	M.fr.VII,80.				
R. Elseyanum ; F. v. M., fragm. I, 44 (1858) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,10	M.fr.VII,80.			

DRYMOPHILA ; R. Brown, prodr. fl. Nov. Holl. 292 (1810).

D. cyanocarpa ; R. Brown, prodr. 292 (1810)...	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,12	M.fr.VII,73; VIII,280; IX,191.			
D. Moorei ; Baker in Journ. Linn. Soc. XIV, 571 (1875) ...	...	...	...	—	—	—	—		N.S.W.	—	—	B.fl.VII,13	M.fr.IX,190.				

DIANELLA ; Lamarck, Encyclop. méthod. II, t. 250 (1786).

D. Tasmanica ; J. Hooker, Fl. Tasman. II, 57, t. 133 (1860) ...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,14	M.fr.VI,121; VIII,279.						
D. longifolia ; R. Brown, prodr. 280 (1810) ..	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,14	M.fr.VI,122.			
D. revoluta ; R. Brown, prodr. 280 (1810) ...	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,15	M.fr.VI,121; XI,20.			
D. caerulea ; Sims, Bot. Mag. t. 505 (1799) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,16	M.fr.VI,123.			
D. ensifolia ; De Candolle & Redouté, Liliac. t. 1 (1802) ...	...	...	...	—	—	—	—		—	Q.	N.A.	B.fl.VII,16	M.fr.VI,123	...	As., Afr., P.		

ASPARAGUS ; Tournefort, inst. 300, t. 154 (1700), from Theophrastos, Dioscorides, and Plinius. (Asparagopsis.)

A. racemosus ; Willdenow, spec. plant. II, 152 (1799) ...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,17	M.fr.VII,73	...	As., Afr.		

EUSTREPHUS ; R. Brown, prodr. 281 (1810).

E. Brownii ; F. v. M., fragm. VII, 73 (1870)...	...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,18	M.fr.VII,73; VIII,280.	P.		

GEITONOPLESIUM ; Cunningham in Bot. Magaz. t. 3131 (1832). (Luzuriago partly, Culcoa.)

G. cymosum ; Cunningham in Bot. Mag. t. 3131 (1832) ...	...	...	...	...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,19	M.fr.VII,74 ...	...	As., P.		

ELACHANTHERA ; F. v. M., in Vict. Naturalist III, 108 (1886).

E. Sewelliae ; F. v. M. in Vict. Natural. III, 108 (1886) ...	...	...	...	—	—	—	—		—	—	N.A.	—					

DRACAENA ; Vandelli, dissertatio de Dracaena t. 1 (1762).

D. angustifolia ; Roxburgh, Fl. Indic. II, 155 (1832) ...	...	...	...	—	—	—	—		—	Q.	N.A.	B.fl.VII,20	M.fr.V,194; VI,120; VIII,279; IX,190.				[P. As.,

CORDYLINE ; Commerçon in R. Hedwig, gen. plant. 243 (1806). (Charlwoodia.)

C. terminalis ; Kunth in act. acad. Berol. 30 (1820) ...	...	...	...	—	—	—	—		N.S.W.	Q.	—	B.fl.VII,21	M.fr.V,196 ...	...	As., P.		
C. Murchisoniae ; F. v. M., fragm. V, 195 (1866) ...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.VII,22	M.fr.V,195				
C. stricta ; Endlicher in Ann. Wien. Mus. I, 162 (1838) ...	...	...	...	—	—	—	—		N.S.W.	Q.	—	B.fl.VII,22	M.fr.V,195; VI,121.				
C. Baueri ; J. Hooker in Gardeners' Chronicle 792 (1860) ...	...	...	...	—	—	—	—		N.S.W.	—	—	—					

PHORMIUM ; R. & G. Forster, charact. gener. 47, t. 24 (1776). (Chlamydia.)

P. tenax ; R. & G. Forster, char. gen. 47, t. 24 (1776) ...	...	...	...	—	—	—	—		N.S.W.	—	—	--	...	...	...	N.Z.	

BLANDFORDIA ; Smith, Exot. Bot. I, 5, t. 4 (1804).

B. marginata ; Herbert in Bot. Regist. Misc. 84 (1842) ...	...	...	...	—	—	T.	—		—	—	—	B.fl.VII,23	M.fr.VII,71.				
B. grandiflora ; R. Brown, prodr. 296 (1810) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VII,23				
B. nobilis ; Smith, Exot. Bot. I, 5, t. 4 (1804) ...	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VII,24	M.fr.VII,70.			
B. flammea ; Hooker, Bot. Mag. t. 4819 (1854) ...	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,24				

HEWARDIA ; Hooker, icon. plant. 858 (1852).

H. Tasmanica ; Hooker, icon. plant. t. 858 (1852) ...	...	...	...	...	...	—	—	T.	—	—	—	—	B.fl.VII,25	M.fr.VII,72.			

ASTELIA ; Banks & Solander in R. Brown, prodr. 291 (1810). (Hamelinia, Funkia, 1805).

A. alpina ; R. Brown, prodr. 291 (1810)	...	...	...	...	...	...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,11	

MILLIGANIA ; J. Hooker in Kew misc. V, 296, t. 9 (1853).

M. densiflora ; J. Hooker, Fl. Tasman. II, 62 (1860)	...	...	...	...	—	—	T.	—	—	—	—	B.fl.VII,26			
M. longifolia ; J. Hooker in Kew misc. V, 296, t. 9 (1853)	...	...	...	—	—	T.	—	—	—	—	B.fl.VII,26				
M. Johnstoni ; F. v. M. in Benth. Fl. Austr. VII, 26 (1878)	...	...	...	—	—	T.	—	—	—	—	B.fl.VII,26				
M. stylosa ; F. v. M. in papers of the R. S. of Tasm. 11 (1876)	...	...	...	—	—	T.	—	—	—	—	B.fl.VII,27				

WURMBSEA ; Thunberg, nov. gen. I, 18 (1781). (Anguillaria, Wurmbea, Melanthium partly, Pleea partly.)

W. tubulosa ; Bentham, Fl. Austr. VII, 28 (1878)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,28	M.fr.VII,75.	
W. dioica ; F. v. M., fragm. X, 119 (1877)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,30	M.fr.VII,74.	

IPHIGENIA ; Kunth, enum. plant. omn. hucusq. cognit. IV, 212 (1843)

I. Indica ; Kunth, enum. plant. IV, 213 (1843)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,31	M.fr.VII,74; IX,193	... As.

SCHELHAMMERA ; R. Brown, prodr. 273 (1810). (Parduyna, 1806).

S. undulata ; R. Brown, prodr. 274 (1810)	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,31	M.fr.VII,71.	
S. multiflora ; R. Brown, prodr. 274 (1810)	...	...	...	...	...	—	—	—	—	—	Q.	—	B.fl.VII,32	M.fr.VII,72.	

KREYSSIGIA ; Reichenbach, iconogr. bot. exotic. III, 11, t. 229 (1829). (Kreysigia, Tripladenia.)

K. Cunninghami ; F. v. M., fragm. VII, 71 (1870)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,32	M.fr.VII,71; VIII,280.	

BURCHARDIA ; R. Brown, prodr. 272 (1810).

B. umbellata ; R. Brown, prodr. 272 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,33	M.fr.VII,77; VIII,280.	

BULBINE ; Linné, hort. Cliffort. 122 (1737). (Anthericum partly, Chrysobactron.)

B. bulbosa ; Haworth, revis. pl. succul. 33 (1821)	...	...	...	...	...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,34	M.fr.VII,70; XI,27.	
B. semibarbata ; Haworth, rev. pl. succ. 33 (1821)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,35	M.fr.VII,70.	

AGROSTOCRINUM ; F. v. M., fragm. phytogr. Austr. II, 95 (1860).

A. stypandroides ; F. v. M., fragm. II, 95 (1860)	..	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,36	M.fr.III,165; VII,65.	

THYSANOTUS ; R. Brown, prodr. 282 (1810). (Thysanella 1806, Chlamysporum 1809, Isandra.)

T. multiflorus ; R. Brown, prodr. 285 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,38	M.fr.VII,69.	
T. triandrus ; R. Brown, prodr. 284 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B fl.VII,39	M.fr.VII,69.	
T. glaucus ; Endlicher in Lehm. pl. Preiss. II, 38 (1846)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII.39			
T. Drummondii ; Baker in Journ. Linn. Soc. XV, 341 (1876)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,39				
T. pauciflorus ; R. Brown, prodr. 285 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,39		
T. asper ; Lindley, Bot. Regist. XXV, App. 58 (1839)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,39			
T. chryantherus ; F. v. M. in Bentham, flor. Hongk. 372 (1861)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,40	M.fr.V,202	...	...	As.
T. exiliflorus ; F. v. M. in Transact. R. S. of S. Austr. IV, 112 (1881)	...	...	—	S.A.	—	—	—	—	—	—	M.fr.XI,139.				
T. isantherus ; R. Brown, prodr. 283 (1810)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,40		
T. tenellus ; Endlicher in Lehm. pl. Preiss. II, 37 (1846)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,40			
T. scaber ; Endlicher in Lehm. pl. Preiss. II, 37 (1846)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,41			
T. tuberosus ; R. Brown, prodr. 282 (1810)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,41	M.fr.VII,69.	
T. exasperatus ; F. v. M., fragm. I, 21 (1858)	...	...	...	...	...	—	S.A.	—	—	—	—	—	B.fl.VII,42	M.fr.I,21.	
T. thyrsoideus ; Baker in Journ. Linn. Soc. XV, 336 (1876)	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,42				
T. Baueri ; R. Brown, prodr. 283 (1810)	...	...	...	...	...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,42	M.fr.I,22; VII,69.	
T. Patersoni ; R. Brown, prodr. 284 (1810)	...	...	...	...	...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,43	M.fr.VII,69; XI,20.	
T. junceus ; R. Brown, prodr. 283 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	—	—	B.fl.VII,43	M.fr.VII,69.	
T. dichotomus ; R. Brown, prodr. 284 (1810)	...	...	...	...	...	W.A.	S.A.	—	V.	—	—	—	B.fl.VII.44	M.fr.VII,69.	
T. arbuscula ; Baker in Journ. Linn. Soc. XV, 339 (1876)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,45			
T. anceps ; Lindley, Bot. Regist. XXV, App. 58 (1839)	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,45			

HODGSONIOLA ; F. v. M., fragm. II, 176 (1861). (Hodgsonia).

H. junciformis ; F. v. M., fragm. II, 176 (1861)	W.A.	—	—	—	—	—	—	B.fl.VII,46	M.fr.II,96,176.	

CAESIA ; R. Brown, prodr. fl. Nov. Holl. 277 (1810).

C. vittata ; R. Brown, prodr. 277 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,47	M.fr.I,63; VII,68.	
C. chlorantha ; F. v. M., fragm. I, 63 (1858)	—	—	—	—	—	Q.	N.A.	—	M.fr.I,63.	
C. parviflora ; R. Brown, prodr. 277 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,47	M.fr.VII,67.	
C. rigidifolia ; F. v. M., fragm. X, 48 (1876)...	W.A.	—	—	—	—	—	—	B.fl.VII,47	M.fr.X,48.	
C. setifera ; Baker in Journ. Linn. Soc. XV, 359 (1876)	—	—	—	—	—	—	N.A.	B.fl.VII,47		

CHAMAESCILLA ; F. v. M., fragm. VII, 68 (1870). (Caesia partly.)

C. corymbosa ; F. v. M., fragm. VII, 68 (1870)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,48	M.fr.VII,68.	
C. spiralis ; F. v. M., fragm. VII, 68 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,48	M.fr.VII,68.	

CORYNOTHECA ; F. v. M., fragm. VII, 68 (1870). (Caesia partly.)

C. lateriflora ; F. v. M., fragm. VII, 68 (1870)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,49	M.fr.I,215.	
C. dichotoma ; F. v. M., fragm. VII, 68 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,50	M.fr.I,215; XI,20.	
C. acanthoclada ; F. v. M., fragm. VII, 68 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,50	M.fr.I,215; XI,20.	

TRICORYNE ; R. Brown, prodr. fl. Nov. Holl. 278 (1810).

T. platyptera ; G. Reichenbach, Beitr. zur syst. Pflanzenk. 72 (1871)	—	—	—	—	—	Q.	—	B.fl.VII,51		
T. anceps ; R. Brown, prodr. 278 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,51	M.fr.VII,67.	
T. muricata ; Baker in Journ. Linn. Soc. XV, 363 (1876)	—	—	—	—	—	Q.	—	B.fl.VII,52		
T. elatior ; R. Brown, prodr. 278 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,52	M.fr.VII,67; XI,20.	
T. simplex ; R. Brown, prodr. 278 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,52	M.fr.VII,67.	
T. humilis ; Endlicher in Lehm. pl. Preiss. II, 36 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,53	M.fr.VII,67.	

STYPANDRA ; R. Brown, prodr. flor. Nov. Holl. 278 (1810).

S. glauca ; R. Brown, prodr. 278 (1810)	W.A.	—	—	V.	N.S.W.	Q.	—	B.fl.VII,53	M.fr.VII,64.	
S. caespitosa ; R. Brown, prodr. 279 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,54	M.fr.VII,65.	

ARTHROPODIUM ; R. Brown, prodr. 276 (1810). (Dichopogon, Siona.)

A. paniculatum ; R. Brown, prodr. 276 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,56	M.fr.VII,66.	
A. minus ; R. Brown, prodr. 276 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,56	M.fr.VII,67; XI,27.	
A. capillipes ; Endlicher in Lehm. pl. Preiss. II, 34 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,57	M.fr.VII,67.	
A. Preissii ; Endlicher in Lehm. pl. Preiss. II, 35 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,57	M.fr.VII,66.	
A. dianellaceum ; F. v. M., fragm. X, 65 (1876)	—	—	—	—	—	Q.	—	B.fl.VII,57	M.fr.X,65.	
A. strictum ; R. Brown, prodr. 276 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,58	M.fr.VII,66.	
A. fimbriatum ; R. Brown, prodr. 276 (1810)...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,59		

CHLOROPHYTUM ; Ker in Bot. Mag. t. 1071 (1808). (Chlorophyton, Phalangium partly.)

C. laxum ; R. Brown, prodr. 277 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,60	M.fr.I,63; VII,71 ...	As., Afr.
C. alpinum ; Baker in Journ. Linn. Soc. XV, 329 (1876)	—	—	T.	—	—	—	—	B.fl.VII,60		

HERPOLIRION ; J. Hooker, Fl. of New Zealand I, 258 (1853).

H. Novae Zelandiae ; J. Hooker, Fl. of New Zeal. I, 258 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,61	M.fr.VII,68; VIII,280.	N.Z.

SOWERBAEA ; Smith in Transact. Linn. Soc. V, 159 (1800). (Sowerbya.)

S. juncea ; Smith in Transact. Linn. Soc. V, 160, t. 6 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,61	M.fr.VI,180.	
S. laxiflora ; Lindley, Botan. Regist. new ser. IV, t. 10 (1841)	W.A.	—	—	—	—	—	—	B.fl.VII,62	M.fr.VI,180; VII,87.	
S. alliacea ; F. v. M., fragm. VI, 180 (1868)	—	—	—	—	—	—	N.A.	B.fl.VII,62	M.fr.VII,87.	

ALLANIA ; Endlicher, gen. plant. 151 (1837). (Alania.)

A. Endlicheri ; Kunth, enum. IV, 644 (1843)...	—	—	—	—	N.S.W.	—	—	B.fl.VII,63	M.fr.VI,180.	

BARTLINGIA; F. v. M., fragm. VII, 88 (1870). (Laxmannia.)

B. grandiflora; F. v. M., Cens. 118 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.VII,64	M.fr.I,159; VII,88.				
B. squarrosa; F. v. M., Cens. 118 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.VII,64					
B. minor; F. v. M., Cens. 118 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,65	M.fr.VII,88,89.				
B. gracilis; F. v. M., fragm. VII, 88 (1870)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,65	M.fr.I,159.				
B. ramosa; F. v. M., fragm. VII, 88 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,66	M.fr.VI,255.				
B. sessiliflora; F. v. M. in pap. R. S. of Tasm. 116 (1877)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,66					
B. brachyphylla; F. v. M., Cens. of Austr. pl. 119 (1882)	W.A.	—	—	—	—	—	—	B.fl.VII,66	M.fr.I,158.				
B. sessilis; F. v. M., Cens. 119 (1882) from Lindley (1839)	W.A.	—	—	—	—	—	—	B.fl.VII,67					
STAWELLIA; F. v. M., fragm. phytogr. Austr. VII, 85 (1870).													
S. dimorphantha; F. v. M., fragm. VII, 85 (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,67	M.fr.VII,85.				
JOHNSONIA; R. Brown, prodr. fl. Nov. Holl. 287 (1810).													
J. lupulina; R. Brown, prodr. 287 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,68	M.fr.VII,86.				
J. pubescens; Lindley, Bot. Regist. XXV, App. 57 (1839)...	W.A.	—	—	—	—	—	—	B.fl.VII,68	M.fr.VI,255; VII,87.				
ARNOCRINUM; Endlicher & Lehmann in pl. Preiss. II, 41 (1846).													
A. Drummondii; Endlicher in Lehm. pl. Preiss. II, 41 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,69	M.fr.VII,73.				
A. Preissii; Lehmann, pl. Preiss. II, 42 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,70					
BORYA; Labillardière, Nov. Holl. pl. specim. 81, t. 107 (1804).													
B. nitida; Labillardière, Nov. Holl. pl. spec. 81, t. 107 (1804)	W.A.	—	—	—	—	—	—	B.fl.VII,71	M.fr.VII,87.				
B. septentrionalis; F. v. M., fragm. V, 41 (1865)	—	—	—	—	—	Q.	—	B.fl.VII,71	M.fr.V,41.				
CALECTASIA; R. Brown, prodr. fl. Nov. Holl. 263 (1810).													
C. cyanea; R. Brown, prodr. 264 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.VII,121					
BAXTERIA; R. Brown in Hook. Lond. Journ. of Bot. II, 492, t. 13-15 (1843).													
B. Australis; R. Brown in Hook. Lond. Journ. II, 492 (1843	W.A.	—	—	—	—	—	—	B.fl.VII,120					
XEROTES; R. Brown, prodr. 257 (1810). (Lomandra 1804, Acanthocarpus, Chamaexeros.)													
X. echinata; Cunningham in Benth. fl. Austr. VII, 112 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,111					
X. Serra; Endlicher in Lehm. pl. Preiss. II, 49 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,111	M.fr.VIII,211; XI,24.				
X. fimbriatus; F. v. M., frsgm. VIII, 211 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,111	M.fr.VIII,211.				
X. Banksii; R. Brown, prodr. 263 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,96	M.fr.VIII,213	P.			
X. dura; F. v. M. in Transact. Vict. Inst. 42 (1854)	—	S.A.	—	—	N.S.W.	—	—	B.fl.VII,97	M.fr.VIII,207.				
X. longifolia; R. Brown, prodr. 262 (1810)	—	S.A.	T.	V.	N.S.W.	Q	N.A.	B.fl.VII,98	M.fr.VIII,210.				
X. rigida; R. Brown, prodr. 262 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,99	M.fr.VIII,209.				
X. Sonderi; F. v. M., fragm. VIII, 206 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,99	M.fr.				
X. odora; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,99					
X. Brownii; F. v. M., fragm. VIII, 206 (1874)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,100	M.fr.VIII,206.				
X. Ordii; F. v. M., fragm. XI, 23 (1878)	W.A.	—	—	—	—	—	—	—	M.fr.XI,137.				
X. sororia; F. v. M., first gen. Report 15 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,100					
X. Endlicheri; F. v. M., fragm. VIII, 205 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,205.				
X. sericea; Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,207.				
X. purpurea; Endlicher in Lehm. pl. Preiss. II, 49 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,101	M.fr.VIII,213.				
X. Preissii; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,102					
X. effusa; Lindley in Mitch. Three Exped. II, 151 (1838)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,102	M.fr.VIII,209.				
X. micrantha; Endlicher in Lehm. pl. Preiss. II, 49 (1846)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,103	M.fr.VIII,212.				
X. Thunbergii; F. v. M., fragm. VIII, 208 (1874)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,103	M.fr.VIII,208.				
X. caespitosa; Bentham, Fl. Austr. VII, 104 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,104					
X. pauciflora; R. Brown, prodr. 261 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,104	M.fr.212.				
X. flexifolia; R. Brown, prodr. 260 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,105	M.fr.VIII,207.				

X. glauca ; R. Brown, prodr. 260 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,105	M.fr.VIII,208.
X. elongata ; Bentham, Fl. Austr. VII, 106 (1878)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,106	
X. rupestris ; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,106	
X. collina ; R. Brown, prodr. 260 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,107	
X. suaveolens ; Endlicher in Lehm. pl. Preiss. II, 50 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,107	M.fr.VIII,207.
X. turbinata ; Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,107	M.fr.VIII,211.
X. spartea ; Endlicher in Lehm. pl. Preiss. II, 51 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,108	M.fr.VIII,211.
X. juncea ; F. v. M. in Transact. Vict. Inst. 135 (1855)	—	S.A.	—	V.	—	—	—	B.fl.VII,108	M.fr.VIII,210.
X. leucocephala ; R. Brown, prodr. 260 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,109	M.fr.VIII,210; IX,195.
X. hastilis ; R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,110	M.fr.VIII,210; XI,54.
DASYPOGON ; R. Brown, prodr. fl. Nov. Holl. 263 (1810).									
D. bromelifolius ; R. Brown, prodr. 263 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,118	M.fr.II,112.
D. Hookeri ; Drummond in Hook. Lond. Journ. II, 168 (1843)	W.A.	—	—	—	—	—	—	B.fl.VII,119	M.fr.II,112; IV,173.
XANTHORRHOEA ; Smith in Transact. Linn. Soc. IV, 219 (1798).									
X. gracilis ; Endlicher in Lehm. pl. Preiss. II, 39 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,114	M.fr.IV,112.
X. pumilio ; R. Brown, prodr. 288 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,114	
X. macronema ; F. v. M., fragm. IV, 112 (1864)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,113	
X. hastilis ; R. Brown, prodr. 288 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,115	M.fr.IV,113.
X. resinosa ; Persoon, syn. pl. I, 370 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,115	M.fr.IV,113.
X. semiplana ; F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.fl.VII,115	M.fr.IV,111.
X. bracteata ; R. Brown, prodr. 288 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,116	
X. minor ; R. Brown, prodr. 288 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,114	M.fr.IV,112.
X. Tateana ; F. v. M. in Zeitschr. Oesterr. Apoth. Ver. XXIII, 294 (1885) ...	—	S.A.	—	—	—	—	—	—	
X. Australis ; R. Brown, prodr. 288 (1810)	—	—	T.	V.	—	—	—	B.fl.VII,116	M.fr.IV,110.
X. quadrangulata ; F. v. M., fragm. IV, 111 (1864)	—	S.A.	—	—	—	—	—	B.fl.VII,117	M.fr.IV,111.
X. Preissii ; Endlicher in Lehm. pl. Preiss. II, 39 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,117	M.fr.IV,110.
KINGIA ; R. Brown in King's Narrat. of survey of Austr. coasts II, 529 (1827).									
K. Australis ; R. Brown in King's Narr. of surv. II, 535 (1827)	W.A.	—	—	—	—	—	—	B.fl.VII,119	

PALMAE.

RAY, method. pl. emend. 135 (1703).—Genera 10, Species 25.

CALAMUS ; Linné, spec. plant. 325 (1753).									
C. Australis ; Martius, hist. nat. palm. III, 213 (1839)	—	—	—	—	—	Q.	—	B.fl.VII,134	M.fr.V,48.
C. Muelleri ; Wendland & Drude in Linnaea XXXIX, 193 (1871)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,135	
C. caryotoides ; A. Cunningham in Martius, hist. nat. palm. III, 212 (1839) ...	—	—	—	—	—	Q.	—	B.fl.VII,135	
BACULARIA ; F. v. M., fragm. VII, 103 (1870). (Linospadix.)									
B. monostachya ; F. v. M., fragm. VII, 103 (1870)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,136	M.fr.VII,82,193; XI,58.
B. minor ; F. v. M., fragm. XI, 58 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,137	M.fr.VIII,235; IX,195; XI,137.
KENTIA ; Blume in Bull. Neerl. 64 (1838). (Grisebachia, Hydriastele, Hedyscepe, Kentiopsis, Veitchia partly, Howea, Howiea.)									
K. Wendlandiana ; F. v. M., fragm. VII, 102 (1870)	—	—	—	—	—	Q.	N.A.	B.fl.VII,137	M.fr.VII,102.
K. Belmoreana ; C. Moore & F. v. M., fragm. VII, 99 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VII,137	M.fr.VIII,234.
K. Forsteriana ; C. Moore & F. v. M., fragm. VII, 100 (1870)	—	—	—	—	N.S.W.	—	—	—	
K. Baueri ; Seemann, fl. Vitiens. 269 (1869)	—	—	—	—	N.S.W.	—	—	—	M.fr.VII,102; VIII,233.
K. Canterburyana ; C. Moore & F. v. M., fragm. VII, 101 (1870)	—	—	—	—	N.S.W.	—	—	B.fl.VII,138	M.fr.VIII,234; X,119.
K. acuminata ; Wendland & Drude in Garcke, Linnaea XXXIX, 207 (1871) ...	—	—	—	—	—	—	N.A.	B.fl.VII,138	

CLINOSTIGMA ; Herm. Wendland in Seemann's Bonplandia X, 196 (1862). (Cyphokentia.)

C. Mooreanum ; F. v. M., fragm. VIII, 235 (1874) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,139	M.fr.VII,101; IX,195; X,120.							

PTYCHOSPERMA ; Labillardière in Mém. de l'Inst. ann. 1808, I, 251, pl. X (1809). (Seaforthia, Laccospadix, Archontophoenix, Drymophlaeus, Calyptrocalyx).

P. Laccospadix ; Bentham, Fl. Austr. VII, 140 (1878) ...	—	—	—	—	—	Q.	—	B.fl.VII,140	M.fr.XI,57.							
P. Beatricae ; F. v. M. in Melb. Chemist. Febr. (1882) ...	—	—	—	—	—	Q.	—	—								
P. Alexandrae ; F. v. M., fragm. V, 47, t. 43 (1865) ...	—	—	—	—	—	Q.	—	B.fl.VII,140	M.fr.V,47,196,213.							
P. Cunninghamii ; Herm. Wendland in Bot. Zeit. 346 (1858) ..	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,141	M.fr.VIII,222.							
P. elegans ; Blume, Rumphia II, 118 (1836) ...	—	—	—	—	—	Q.	—	B.fl.VII,141	M.fr.VIII,222.							
P. Normanbyi ; F. v. M., fragm. XI, 56 (1878) ...	—	—	—	—	—	Q.	—	B.fl.VII,142	M.fr.VIII,235.							

ARECA ; Linné, spec. plant. 1189 (1753) from Ray (1688).

A. Alicae ; F. v. M. in Regel's Gartenflora, 199-201 (1879)...	—	—	—	—	—	Q.	—	—	M.fr.XI,88.							

COCOS ; Linné, spec. plant. 1188 (1753).

C. nucifera ; Linné, spec. plant. 1188 (1753) ...	—	—	—	—	—	Q.	—	B.fl.VII,143	...	...	As., Afr., Am., P.					

CARYOTA ; Linné, gen. plant. 355 (1737).

C. Rumphiana ; Martius, hist. nat. palm. III, 195 (1839) ...	—	—	—	—	—	Q.	—	B.fl.VII,144	...	...	As., P.					

LICUALA ; Rumphius, herb. Amboin. I, 44, t. 9 (1741).

L. Muelleri ; Wendland & Drude in Garcke, Linnaea XXXIX, 223 (1871) ...	—	—	—	—	—	Q.	—	B.fl.VII,145	M.fr.VIII,221; X,120.	As.						

LIVISTONA ; R. Brown, prodr. fl. Nov. Holl. 267 (1810). (Corypha partly.)

L. Leichhardtii ; F. v. M., fragm. V, 49 (1865) ...	—	—	—	—	—	—	N.A.	B.fl.VII,146	M.fr.VIII,221; XI,55.							
L. Mariae ; F. v. M. in Giles's Geog. Trav. in Centr. Austr. 222 (1875) ...	—	S.A.	—	—	—	—	N.A.	—	M.fr.VIII,283; XI,54,55.							
L. Australis ; Martius, hist. nat. palm. III, 241 (1839) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,146	M.fr.XI,55.							

NIPACEAE.

Brogniart, enum. des genr. de plantes 15 (1843).—Genus 1, Species 1.

NIPA ; Wurmb in Verhandl. Batav. Genootsch. I, 349 (1779) from Camellus, Ray & Rumphius. (Nypa.)

N. fruticans ; Wurmb in Verhandl. Batav. Genootsch. I, 349 (1779) ...	—	—	—	—	—	Q.	—	—	M.fr.XI,128.	...	As., P.					

PANDANEAE.

R. Brown, prodr. fl. Nov. Holl. (1810).—Genera 2, Species 11.

PANDANUS ; Rumphius, herb. Amboin. IV, 139, t. 74 (1750). (Athrodactylis, Fiquetia.)

P. odoratissimus ; Linné, fil. suppl. 424 (1781) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,148	M.fr.IV,40; XI,55	...	As., P.					
P. pedunculatus ; R. Brown, prodr. 341 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,149	M.fr.IV,40; VIII,220; IX,193.							
P. Moorei ; F. v. M., Cens. 140 (1882)...	—	—	—	—	N.S.W.	—	—	—								
P. Forsteri ; Moore & F. v. M., fragm. VIII, 220 (1874) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,149	M.fr.VIII,220.							
P. Solms-Laubachii ; F. v. M. in der bot. Zeitung (1887) ...	—	—	—	—	—	Q.	—	—								
P. aquaticus ; F. v. M., fragm. V, 40 (1865) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,148	M.fr.VIII,220.							
P. monticola ; F. v. M., fragm. V, 40 (1865) ...	—	—	—	—	—	Q.	—	B.fl.VII,150	M.fr.VII,63; VIII,220; IX,193.							

FREYCINETIA ; Gaudichaud in Annal. des scienc. natur. III, 509 (1824).

F. insignis ; Blume, Rumphia I, 158, t. 42 (1835) ...	—	—	—	—	—	Q.	—	—	...	...	...	As.				
F. Gaudichaudii ; Brown & Bennett, pl. Jav. rar. 31, t. 9 (1838) ...	—	—	—	—	—	Q.	—	B.fl.VII,151	...	...	...	As.				
F. excelsa ; F. v. M., fragm. V, 39 (1865) ...	—	—	—	—	—	Q.	—	B.fl.VII,151	M.fr.V,39.							
F. Baueriana ; Endlicher, prodr. fl. ins. Norf. 25 (1833) ...	—	—	—	—	N.S.W.	—	—	—								

AROIDEAE.

A. L. de Jussieu, gen. plant. 23 (1789), from B. de Jussieu (1759).—Genera 6, Species 10.

TYPHONIUM ; Schott in Wien. Zeitschr. III, 72 (1829). (Arum partly.)

T. liliifolium ; F. v. M. in Schott, prodr. syst. aroid. 107 (1860) ... ... — — — — — — N.A. B.fl.VII,153 M.fr.VIII,187.
T. alismifolium ; F. v. M., fragm. VIII, 186 (1874) ... ... — — — — — Q. — B.fl.VII,153 M.fr.VIII,186.
T. Brownii ; Schott, synops. Aroid. I, 11 (1855) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,154 M.fr.VIII,187.
T. angustilobum ; F. v. M., fragm. X, 66 (1876) ... ... — — — — — Q. — B.fl.VII,154 M.fr.X,66.

AMORPHOPHALLUS ; Blume in Nouv. Ann. du Mus. III, 366 (1834). (Brachyspatha.)

A. variabilis ; Blume, Rumphia I, 146, t. 35 (1835) ... . — — — — — — N.A. B.fl.VII,155 M.fr.VIII,87. As.

COLOCASIA ; Ray, Method. 157 (1682). (Caladium partly, Alocasia.)

C. antiquorum ; Schott, Meletemat. botan. 18 (1832) ... ... — — — — — Q. N.A. B.fl.VII,155 M.fr.VIII,87. As., Afr., P.
C. macrorrhiza ; Schott, Meletemat. botan. 18 (1832) ... ... — — — — N.S.W. Q. — B.fl.VII,155 M.fr.VIII,87. As., P.

RHAPHIDOPHORA ; Hasskarl in Fl. Regensb. Beibl. II, 11 (1842). (Epipremnum partly.)

R. pertusa ; Schott in Seem. Bonplandia V, 45 (1857) ... ... — — — — — Q. — B.fl.VII,156 M.fr.VIII,87. As., P.

GYMNOSTACHYS ; R. Brown, prodr. fl. Nov. Holl. 337 (1810).

G. anceps ; R. Brown, prodr. 337 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,157 M.fr.VIII,87.

POTHOS ; Linné, amœn. acad. I, 410 (1747).

P. Loureiri ; Hooker & Arnott, Bot. of Beech. Voy. 220 (1841) ... ... — — — — N.S.W. Q. — B.fl.VII,158 M.fr.I,62; III,163; VIII, 87. As., P.

TYPHACEAE.

A. L. de Jussieu, Gen. plant. 25 (1789).—Genera 2, Species 2.

TYPHA ; Tournefort, inst. 530, t. 301 (1700) from Theophrastos, Dioscorides and Plinius.

T. angustifolia ; Linné, spec. plant. 971 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,159 M.fr.VII,116. E., As., Afr., Am., P., N.Z.

SPARGANIUM ; Tournefort, inst. 530, t. 302 (1700).

S. angustifolium ; R. Brown, prodr. 338 (1810) ... ... — — — V. N.S.W. Q. — B.fl.VII,160 M.fr.VII,117.

LEMNACEAE.

J. E. Gray, Nat. arrang. of Brit. pl. II, 729 (1821).—Genera 2, Species 6.

LEMNA ; Linné, syst. nat. 9 (1735); Fl. Lappon. 351 (1737). (Telmatophace, Spirodela.)

L. trisulca ; Linné, spec. plant. 970 (1753) ... ... — S.A. T. V. N.S.W. Q. — B.fl.VII,162 M.fr.VIII,188. E., As., Afr., Am.
L. minor ; Linné, spec. plant. 970 (1753) ... ... W.A. S.A. T. V. N.S.W. Q. N.A. B.fl.VII,163 M.fr.VIII,188. E., As., Afr., Am., P., N.Z.
L. oligorrhiza ; Kurz in Journ. Linn. Soc. IX, 267 (1867) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,163 M.fr.VIII,188. As., P.
L. polyrrhiza ; Linné, spec. plant. 970 (1753) ... ... — — — V. N.S.W. — — B.fl.VII,164 M.fr.VIII,188. E., As., Afr., Am.
L. gibba ; Linné, spec. plant. 970 (1753) ... ... W.A. — — — — — — B.fl.VII,163 M.fr.VIII,188. E., As., Afr., Am., N.Z.

WOLFFIA ; Horkel & Schleiden in Schlecht. Linnaea XIII, 389 (1839). (Wolfia.)

W. Michelii ; Schleiden, Beitr. zur Botan. 233 (1844) ... ... — S.A. — V. N.S.W. — — B.fl.VII,162 M.fr.VIII,187. E., As., Afr.

FLUVIALES.

Ventenat, Tabl. du règn. véget. II, 80 (1799).—Genera 10, Species 36.

NAJAS ; Linné, syst. nat. 9 (1735); gen. plant. 278 (1737). (Caulinia partly.)

N. major ; Allioni, Fl. Pedem. II, 221 (1785)... ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,181 M.fr.VIII,218. E., As., Afr., Am.
N. tenuifolia ; R. Brown, prodr. 345 (1810) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,181 M.fr.I,94; VIII,219. As., Afr.

LEPILAENA ; Drummond & Harvey in Hook. Kew misc. VII, 57 (1855). (Zannichellia partly, Hexatheca, Altheniae subgenus.)

L. Australis ; J. Drummond in Hook. Kew misc. VII, 58 (1855)	W.A.	S.A.	—	V.	—	—	—	B.fl.VII,179	M.fr.VIII,217.				
L. Preissii ; F. v. M., fragm. VIII, 217 (1874)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,180	M.fr.VIII,217.				

DIPLANTHERA ; Thouars, gen. nov. Madag. 3 (1806). (Halodule).

D. tridentata ; Thouars, gen. nov. Madag. 3 (1806)	—	—	—	—	—	—	N.A.	—	M.fr.VIII,218.	As., Afr.			

CYMODOCEA ; C. Koenig, Annals of Botany II, 96, t. 7 (1806). (Amphibolis, Graumuellera, Kernera partly.)

C. zosterifolia ; F. v. M., Cens. 121 (1882) from Agardh. (1822)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,177	M.fr.IV,113; VIII,218; IX,196.				
C. ciliata ; Ehrenberg in Garcke, Linnaea XXXV, 162 (1867)	—	—	—	—	—	Q.	N.A.	B.fl.VII,178	M.fr.VIII,218.	As., Afr.			
C. rotundata ; Ascherson & Schweinfurth in Neum. Anl. zu wiss. Beob. 361 (1875)	—	—	—	—	—	—	N.A.	—	M.fr.X,162.	As.			
C. serrulata ; Ascherson & Magnus in Neum. Anleit. zu wiss. Beob. 362 (1875)	—	—	—	—	—	Q.	—	B.fl.VII,178	M.fr.VIII,218.				
C. isoetifolia ; Ascherson in Garcke, Linnaea XXXV, 163 (1867)	—	—	—	—	—	Q.	N.A.	B.fl.VII,178	...	...	As.		

ZOSTERA ; Linné, Waesgoeta-Resa, 166-168 et fig. (1747).

Z. nana ; F. K. Mertens in Roth, enum. pl. phaner. Germ. 8 (1827)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,176	M.fr.VIII,218.	E., As., Afr., Am.			
Z. Tasmanica ; G. v. Martens in Garcke, Linnaea XXXV, 68 (1867)	—	S.A.	T.	V.	—	—	—	B.fl.VII,176	M.fr.VIII,218.				

POSIDONIA ; Koenig, Annals of Botany II, 95, t. 6 (1806). (Caulinia partly, Kernera partly.)

P. Australis ; J. Hooker, Fl. Tasman. II, 43 (1860)...	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,175	M.fr.VI,198; VIII,218.				

RUPPIA ; Linné, syst. nat. 9 (1735); gen. plant. 277 (1737).

R. maritima ; Linné, spec. plant. 127 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,174	M.fr.VIII,217.	E., As., Afr., Am., N.Z.			

POTAMOGETON ; Fuchs, histor. stirp. comment. 651 (1542).

P. tricarinatus ; F. v. M. & A. Bennett in Brit. Journ. XXV, 178 (1887) ...	—	—	—	—	—	Q.	N.A.	—					
P. natans ; Linné, spec. plant. 126 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,170	M.fr.VIII,217.	E., As., Afr., Am., P., N.Z.			
P. coloratus ; Hornemann, fl. Danic. t. 1449 (1813)	—	—	—	V.	—	—	—	—	...	...	E., As., Afr.		
P. polygonifolius ; Pourret in Mém. de Toulouse (1788)	—	—	T.	V.	N.S.W.	—	—	—	...	...	E., As., Afr.		
P. lucens ; Linné, spec. plant. 126 (1753)	—	—	—	V.	—	—	—	—	...	...	E., As., Afr., Am.		
P. Tepperi ; A. Bennett in Brit. Journ. XXV, 177 (1887)	—	S.A.	—	—	—	Q.	—	—					
P. Cheesemanii ; A. Bennett in. Brit. Journ. of Bot. XXI, 66 (1883)	—	—	T.	—	—	—	—	—	...	...	N.Z.		
P. Javanicus ; Hasskarl in Act. Soc. Ind. Neerl. I, 26 (1857)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,171	M.fr.VIII,217.	As.			
P. Drummondii ; Bentham, Fl. Austr. VII, 171 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,171					
P. perfoliatus ; Linné, spec. plant. 126 (1753)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,172	M.fr.VIII,217.	E., As., Afr., Am.			
P. crispus ; Linné, spec. plant. 126 (1753)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,172	M.fr.VIII,217.	E., As., Afr., Am.			
P. obtusifolius ; Mertens & Koch, Deutschl. Fl. I, 855 (1823)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,172	M.fr.VIII,216.	E., As., Afr., Am., N.Z.			
P. acutifolius ; Link in Roem. & Schult. syst. veg. III, 513 (1818)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,173	M.fr.VIII,216.	E., As.			
P. pectinatus ; Linné, spec. plant. 127 (1753)...	—	S.A.	T.	V.	—	—	—	B.fl.VII,173	M.fr.VIII,217.	E., As., Afr., Am., N.Z.			

APONOGETON ; Linné, fil. suppl. plant. 32 (1781).

A. monostachyus ; Linné fil., suppl. 214 (1781)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,188	M.fr.VIII,215.	As.			
A. elongatus ; F. v. M. in Benth. Fl. Austr. VII, 188 (1878)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,188	M.fr.VIII,216.				

TRIGLOCHIN ; Rivinus in Ruppius, fl. Jenens. 54 et 366 (1718) from Dalechamps (1586). (Cycnogeton, Maundia.)

T. calcitrapa ; Hooker, icon. pl. t. 731 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	—	—	M.fr.VI,82.				
T. centrocarpa ; Hooker, icon. plant. t. 728 (1848)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,167	M.fr.VI,82.				
T. striata ; Ruiz & Pavon, Fl. Per. et Chil. III, 72 (1802)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,166	M.fr.VI,83.	Am., N.Z.			
T. mucronata ; R. Brown, prodr. 343 (1810)	W.A.	S.A.	—	V.	—	—	—	B.fl.VII,168	M.fr.VI,81.				
T. procera ; R. Brown, prodr. 343 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,168	M.fr.VI,83.				
T. Maundii ; F. v. M., fragm. VI, 83 (1867)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,169	M.fr.I,23.				

ALISMACEAE.

VENTENAT, Tabl. du règn. véget. II, 157 (1799).—Genera 3, Species 6.

ALISMA ; Rivinus in Ruppius, fl. Jen. 54 (1718), from Dioscorides and Plinius. (Caldesia ; Echinodorus partly.)

A. Plantago ; Linné, spec. plant. 842 (1753) ...	...	...	...	...	...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,185	M.fr.VIII,215. E., As., Afr., Am.
A. acanthocarpum ; F. v. M., fragm. I, 23 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,185	M.fr.VIII,214.
A. oligococcum ; F. v. M., fragm. I, 23 (1858)	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,185	M.fr.VI,169; VIII,214. As., Afr.
A. parnassifolium ; Bassi in Comment. Scienc. inst. Bonon. VI, 13, t. 1 (1768)						—	—	—	—	—	Q.	—	B.fl.VII,186	M.fr.VIII,214. E., As., Afr.

DAMASONIUM ; Tournefort, inst. 256 (1700). (Actinocarpus.)

D. Australe ; Salisbury in Transact. hort. Soc. Lond. I, 268 (1812)	...	...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,186	M.fr.VIII,215.			

TENAGOCHARIS ; Hochstetter in Regensb. bot. Zeit. June (1841). (Butomopsis.)

T. Cordofana ; Hochstetter in Regensb. bot. Zeit. June (1841)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,187	M.fr.X,103.		As., Afr.

PONTEDERIACEAE.

HUMBOLDT, Bonpland & Kunth, nov. gen. et spec. plant. I, 265 (1815).—Genus 1, Species 1.

MONOCHORIA ; Presl, reliq. Haenk. I, 127 (1827). (Limnostachys.)

M. cyanea ; F. v. M., fragm. VIII, 44 (1872)...	...	...	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,74	M.fr.VIII,44.

PHILHYDREAE.

R. BROWN in Flinders' Voy. II, 57 (1814).—Genera 3, Species 4.

PHILHYDRUM ; Banks in Gaertner, de fruct. I, 62, t. 16 (1788). (Philydrum.)

P. lanuginosum ; Banks in Gaertn. de fruct. I. 62, t. 16 (1788)	...	...	...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,73	M.fr.V,203.		As.

PRITZELIA ; F. v. M., Papuan plants 13 (1875). (Hetaeria, Philydrella.)

P. pygmaea ; F. v. M., Papuan plants 13 (1875)	...	...	...	...	...	W.A.	—	—	—	—	—	—	B.fl.VII,74	M.fr.V,203.

HELMHOLTZIA ; F. v. M., fragm. V, 202 (1866).

H. acorifolia ; F. v. M., fragm. V, 203 (1866)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,75	M.fr.V,203.
H. glaberrima ; Caruel in A. & C. de Candolle, monogr. phanerog. III, 6 (1881)						—	—	—	—	—	Q.	—	—	

COMMELINEAE.

R. BROWN, prodr. fl. Nov. Holl. 268 (1810).—Genera 6, Species 19.

ZYGOMENES ; Salisbury in Transact. hort. Soc. I, 271 (1812). (Cyanotis).

Z. axillaris ; Salisbury in Transact. hort. Soc. I, 271 (1812)	...	...	...	—	—	—	—	—	Q.	N.A.	B.fl.VII,82	M.fr.VIII,62.		As.

COMMELINA ; Plumier, nov. pl. Amer. gen. 48, t. 38 (1703). (Commelynia.)

C. ensifolia ; R. Brown, prodr. 269 (1810)	...	...	...	...	...	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,83	M.fr.VIII,60; IX,191. As., P.
C. cyanea ; R. Brown, prodr. 269 (1810)	...	...	...	...	...	...	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,84	M.fr.VIII,59; IX,191. P.
C. lanceolata ; R. Brown, prodr. 269 (1810)	...	...	...	...	...	...	—	—	—	—	Q.	N.A.	B.fl.VII,84	M.fr.VIII,59.
C. agrostophylla ; F. v. M. in J. Hooker, Fl. Tasman. p. 47 (1860)		...	...	—	—	—	—	—	Q.	N.A.	—	M.fr.VIII,59.		

ANEILEMA ; R. Brown, prodr. fl. Nov. Holl. 270 (1810).

A. acuminatum ; R. Brown, prodr. 270 (1810)	...	...	...	...	...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,85	M.fr.VIII,61.
A. biflorum ; R. Brown, prodr. 270 (1810)	...	..	...	...	...	...	—	—	—	N.S.W.	Q.	—	B.fl.VII,86	
A. sclerocarpum ; F. v. M., fragm. VIII, 61 (1873)...		...	...	...	...	...	—	—	—	—	Q.	—	B.fl.VII,86	M.fr.VIII,61.

A. siliculosum; R. Brown, prodr. 270 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 87	M.fr. VIII, 61; IX, 191.					
A. gramineum; R. Brown, prodr. 270 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 87	M.fr. VIII, 62.					
A. giganteum; R. Brown, prodr. 271 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 88	M.fr. VIII, 62.	As., Afr.				
FLORISCOPA; Loureiro, Fl. Cochinch. I, 192 (1790). (Floscopa, Dithyrocarpus.)														
F. scandens; Loureiro, fl. Cochinch. I, 193 (1790)	—	—	—	—	—	Q.	—	B.fl. VII, 89	M.fr. VIII, 63.	As., P.				
POLLIA; Thunberg, nov. gen. plant. I, 11 (1781).														
P. cyanococca; F. v. M., fragm. V, 40 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 90	M.fr. VIII, 63.					
CARTONEMA; R. Brown, prodr. fl. Nov. Holl. 271 (1810).														
C. philhydroides; F. v. M., fragm. I, 62 (1858)	W.A.	—	—	—	—	—	—	B.fl. VII, 91	M.fr. VIII, 64.					
C. spicatum; R. Brown, prodr. 271 (1810)	—	—	—	—	—	—	N.A.	B.fl. VII, 91	M.fr. I, 62; VIII, 64.					
C. parviflorum; Hasskarl in Flora Regensb. 365 (1869)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 91						
C. trigonospermum; Clarke in De Candolle, mon. phaner. III, 24 (1881)	—	—	—	—	—	—	N.A.	—	M.fr. XI, 139.					
C. brachyantherum; Bentham, Fl. Austr. VII, 92 (1878)	—	—	—	—	—	Q.	—	B.fl. VII, 92						
C. tenue; Bentham in De Candolle, monogr. phaner. III, 264 (1881)	—	—	—	—	—	—	N.A.	—	M.fr. XI, 139.					

XYRIDEAE.

SALISBURY in Transact. of the Hortic. Soc. I, 326 (1812).—Genus 1, Species 9.

XYRIS; Gronovius in Linné, genera plantar. 11 (1737).														
X. elongata; Rudge in Trans. Linn. Soc. X, 289, t. 15 (1809)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 77	M.fr. VIII, 205.					
X. pauciflora; Willdenow, phytogr. 2, t. 1 (1794)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 77	M.fr. VIII, 205.	As.				
X. lacera; R. Brown, prodr. 257 (1810)	W.A.	—	—	—	—	—	—	B.fl. VII, 78	M.fr. VIII, 204.					
X. flexifolia; R. Brown, prodr. 256 (1810)	W.A.	—	—	—	—	—	—	B.fl. VII, 78	M.fr. VIII, 204.					
X. gracilis; R. Brown, prodr. 256 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII, 79	M.fr. VIII, 204.					
X. operculata; Labillardière, Nov. Holl. pl. spec. I, 14, t. 10 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII, 79	M.fr. VIII, 204.					
X. lanata; R. Brown, prodr. 257 (1810)	W.A.	—	—	—	—	—	—	B.fl. VII, 80	M.fr. VIII, 204.					
X. laxiflora; F. v. M., fragm. VIII, 203 (1874)	W.A.	—	—	—	—	—	—	B.fl. VII, 80	M.fr. VIII, 203.					
X. gracillima; F. v. M., fragm. VIII, 203 (1874)	W.A.	—	—	—	—	—	—	B.fl. VII, 80	M.fr. VIII, 203.					

FLAGELLARIACEAE.

ENDLICHER, gener. plantar. 131 (1836).—Genus 1, Species 1.

FLAGELLARIA; Linné, Amoenit. academ. 396 (1747).														
E. Indica; Linné, spec. plant. 333 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 10	...	...	As., Afr., P.			

JUNCEAE.

R. BROWN, prodr. fl. Nov. Holl. 257 (1810).—Genera 2, Species 16.

LUZULA; De Candolle, Fl. franc. III, 158 (1805).														
L. campestris; De Candolle, Fl. franc. III, 158 (1805)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII, 123	—	...	E., As., Afr., Am., P., N.Z.			
L. longiflora; Bentham, Fl. Austr. VII, 123 (1878)	—	—	—	—	N.S.W.	—	—	B.fl. VII, 123						
JUNCUS; Tournefort, inst. 246, t. 127 (1700), from Camerarius, J. & C. Bauhin, Ray & Morison.														
J. gracilis; R. Brown, prodr. 259 (1810)	W.A.	—	—	—	—	—	—	B.fl. VII, 125						
J. planifolius; R. Brown, prodr. 259 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII, 125	...	...	N.Z.			
J. caespititius; E. Meyer in Lehm. pl. Preiss. II, 47 (1846)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII, 126						
J. falcatus; E. Meyer, synops. luzular. 34 (1823)	—	—	T.	V.	N.S.W.	—	—	B.fl. VII, 126	...	...	Am.			

J. bufonius; Linné, spec. plant. 328 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,127			E., As., Afr., Am., P., N.Z.
J. homalocaulis; F. v. M., first gen. Rep. 19 (1853)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,128			
J. Brownii; F. v. M., first gen. Rep. 19 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,128	M.fr.IX,78.		
J. communis; E. Meyer, synops. Juncor. 12 (1822)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,128			E., As., Afr., Am., P., N.Z.
J. vaginatus; R. Brown, prodr. 258 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,129			
J. pauciflorus; R. Brown, prodr. 259 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,129			
J. pallidus; R. Brown, prodr. 258 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,130			N.Z.
J. maritimus; Lamarck, Encycl. méth. III, 264 (1789)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,130			As., Afr., N.Z.
J. prismatocarpus; R. Brown, prodr. 259 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,131			
J. stipulatus; Meyen & Nees in Act. Acad. Caes. L.C. XVIII, suppl. 127 (1843)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,132			N.Z.

ERIOCAULEAE.

Humboldt, Bonpland & Kunth, Nov. gen. et spec. plant. I, 251 (1815).—Genus 1, Species 18.

ERIOCAULON; Linné, gen. plant. ed. sec. 35 (1742). (Electrosperma)

E. setaceum; Linné, spec. plant 87 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,191			As.
E. Australe; R. Brown, prodr. 254 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,192			
E. quinquangulare; Linné, spec. plant. 87 (1753)	—	—	—	—	—	—	N.A.	B.fl.VII,192			As.
E. Smithii; R. Brown, prodr. 254 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,192			
E. nanum; R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,193			
E. cinereum; R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,193	M.fr.I,95.		As.
E. pusillum; R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,194			
E. pallidum; R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,194			
E. nigricans; R. Brown, prodr. 254 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,194			
E. electrospermum; F. v. M., Cens. 123 (1882)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,195			
E. lividum; F. v. M., fragm. I, 92 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,195	M.fr.I,92.		
E. concretum; F. v. M., fragm. I, 92 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,195	M.fr.I,92.		
E. Schultzii; Bentham, Fl. Austr. VII, 195 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,195			
E. tortuosum; F. v. M., fragm. I, 94 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,196	M.fr.I,91.		
E. monoscapum; F. v. M., fragm. I, 94 (1858)	—	—	—	—	—	—	N.A.	B.fl.VII,196	M.fr.I,94.		
E. spectabile; F. v. M., fragm. I, 95 (1858)	—	—	—	—	—	Q.	N.A.	B.fl.VII,196	M.fr.I,95.		
E. scariosum; R. Brown, prodr. 255 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,197			
E. depressum; R. Brown, prodr. 255 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,197	M.fr.I,92.		

RESTIACEAE.

R. Brown, prodr. flor. Nov. Holl. 243 (1810).—Genera 14, Species 93.

TRITHURIA; J. Hooker, Fl. Tasman. II, 79, t. 138 (1860). (Juncella, 1854.)

T. submersa; J. Hooker, Fl. Tasman. II, 79, t. 138 (1860)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,199	M.fr.I,96; VIII,237.		

APHELIA; R. Brown, prodr. fl. Nov. Holl. 251 (1810). (Brizula.)

A. cyperoides; R. Brown, prodr. 252 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,200	M.fr.VIII,236.		
A. nutans; J. Hooker in Benth. Fl. Austr. VII, 200 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,200			
A. gracilis; Sonder in Schlecht. Linnaea XXVIII, 227 (1855)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,201	M.fr.VIII,237.		
A. Drummondii; Bentham, Fl. Austr. VII, 201 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,201			
A. Pumilio; F. v. M., first gen. Report 20 (1853)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,201	M.fr.VIII,237.		
A. Brizula; F. v. M., fragm. V, 203 (1866)	W.A.	—	—	—	—	—	—	B.fl.VII,202	M.fr.VIII,237.		

CENTROLEPIS; Labillardière, Nov. Holl. pl. specim. I, 7 (1804). (Desvauxia, Devauxia, Alepyrum.)

C. humillima; F. v. M. in Benth. Fl. Austr. VII, 203 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,203			
C. polygyna; Hieronymus in Abh. naturf. Ges. Halle XII, 96 (1873)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,203	M.fr.VIII,237.		

C. alepyroides; Hieronymus in Abh. naturf. Ges. Halle XII, 96 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,204	M.fr.VIII,237.
C. mutica; Hieronymus in Abh. nat. Ges. Halle XII, 97 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,204	
C. glabra; F. v. M. in Abh. nat. Ges. Halle XII, 95 (1873) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,204	M.fr.VIII,237.
C. muscoides; Hieronymus in Abh. nat. Ges. Halle XII, 95 (1873) ...	—	—	T.	—	—	—	—	B.fl.VII,205	
C. monogyna; Bentham, Fl. Austr. VII, 205 (1878) ...	—	—	T.	—	—	—	—	B.fl.VII,205	
C. pulvinata; Desvaux in Ann. des sc. nat. XIII, 42, t. 2 (1828)...	—	—	T.	—	—	—	—	B.fl.VII,205	
C. pusilla; Roemer & Schultes, syst. veg. I, 44 (1817) ...	—	—	—	—	—	Q.	—	B.fl.VII,205	
C. aristata; Roemer & Schultes, syst. veg. I, 44 (1817) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,206	M.fr.VIII,237.
C. Drummondii; Hieronymus in Abh. nat. Ges. Halle XII, 98 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,206	M.fr.VIII,237.
C. Banksii; Roemer & Schultes, syst. veg. I, 44 (1817) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,207	M.fr.VIII,237.
C. fascicularis; Labillardière, Nov. Holl. pl. spec. I, 7, t. 1 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,207	
C. pilosa; Hieronymus in Abh. nat. Ges. Halle 102 (1873)...	W.A.	—	—	—	—	—	—	B.fl.VII,207	
C. strigosa; Roemer & Schultes, syst. veg. I, 43 (1817 ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,207	
C. exserta; Roemer & Schultes, syst. veg. I, 44 (1817) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,208	M.fr.VIII,237.
LYGINIA; R. Brown, prodr. fl. Nov. Holl. 248 (1810). (Schoenodum partly.)									
L. barbata; R. Brown, prodr. 248 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,210	M.fr.VIII,79.
ECDEIOCOLEA; F. v. M., fragm. phytogr. Austr. VIII, 236 (1874).									
E. monostachya; F. v. M., fragm. VIII, 236 (1874)...	W.A.	—	—	—	—	—	—	B.fl.VII,211	M.fr.VIII,236.
ANARTHRIA; R. Brown, prodr. fl. Nov. Holl. 248 (1810).									
A. scabra; R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,212	M.fr.VIII,80.
A. laevis; R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,212	M.fr.VIII,81.
A. gracilis; R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,213	M.fr.VIII,72,81.
A. prolifera; R. Brown, prodr. 249 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,213	M.fr.VIII,82.
A. polyphylla; Nees in Lehm. pl. Preiss. II, 63 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,214	M.fr.VIII,83.
LEPYRODIA; R. Brown, prodr. fl. Nov. Holl. 247 (1810).									
L. scariosa; R. Brown, prodr. 248 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,215	M.fr.VIII,72.
L. Muelleri; Bentham, Fl. Austr. VII, 215 (1878) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,215	
L. anarthria; F. v. M., fragm. VIII, 73 (1873) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,216	
L. gracilis; R. Brown, prodr. 247 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,216	M.fr.VIII,73.
L. Tasmanica; J. Hooker, Fl. Tasman. II, 72, t. 135 (1860) ...	—	—	T.	V.	—	—	—	B.fl.VII,216	M.fr.VIII,73,75; IX,194.
L. interrupta; F. v. M., fragm. VIII, 74 (1873) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,217	M.fr.VIII,74.
L. hermaphrodita; R. Brown, prodr. 248 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,217	M.fr.VIII,75.
L. monoica; F. v. M., fragm. VIII, 76 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,217	M.fr.VIII.76.
L. Muirii; F. v. M., fragm. VIII, 78 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,218	M.fr.VIII,76.
L. stricta; R. Brown, prodr. 248 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,218	M.fr.VIII,74.
L. macra; Nees in Lehm. pl. Preiss. II, 60 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,218	
L. Drummondiana; Steudel, syn. pl. glum. II, 248 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.VII,219	M.fr.VIII,75.
L. glauca; F. v. M., fragm. VIII, 77 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,219	M.fr.VIII,77.
L. anaectocolea; F. v. M., fragm. VIII, 78 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,220	M.fr.VIII,78.
RESTIO; Linné, syst. nat. edit. duodec. II, 735 (1767). (Megalotheca.)									
R. fastigiatus; R. Brown, prodr. 246 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,222	M.fr.VIII,67.
R. Megalotheca; F. v. M., fragm. VIII, 99 (1873) ...	W.A.	—	—	—	—	—	—	B.fl.VII,222	
R. tropicus; R. Brown, prodr. 246 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,223	
R. applanatus; Sprengel, syst. I, 185 (1825) ...	W.A.	—	—	—	—	—	—	B.fl.VII,223	M.fr.VIII,69.
R. monocephalus; R. Brown, prodr. 245 (1810) ...	—	—	T.	—	—	—	—	B.fl.VII,224	M.fr.VIII,68.
R. dimorphus; R. Brown, prodr. 246 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,224	M.fr.VIII,68.
R. conferto-spicatus; Steudel, syn. glum. II, 256 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.VII,224	
R. sphacelatus; R. Brown, prodr. 246 (1810)...	W.A.	—	—	—	—	—	—	B.fl.VII,225	

R. deformis; R. Brown, prodr. 245 (1810)	W.A.	—	—	--	—	—	—	B.fl.VII,225	M.fr.VIII,65.
R. crispatus; R. Brown, prodr. 246 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,225	
R. nitens; Nees in Lehm. pl. Preiss. II, 59 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,226	M.fr.VIII,70.
R. gracilior; F. v. M. in Benth. Fl. Austr. VII, 226 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,226	
R. chaunocoleus; F. v. M., fragm. VIII, 64 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,226	M.fr.
R. Australis; R. Brown, prodr. 245 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,227	M.fr.VIII,69.
R. gracilis; R. Brown, prodr. 245 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,227	M.fr.VIII,69.
R. complanatus; R. Brown, prodr. 245 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,228	M.fr.VIII,67.
R. tetraphyllus; Labillardière, Nov. Holl. pl. spec. II, 77 (1806)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,228	M.fr.VIII,66.
R. laxus; R. Brown, prodr. 245 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,228	M.fr.VIII,71.
R. ornatus; Steudel, syn. pl. glum. II, 256 (1855)	W.A.	—	—	—	—	—	—	B.fl.VII,229	M.fr.VIII,70.
R. leptocarpoides; Bentham, Fl. Austr. VII, 229 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,229	M.fr.VIII,65.
R. amblycoleus; F. v. M., fragm. VIII, 65 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,230	M.fr.VIII,65.
R. tremulus; R. Brown, prodr. 230 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,230	M.fr.VIII,70.
LOXOCARYA; R. Brown, prodr. fl. Nov. Holl. 249 (1810). (Desmocladus.)									
L. densa; Bentham, Fl. Austr. VII, 241 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,241	
L. Benthami; F. v. M., Cens. 124 (1882) from Masters (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,242	
L. pubescens; Bentham, Fl. Austr. VII, 242 (1878)...	W.A.	—	—	—	—	—	—	B.fl.VII,242	M.fr.VIII,100.
L. fasciculata; Bentham, Fl. Austr. VII, 242 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,242	M.fr.VIII,98.
L. flexuosa; Bentham, Fl. Austr. VII, 243 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,243	M.fr.VIII,100.
L. cinerea; R. Brown, prodr. 249 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,243	
CALOSTROPHUS; Labillardière, Nov. Holl. pl. spec. II, 78 (1806). (Calorophus, Hypolaena.)									
C. elongatus; Labillardière, Nov. Holl. pl. spec. II, 78 (1806)	—	—	T.	—	—	—	—	B.fl.VII,238	M.fr.VIII,86; IX,94; X,120.
C. lateriflorus; F. v. M., fragm. VIII, 87 (1873)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,238	M.fr.VIII,87.
C. gracillimus; F. v. M., fragm. VIII, 88 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,239	M.fr.VIII,88.
C. fastigiatus; F. v. M. in pap. Roy. Soc. Tasm. 117 (1878)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,239	M.fr.VIII,84.
C. exsulcus; F. v. M., Cens. 124 (1882) from R. Brown (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,240	M.fr.VIII,86.
LEPTOCARPUS; R. Brown, prodr. 250 (1810). (Schoenodum partly.)									
L. scariosus; R. Brown, prodr. 250 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,232	M.fr.VIII,94.
L. tenax; R. Brown, prodr. 250 (1810)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,232	M.fr.VIII,96.
L. Brownii; J. Hooker, Fl. Tasman. II, 73, t. 136 (1860)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,233	M.fr.VIII,91.
L. canus; Nees in Ann. of nat. hist. VI, 50 (1843)	W.A.	—	—	—	—	—	—	B.fl.VII,234	M.fr.VIII,95.
L. coangustatus; Nees in Lehm. pl. Preiss. II, 65 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,234	M.fr.VIII,94.
L. aristatus; R. Brown, prodr. 250 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,235	M.fr.VIII,90.
L. erianthus; Bentham, Fl. Austr. VII, 235 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,235	
L. ramosus; R. Brown, prodr. 250 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,236	M.fr.VIII,92.
L. elatior; R. Brown, prodr. 250 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,236	
L. spathaceus; R. Brown, prodr. 250 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,236	M.fr.VIII,93.
L. Schultzii; Bentham, Fl. Austr. VII, 237 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,237	M.fr.VIII,93.
LEPIDOBOLUS; Nees in Lehm. pl. Preiss. II, 66 (1846).									
L. drapetocoleus; F. v. M., fragm. VIII, 84 (1873)...	—	S.A.	—	V.	—	—	—	B.fl.VII,244	M.fr.VIII,84.
L. Preissianus; Nees in Lehm. pl. Preiss. II, 66 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,245	M.fr.VIII,83.
L. chaetocephalus; F. v. M., fragm. VIII, 84 (1873)	W.A.	—	—	—	—	—	—	B.fl.VII,245	
CHAETANTHUS; R. Brown, prodr. 251 (1810). (Prionosepalum.)									
C. leptocarpoides; R. Brown, prodr. 251 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,246	M.fr.VIII,97.
ONYCHOSEPALUM; Steudel, syn. pl. glum. II, 249 (1855).									
O. laxiflorum; Steudel, syn. pl. glum. II, 249 (1855)	W.A.	—	—	—	—	—	—	B.fl.VII,246	M.fr.IX,51.

ACALYCEAE HYPOGYNAE.

F. v. M. in Woolls's plants of the neighb. of Sydney 48 (1880).

CYPERACEAE.

Haller, enum. stirp. Helvet. I, 234 (1742).—Genera 29, Species 380.

KYLLINGIA; Rottboell, descr. et icon. rar. et nov. plant. 13, t. 4 (1773). (Killiogia.)

K. intermedia; R. Brown, prodr. 219 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII,251	M.fr. VIII,27	...	As., P.
K. monocephala; Rottboell, descr. et icon. pl. 13, t. 4 (1773) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl. VII,251	M.fr. VIII,271; IX,54.		As., Afr., Am., P.
K. cylindrica; Nees in Wight's Contr. to Bot. of Ind. 91 (1834) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,252	M.fr. VIII,271	...	As.
K. triceps; Rottboell, descr. et icon. pl. 14, t. 4 (1773) ...	—	—	—	—	—	Q.	—	B.fl. VII,252	...	...	As., Afr.

CYPERUS; Tournefort, inst. 527, t. 299 (1700) from Hippocrates, Theophrastos & Plinius. (Pycreus, Mariscus, Auosporum, Sorostachys.)

C. pumilus; Linné, amoen. acad. IV, 302 (1759) ...	—	—	—	—	—	Q.	—	B.fl. VII,258	M.fr. VIII,267	...	As., Afr.
C. Eragrostis; Vahl, enum. plant. II, 322 (1806) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII,258	M.fr. VIII,260; IX,52.		As., Afr., P.
C. flavescens; Linné, spec. plant. 46 (1753) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,259	...	...	E., As., Afr.
C. globosus; Allioni, actuar. ad fl. Pedem. 49 (1789) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII,260	M.fr. VIII,260; IX,52.		E., As., Afr.
C. unioloides; R. Brown, prodr. 216 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII,260	M.fr. VIII,259; IX,52.		As., Afr., Am.
C. polystachyus; Rottboell, descr. et icon. pl. 39, t. 11 (1773) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII,261	M.fr. VIII,265,270; XI,53; XII,25		E., As.
C. Hochstetteri; Nees in Regensb. Flora 755 (1845) ...	—	—	—	—	—	—	N.A.	B.fl. VII,261	M.fr. VIII,265; IX,53.		As., Afr., Am.
C. lucidulus; Klein in Link, Jahrb. III, 86 (1820) ...	—	—	—	—	—	Q.	—	—	...	...	As.
C. pygmaeus; Rottboell, descr. et icon. pl. 20, t. 14 (1773) ...	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl. VII,262	M.fr. VIII,268; IX,53.		E., As., Afr., Am.
C. cephalotes; Vahl, enum. II, 311 (1806) ...	—	—	—	—	—	Q.	—	B.fl. VII,263	M.fr. VIII,272; IX,54.		As.
C. laevigatus; Linné, mantissa alt. 179 (1771) ...	W.A.	—	—	—	—	—	—	B.fl. VII,263	M.fr. VIII,266; IX,53.		E., As., Afr., Am.
C. platystylis; R. Brown, prodr. 241 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl. VII,264			
C. alopecuroides; Rottboell, descr. et icon. pl. 38, t. 8 (1773) ...	—	—	—	—	—	Q.	—	B.fl. VII,264	M.fr. VIII,263; IX,52.		As., Afr.
C. leucocephalus; Retzius, observ. V, 11 (1788) ...	—	—	—	—	—	Q.	N.A.	B.fl. VII,265	M.fr. VIII,271; XII,25.		As.
C. tenellus; Linné, fil. suppl. 103 (1781) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl. VII,265	M.fr. VIII,261; IX,53; XII,25.		Afr., N.Z.
C. gracilis; R. Brown, prodr. 213 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII,265	M.fr. VIII,264; IX,53; XII,25.		P.
C. enervis; R. Brown, prodr. 213 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,266			
C. debilis; R. Brown, prodr. 213 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl. VII,266			
C. castaneus; Willdenow, spec. plant. I, 278 (1797) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,267	...	...	As.
C. cuspidatus; Humboldt, Bonpland & Kunth, nov. gen. et sp. pl. I, 204 (1815)	—	—	—	—	—	Q.	—	B.fl. VII,267	...	...	As., Afr., Am.
C. squarrosus; Linné, amoen. acad. IV, 303 (1759) ...	—	—	—	—	—	—	N.A.	—			
C. ixiocarpus; F. v. M. in proc. R. S. Vict. XXIII, 53 (1885) ...	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl. VII,268	M.fr. VIII,262; IX,53; XII,25.		As., Afr., Am.
C. difformis; Linné, amoen. acad. IV, 302 (1759) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII,268	M.fr. VIII,262; IX,52.		E., As., Afr.
C. tetraphyllus; R. Brown, prodr. 214 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,269	M.fr. VIII,264.		
C. trinervis; R. Brown, prodr. 213 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII,269	M.fr. VIII,267; IX,53.		
C. Haspan; Linné, spec. plant. 45 (1753) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII,270	M.fr. VIII,260; IX,52.		As., Afr., Am., P.
C. concinnus; R. Brown, prodr. 214 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII,270	M.fr. VIII,261.		
C. filipes; Bentham, Fl. Austr. VII, 271 (1878) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,271			
C. pedunculosus; F. v. M., fragm. VIII, 266 (1874) ...	—	—	—	—	—	Q.	—	B.fl. VII,272	M.fr.		
C. vaginatus; R. Brown, prodr. 213 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII,272	M.fr. VIII,261; IX,53.		
C. Holoschoenus; R. Brown, prodr. 215 (1810) ...	—	S.A.	—	—	—	Q.	N.A.	B.fl. VII,273	M.fr. I,200; VIII,262.		
C. dactylotes; Bentham, Fl. Austr. VII, 273 (1878) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII,273			
C. Gilesii; Bentham, Fl. Austr. VII, 274 (1878) ...	—	S.A.	—	—	N.S.W.	Q.	—	B.fl. VII,274			
C. fulvus; R. Brown, prodr. 215 (1810) ...	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl. VII,274	M.fr. VIII,268.		
C. carinatus; R. Brown, prodr. 216 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII,274			
C. alterniflorus; R. Brown, prodr. 216 (1810) ...	W.A.	S.A.	—	—	—	Q.	—	B.fl. VII,275			
C. pilosus; Vahl, enum. II, 354 (1806) ...	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,275	M.fr. VIII,260; IX,52.		As.

C. procerus; Rottboell, descr. et. icon. 29, t. 5 (1773)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,276				
C. Iria; Linné, spec. plant. 45 (1753)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,276	M.fr.VIII,266; IX,53; XII,25.		As., Afr.	
C. eleusinoides; Wallich in Kunth. enum. II, 39 (1837)	—	—	—	—	—	Q.	N.A.	B.fl.VII,277	M.fr.VIII,264; IX,53.		As., Afr.	
C. distans; Linné, fil. suppl. 103 (1781)	—	—	—	—	—	Q.	—	B.fl.VII,277	M.fr.VIII,266; IX,53.		As., Afr., Am., P.	
C. Malaccensis; Lamarck, ill. des gen. I, 146 (1791)	—	—	—	—	—	—	N.A.	B.fl.VII,278	M.fr.VIII,260; IX,52.		As.	
C. articulatus; Linné, spec. plant. 44 (1753)	—	—	—	—	—	—	N.A.	B.fl.VII,278	...	...	...	As., Afr., Am.
C. corymbosus; Rottboell, desc. et icon. pl. 42, t. 7 (1773)	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,279				[Am., P.
C. rotundus; Linné, spec. plant. 45 (1755)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,279	M.fr.VIII,269; IX,53; XII,25.		As., Afr.,	
C. stenostachyus; Bentham, Fl. Austr. VII, 280 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,280				
C. Lhotzkyanus; Boeckeler in Engler's bot. Jahrb. 498 (1884)	—	—	—	—	N.S.W.	—	—	—				
C. flavidus; Retzius, obs. bot. V, 13 (1788)	—	—	—	—	—	Q.	—	—				
C. tenuiflorus; Rottboell, descr. et icon. pl. 30, t. 14, f. I (1773)	—	—	—	—	—	Q.	N.A.	—				
C. congestus; Vahl, enum. II, 350 (1806)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.VII,280	M.fr.VIII,269; IX,53.		Afr.	
C. subulatus; R. Brown, prodr. 217 (1810)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,281				
C. sporobolus; R. Brown, prodr. 215 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,281				
C. angustatus; R. Brown, prodr. 214 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,282	M.fr.IX,54.			
C. inornatus; Boeckeler in Fl. Regensb. 86 (1875)	—	—	—	—	—	Q.	—	B.fl.VII,282	M.fr.IX,54.			
C. lucidus; R. Brown, prodr. 218 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,283	M.fr.VIII,270; IX,53.			
C. pennatus; Lamarck, illustr. des genr. I, 144 (1799)	—	—	—	—	—	Q.	N.A.	B.fl.VII,284	M.fr.VIII,263; IX,53.		As., P.	
C. exaltatus; Retzius, observ. bot. V, 11 (1789)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,285	M.fr.VIII,263; IX,53.		As., P.	
C. haematodes; Endlicher, prodr. fl. Norf. 22 (1839)	—	—	—	—	N.S.W.	—	—	B.fl.VII,285	M.fr.IX,54.			
C. digitatus; Roxburgh, fl. Ind. I, 209 (1820)	—	—	—	—	—	Q.	—	B.fl.VII,286	M.fr.VIII,263	...	As., Afr., P.	
C. ferax; Cl. Richard in act. soc. hist. nat. Par. I, 106 (1792)	—	—	—	—	—	Q.	—	B.fl.VII,286	...	...	...	As., Afr., Am.
C. stoloniferus; Retzius, observ. IV, 10 (1786)	—	—	—	—	—	—	N.A.	—	...	..	...	As.
C. esculentus; Linné, sp. pl. 45 (1753)	—	—	—	—	—	Q.	—	—	..	...	...	E., As., Afr.
C. Bowmani; F. v. M. in Benth. Fl. Austr. VII, 287 (1878)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,287				
C. trichostachys; Bentham, Fl. Austr. VII, 287 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,287				
C. leiocaulis; Bentham, Fl. Austr. VII, 287 (1878)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,287				
C. scaber; Bentham, Fl. Austr. VII, 288 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.VII,288				
C. decompositus; F. v. M., fragm. VIII, 267 (1874)	—	—	—	—	—	Q.	N.A.	B.fl.VII,288	M.fr.XII,25.			
C. Armstrongii; Bentham, Fl. Austr. VII, 289 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.VII,289				
C. umbellatus; Bentham, Fl. Hongk. 386 (1861)	—	—	—	—	—	Q.	—	B.fl.VII,289	M.fr.VII,267; IX,53.		As., Afr., P.	
C. conicus; Boeckeler in Schlecht. Lionaea XXXVIII, 371 (1874)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VII,290	M.fr.VIII,268; IX,53.			
HELEOCHARIS; R. Brown, prodr. 224 (1810). (Eleocharis, Eleochiton, Scirpidium.)												
H. sphacelata; R. Brown, prodr. 224 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,292	M.fr.VI,94; VIII,239.		N.Z., P.	
H. spiralis; R. Brown, prodr. 224 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,292	...	..	...	As., Afr., Am.
H. compacta; R. Brown, prodr. 224 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,293	M.fr.VIII,239		...	As., Afr., Am.
H. fistulosa; J. A. & J. H. Schultes, mantiss. II, 89 (1824)	—	—	—	—	—	Q.	—	B.fl.VII,293	M.fr.VI,93; VIII,239.		As., Afr., Am.	
H. tetraquetra; Nees in Wight's Contrib. Bot. of Ind. 113 (1834)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,294	M.fr.VIII,239		...	As.
H. cylindrostachys; Boeckeler in Regensb. Flora, 108 (1875)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,294	M.fr.VIII,240.			
H. acuta; R. Brown, prodr. 224 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,294	M.fr.VIII,240.		...	Am., N.Z.
H. multicaulis; Smith, Engl. Flor. II, 64 (1824)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,296	...	...	..	E., Am., N.Z.
H. atricha; R. Brown, prodr. 225 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,295	M.fr.VII,252; IX,100.			
H. capitata; R. Brown, prodr. 225 (1810)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VII,296	M.fr.VI,94; VIII,240.		As., Afr., Am.	
H. atropurpurea; Kunth, enum. II, 151 (1837)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,296	M.fr.VIII,240		...	E., As., Afr., Am.
H. acicularis; R. Brown, prodr. 224 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,297	M.fr.VIII,240		...	E., As., Afr., Am.
FIMBRISTYLIS; Vahl, enum. plant. II, 285 (1806). (Abildgaardia, Trichelostylis, Oncostylis.)												
F. acicularis; R. Brown, prodr. 226 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,301	M.fr.IX,54.			
F. acuminata; Vahl, enum. II, 285 (1806)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,301	M.fr.VIII,274; IX,54.		As.	

F. punctata ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,302					
F. rhyticarya ; F. v. M., fragm. I, 215 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.VII,302	M.fr.I,215.				
F. leucostachya ; Boeckeler in Garcke, Linnaea XXXVIII, 385 (1874)	—	—	—	—	—	—	N.A.	B.fl.VII,302	M.fr.IX,54.				
F. nutans ; Vahl, enum. II, 285 (1806)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,303	M.fr.VIII,274; IX,54.			As. P.	
F. pauciflora ; R. Brown, prodr. 225 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,303	...	...	...	As.	
F. cardiocarpa ; F. v. M., fragm. I, 194 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,303	M.fr.I,194.				
F. leucocolea ; Bentham, Fl. Austr. VII, 304 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,304					
F. polytrichoides ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,304	...	...	...	As.	
F. androgyna ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,304					
F. subbulbosa ; Bentham, Fl. Austr. VII, 305 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,305					
F. tetragona ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,305	M.fr.I,194; VIII,274.			As.	
F. trigastrocarya ; F. v. M., fragm. I, 194 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,305	M.fr.I,194.				
F. monandra ; F. v. M., fragm. I, 195 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,306	M.fr.I,195.				
F. pterygosperma ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,306	M.fr.I,193; IX,55.				
F. sphaerocephala ; Bentham, Fl. Austr. VII, 306 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.VII,306					
F. xyridis ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,307	M.fr.VIII,274; IX,54.				
F. oxystachya ; F. v. M., fragm. I, 195 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,307	M.fr.VIII,272.				
F. macrantha ; Boeckeler in Garcke, Linnaea XXXVIII, 388 (1874)	—	—	—	—	—	—	N.A.	B.fl.VII,307	M.fr.IX,55.				
F. aquarrulosa ; F. v. M., fragm. I, 216 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.VII,308	M.fr.I,216.				
F. monostachya ; Hasskarl, pl. Jav. rar. 61-64 (1848)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,308	M.fr.VIII,272; IX,54.			As., Afr., Am.	
F. Brownii ; Bentham, Fl. Austr. VII, 308 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,308	M.fr.VIII,273.				
F. Dallachyi ; F. v. M. in Benth. Fl. Austr. VII, 309 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,309	M.fr.VIII,273.				
F. velata ; R. Brown, prodr. 227 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII.309	M.fr.IX,11,54	...		N.Z.	
F. aestivalis ; Vahl, enum. II, 288 (1806)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,310	M.fr.IX,11,54	...		As., Am., P.	
F. dichotoma ; Vahl, enum. II, 287 (1806)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,310	M.fr.IX,10,54	...		As., Afr., Am., P.	
F. depauperata ; R. Brown, prodr. 227 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,311					
F. spirostachya ; F. v. M. in Benth. Fl. Austr. VII, 311 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,311					
F. communis ; Kunth, enum. II, 234 (1837)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,311	M.fr.I,196; IX,10,54.			As., Afr., Am., P.	
F. ferruginea ; Vahl, enum. 291 (1806)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,312	M.fr.I,197; IX,10,54.			As., Afr., Am., P.	
F. denudata ; R. Brown, prodr. 227 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,313	M.fr.IX,9.				
F. elata ; R. Brown, prodr. 227 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,313					
F. caespitosa ; R. Brown, prodr. 228 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,313	M.fr.I,199.				
F. spiralis ; R. Brown, prodr. 226 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,314					
F. subaristata ; Bentham, Fl. Austr. VII, 314 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,314					
F. leptoclada ; Bentham, Fl. Austr. VII, 314 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,314					
F. debilis ; F. v. M., fragm. I, 198 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,315	M.fr.I,198.				
F. corynocarya ; F. v. M., fragm. I, 197 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,315	M.fr.I,187.				
F. solidifolia ; F. v. M., fragm. I, 198 (1859)	—	—	—	—	—	—	N.A.	B.fl.VII,315	M.fr.I,198.				
F. obtusangula ; F. v. M., fragm. I, 198 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.VII,315	M.fr.I,198.				
F. miliacea ; Vahl, enum. II, 287 (1806)	—	—	—	—	—	Q.	N.A.	B.fl.VII,316	M.fr.I,199; IX,12,54.			As., P.	
F. rara ; R. Brown, prodr. 227 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,316					
F. microcarya ; F. v. M., fragm. I, 200 (1859)	—	—	—	—	—	Q.	N.A.	B.fl.VII,316	M.fr.I,200.				
F. quinquangularis ; Kunth, enum. II, 229 (1837)	—	—	—	—	—	—	N.A.	B.fl.VII,317	...	...	...	As., Afr.	
F. cyperoides ; R. Brown, prodr. 228 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,317	M.fr.VIII,273; IX,11,54.			As.	
F. furva ; R. Brown, prodr. 228 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,318					
F. cymosa ; R. Brown, prodr. 228 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,318					
F. multifolia ; Boeckeler in Garcke, Linnaea XXXVIII, 397 (1874)	—	—	—	—	—	—	N.A.	B.fl.VII,319	M.fr.IX,55.				
F. sericea ; R. Brown, prodr. 228 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,319					
F. macrostachya ; Boeckeler in Garcke, Linnaea XXXVIII, 386 (1874)	—	—	—	—	—	—	N.A.	B.fl.VII,319	M.fr.IX,54.				
F. Neilsoni ; F. v. M., fragm. IX, 79 (1875)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,320	M.fr.IX,79.				
F. capitata ; R. Brown, prodr. 228 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,320	M.fr.I,196.				

F. Schultzii ; Boeckeler in Garcke, Linnaea XXXVIII, 391 (1874)	—	—	—	—	—	—	—	N.A.	B.fl.VII,321	M.fr.IX,55.					
F. barbata ; Bentham, Fl. Austr. VII, 321 (1878)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,321	M.fr.IX,55	...	...	As., Afr., P.			
F. capillaris ; A. Gray, Man. of Bot. N.U.S. fifth ed. 567 (1867)	W.A.	—	—	—	—	Q.	N.A.	B.fl.VII,322	M.fr.IX,55	...	...	As., Afr., Am.			
SCIRPUS ; Tournefort, inst. 528, t. 300 (1700) from Terentius. (Isolepis, Malacochaete.)															
S. humillimus ; Bentham, Fl. Austr. VII, 324 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,324	M.fr.IX,7.						
S. fluitans ; Linné, spec. plant. 48 (1753)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,325	M.fr.IX,55	...	...	E., As., Afr.			
S. arenarius ; Bentham, Fl. Austr. VII, 325 (1878)	W.A.	—	—	V.	—	—	—	B.fl.VII,325							
S. lenticularis ; Sprengel, syst. I, 208 (1825)	—	—	—	—	N.S.W.	—	—	B.fl.VII,326	M.fr.IX,55						
S. crassiusculus ; J. Hooker in Benth. Fl. Austr. VII, 326 (1878)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,326							
S. cyperoides ; Sprengel, syst. I, 208 (1825)	W.A.	—	—	—	—	—	—	B.fl.VII,326							
S. setaceus ; Linné, spec. plant. 49 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,327	M.fr.IX,55	...	...	E., As., Afr.			
S. riparius ; Sprengel, syst. I, 208 (1825)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,327	...	...	...	E., As., Afr., Am.			
S. cartilaginens ; Sprengel, syst. I, 208 (1825)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,328	M.fr.IX,55	...	...	Afr., N.Z.			
S. squarrosus ; Linné, mant. plant. alt. 181 (1771)	—	—	—	—	—	Q.	N.A.	B.fl.VII,329	...	...	...	As., Afr.			
S. inundatus ; Sprengel, syst. I, 207 (1825)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,329	M.fr.IX,55	...	...	N.Z.			
S. prolifer ; Rottboell, descr. et icon. pl. 55, t. 17 (1773)	—	—	—	—	N.S.W.	—	—	B.fl.VII,330	M.fr.IX,55	...	...	Afr.			
S. articulatus ; Linné, spec. plant. 47 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,331	M.fr.IX,55	...	...	As., Afr.			
S. nodosus ; Rottboell, descr. et icon. pl. 52, t. 8 (1773)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,331	M.fr.IX,55	...	...	Afr., Am., N.Z.			
S. supinus ; Linné, spec. plant. 49 (1753)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,330	M.fr.IX,55; XII,25	...	E., As., Afr., Am.				
S. debilis ; Pursch, flor. Amer. Sept. I, 55 (1814)	—	—	—	—	—	Q.	N.A.	B.fl.VII,332	...	...	...	As., Am.			
S. mucronatus ; Linné, spec. plant. 50 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,332	M.fr.IX,8,56	...	...	E., As.,			
S. pungens ; Vahl, enum. II, 255 (1806)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,333	M.fr.IX,8,56	...	...	E., Afr., Am., N.Z.			
S. lacustris ; Linné, spec. plant. 48 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,333	M.fr.IX,7,56	...	E., As., Afr., Am., N.Z.				
S. litoralis ; Schrader, Fl. Germ. I, 142, t. 5 (1806)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,334	M.fr.IX,7,56; XII,25	...	E., As., Afr.				
S. maritimus ; Linné, spec. plant. 51 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,335	M.fr.IX,8,56	...	E., As., Afr., Am., N.Z.				
S. grossus ; Linné, fil. suppl. plant. 104 (1781)	—	—	—	—	—	Q.	—	—	M.fr.XII,24	...	...	As.			
S. polystachyus ; F. v. M. in Transact. phil. Soc. Vict. I, 108 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,335	M.fr.IX,9,56.						
LIPOCARPHA ; R. Brown in Tuckey's narrat. of an Exped. to Congo, 459 (1818). (Hypaelyptum.)															
L. argentea ; R. Brown in Tuck. Congo, 459 (1818)	—	—	—	—	—	Q.	—	B.fl.VII,336	M.fr.VIII,238; IX,57.	As., Afr.					
L. microcephala ; R. Brown in Tuck. Congo, 459 (1818)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,337	M.fr.VIII,238; IX,57.	As., P.					
FUIRENA ; Rottboell, descr. et icon. rar. pl. illustr. 70 (1773).															
F. umbellata ; Rottboell, descr. et icon. pl. 70, t. 19 (1773)	—	—	—	—	—	Q.	N.A.	B.fl.VII,337	M.fr.VIII,238; IX,57.	As., Afr., Am., P.					
F. glomerata ; Lamarck, Illustr. des genr. I, 150 (1791)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,338	M.fr.VIII,238; IX,57.	As., Afr., P.					
HYPELYTRUM ; L. C. Richard in Persoon, synops. I, 70 (1805). (Hypolytrum.)															
H. latifolium ; L. C. Richard in Pers. syn. I, 70 (1805)	—	—	—	—	—	Q.	—	B.fl.VII,339	M.fr.VIII,238; IX,57.	As., Afr., Am., P.					
EXOCARYA ; Bentham in J. Hooker, icon. plant. t. 1206 (1877).															
E. scleroides ; Bentham in Hook. icon. pl. t. 1206 (1877)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,339	M.fr.IX,12.						
MAPANIA ; Aublet, Hist. pl. Guian. 47, t. 17 (1775). (Pandanophyllum.)															
M. hypelytroides ; F. v. M., fragm. IX, 16 (1875)	—	—	—	—	—	Q.	—	B.fl.VII,341	M.fr.IX,16	...	...	P.			
SCIRPODENDRON ; Zippelius in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869).															
S. costatum ; Kurz in Journ. Asiat. Soc. Beng. XXXVIII, 85 (1869)	—	—	—	—	—	Q.	—	B.fl.VII,341	M.fr.X,104	...	...	As., P.			
LEPIRONIA ; L. C. Richard in Persoon, synops. I, 70 (1805). (Chondrachne.)															
L. mucronata ; L. C. Richard in Persoon, syn. I, 70 (1805)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,342	M.fr.IX,17,58	...	As., Afr., P.				
CHORIZANDRA ; R. Brown, prodr. 221 (1810). (Chorisandra.)															
C. sparocephala ; R. Brown, prodr. 221 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,344	M.fr.IX,18,58.						
C. enodis ; Nees in Lehm. pl. Preiss. II, 73 (1846)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,344	M.fr.IX,18.						

C. multiarticulata; Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842)	W.A.	—	—	—	—	—	—	B.fl.VII,345	M.fr.IX,18.				
C. cymbaria; R. Brown, prodr. 221 (1810)	W.A.	—	—	V.	N.S.W.	Q.	—	B.fl.VII,345	M.fr.IX,18,58	...	P.		
OREOBOLUS; R. Brown, prodr. fl. Nov. Holl. 235 (1810).													
O. Pumilio; R. Brown, prodr. 226 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,346	M.fr.IX,20,58	...	Am., P., N.Z.		
REMIREA; Aublet, Hist. des pl. de la Guian. franc. I, 44, t. 16 (1775).													
R. maritima; Aublet, Hist. pl. Guian. I, 45, t. 16 (1775)	—	—	—	—	—	Q.	—	B.fl.VII,347	M.fr.V,92; IX,20,58.		As., Afr.. Am., P.		
RHYNCHOSPORA; Vahl, enum. plant. II, 229 (1806). (Morisia.)													
R. aurea; Vahl, enum. II, 229 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,348	M.fr.IX,17,57	...	As., Afr., Am., P.		
R. glauca; Vahl, enum. II, 229 (1806)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,349	M.fr.IX,17,57	..	As., Afr., Am.		
R. Wallichiana; Kunth, enum. II, 289 (1837)	—	—	—	—	—	Q.	N.A.	B.fl.VII,349	M.fr.IX,17,52	...	As., Afr., Am.		
R. longisetis; R. Brown, prodr. 230 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,350	M.fr.IX,17,58.				
R. tenuifolia; Bentham, Fl. Austr. VII, 350 (1878)....	—	—	—	—	—	Q.	N.A.	B.fl.VII,350					
CYATHOCHAETE; Nees in Lehm. pl. Preiss. II, 86 (1846). (Tetralepis.)													
C. clandestina; Bentham, Fl. Austr. VII, 351 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,351	M.fr.IX,40; XII,25.				
C. avenacea; Bentham, Fl. Austr. VII, 351 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,351	M.fr.IX,40.				
C. diandra; Nees in Lehm. pl. Preiss. II, 86 (1846)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,352	M.fr.IX,39; XII,25.				
CARPHA; Banks & Solander in R. Brown, prodr. 230 (1810).													
C. alpina; R. Brown, prodr. 280 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,381	M.fr.IX,39 ...	...	P., N.Z.		
SCHOENUS; Linné, coroll. gen. 2 (1737). (Chaetospora, Elynanthus, Tricostularia, Helothrix, Isoschoenus, Gymnochaete, Gymnoschoenus, Mesomelaena, Discopodium.)													
S. cruentus; F. v. M., fragm. IX, 36 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,357	M.fr.IX,37.				
S. Benthami; F. v. M., syst. Cens. of Austr. pl. 127 (1882)	W.A.	—	—	—	—	—	—	B.fl.VII,357	M.fr.IX,37.				
S. lanatus; Labillardière, Nov. Holl. pl. spec. I, 19, t. 20 (1804)	W.A.	—	—	—	—	—	—	B.fl.VII,357	M.fr.IX,37.				
S. curvifolius; Poiret, Encycl. méth. suppl. II, 251 (1811)...	W.A.	—	—	—	—	—	—	B.fl.VII,358	M.fr.IX,36.				
S. capitatus; F. v. M., fragm. IX, 58 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,358	M.fr.IX,37.				
S. setifolius; Bentham, Fl. Austr. VII, 359 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,359					
S. Drummondii; Bentham, Fl. Austr. VII, 359, non. Steud. (1870)	W.A.	—	—	—	—	—	—	B.fl.VII,359	M.fr.IX,37.				
S. turbinatus; Poiret, Encycl. méth. suppl. II, 251 (1811)	—	—	—	—	N.S.W.	—	—	B.fl.VII,359	M.fr.IX,33.				
S. barbatus; Boeckeler in Garcke, Linnaea XXXVIII, 277 (1874)...	W.A.	—	—	—	—	—	—	B.fl.VII,360	M.fr.IX,30.				
S. flavus; Boeckeler in Garcke, Linnaea XXXVIII, 278 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,360	M.fr.IX,30.				
S. brevisetis; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	—	—	—	—	—	—	B.fl.VII,360	M.fr.IX,30.				
S. Armeria; Boeckeler in Garcke, Linnaea XXXVIII, 279 (1874)...	W.A.	—	—	—	—	—	—	B.fl.VII,361	M.fr.IX,30.				
S. aphyllus; Boeckeler in Garcke, Linnaea XXXVIII, 280 (1874)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,361	M.fr.IX,28; XII,26.				
S. imberbis; R. Brown, prodr. 231 (1810)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,361	M.fr.IX,28.				
S. ericetorum; R. Brown, prodr. 231 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,362	M.fr.IX,58; XII,26.				
S. nitens; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,362	M.fr.IX,35	...	Am., N.Z.		
S. cygneus; Nees in Lehmann, pl. Preiss. II, 81 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,363	M.fr.IX,80.				
S. minutulus; F. v. M., fragm. IX, 32 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,363	M.fr.IX,32.				
S. trachycarpus; F. v. M., fragm. IX, 33 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,363	M.fr.IX,33.				
S. Tepperi; F. v. M., fragm. XI, 106 (1880)	—	S.A.	T.	—	—	—	—	—	M.fr.XI.106.				
S. nanus; F. v. M., fragm. IX, 36 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,364	M.fr.IX,28.				
S. pleiostemoneus; F. v. M., fragm. IX, 52 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,364	M.fr.IX,52.				
S. breviculmis; Bentham, Fl. Austr. VII, 364 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,364					
S. deformis; Poiret, Encycl. méth. suppl. II, 251 (1811)	—	S.A.	—	—	—	—	—	B.fl.VII,364	M.fr.IX,39.				
S. unispiculatus; F. v. M. in Benth. Fl. Austr. VII, 365 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,365					
S. obtusifolius; Boeckeler in Garcke, Linnaea XXXVIII, 281 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,366	M.fr.IX,31.				
S. grammatophyllus; F. v. M., fragm. IX, 31 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,366	M.fr.IX,31.				
S. asperocarpus; F. v. M., fragm. IX, 29 (1875)	W.A.	—	—	—	—	—	—	B.fl.VII,366	M.fr.IX,29.				

S. Moorei; Bentham, Fl. Austr. VII, 367 (1878)	—	—	—	—	N.S.W.	—	—	B.fl. VII,367				
S. villosus; R. Brown, prodr. 231 (1810)	—	—	—	—	N.S.W.	—	—	B.fl. VII,367	M.fr. IX,28.			
S. grandiflorus; F. v. M., fragm. IX, 30 (1875)	W.A.	—	—	—	—	—	—	B.fl. VII,367	M.fr. IX,30.			
S. distaus; F. v. M., syst. Cens. of Austr. pl. 127 (1882)	W.A.	—	—	—	—	—	—	—	M.fr. IX,35.			
S. calostachyus; Poiret, Encycl. méth. suppl. II, 251 (1811)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,368	M.fr. IX,35.			
S. scabripes; Bentham, Fl. Austr. VII, 368 (1878)	—	—	—	—	—	Q.	—	B.fl. VII,368				
S. multiglumis; Bentham, Fl. Austr. VII, 368 (1878)	W.A.	—	—	—	—	—	—	B.fl. VII,368				
S. efoliatus; F. v. M., fragm. IX, 32 (1875)	W.A.	—	—	—	—	—	—	B.fl. VII,369	M.fr. IX,32.			
S. acuminatus; R. Brown, prodr. 231 (1810)	W.A.	—	—	—	—	—	—	B.fl. VII,369	M.fr. IX,80.			
S. pedicellatus; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	—	—	—	—	—	—	B.fl. VII,369				
S. fascicularis; Nees in Ann. and Mag. of nat. Hist. VI, 48 (1842)	W.A.	—	—	—	—	—	—	B.fl. VII,370				
S. brevifolius; R. Brown, prodr. 231 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII,370	M.fr. IX,29,80.			
S. melanostachys; R. Brown, prodr. 231 (1810)	—	—	—	V.	N.S.W	Q.	—	B.fl. VII,370	M.fr. IX,29.			
S. sparteus; R. Brown, prodr. 231 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII,371				
S. vaginatus; F. v. M. in Benth. Fl. Austr. VII, 371 (1878)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,371				
S. falcatus; R. Brown, prodr. 232 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII,372	M.fr. IX,29.			
S. punctatus; R. Brown, prodr. 232 (1810)	—	—	—	—	—	—	N.A.	B.fl. VII,372				
S. indutus; F. v. M. in Benth. Fl. Austr. VII, 372 (1878)	W.A.	—	—	—	—	Q.	—	B.fl. VII,372				
S. bifidus; Boeckeler in Garcke, Linnaea XXXVIII, 282 (1874)	W.A.	—	—	—	—	—	—	B.fl. VII,373	M.fr. IX,38,80.			
S. apogon; Roemer et Schultes, syst. veg. II, 77 (1817)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl. VII,373	M.fr. IX,56; X,120.	N.Z.		
S. odontocarpus; F. v. M., fragm. IX, 32 (1875)	W.A.	—	—	—	—	—	—	B.fl. VII,374	M.fr. IX,32.			
S. humilis; Bentham, Fl. Austr. VII, 374 (1878)	W.A.	—	—	—	—	—	—	B.fl. VII,374				
S. sculptus; Boeckeler in Garcke, Linnaea XXXVIII, 286 (1874)	W.A.	S.A.	—	—	—	—	—	B.fl. VII,374	M.fr. IX,30.			
S. axillaris; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII,375	M.fr. IX,34	...	N.Z.	
S. tenellus; Bentham, Fl. Austr. VII, 375 (1878)	W.A.	—	—	—	—	—	—	B.fl. VII,375				
S. natans; F. v. M., fragm. IX, 36 (1875)	W.A.	—	—	—	—	—	—	B.fl. VII,375	M.fr. IX,38.			
S. fluitans; J. Hooker, Fl. Tasman. II, 81, t. 141 (1860)	—	—	T.	—	—	—	—	B.fl. VII,376	M.fr. IX,28.			
S. octandrus; F. v. M., fragm. IX, 31 (1875)	W.A.	—	—	—	—	—	—	B.fl. VII,377	M.fr. IX,31.			
S. capillaris; F. v. M., syst. Cens. of Austr. pl. 128 (1882)	—	S.A.	T.	V.	—	—	—	B.fl. VII,377	M.fr. IX,34,80.			
S. stygius; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	—	—	—	—	—	—	B.fl. VII,378	M.fr. IX,36,34,80.			
S. deustus; F. v. M., syst. Cens. of Austr. pl. 128 (1882) from R. Brown (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,379	M.fr. IX,39.			
S. tetragonus; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	—	—	—	—	—	—	B.fl. VII,380	M.fr. IX,36.			
S. sphaerocephalus; Poiret, Encycl. méth. suppl. II, 251 (1811)	—	—	T.	V.	N.S.W.	—	—	B.fl. VII,380	M.fr. IX,33.			
S. anceps; Poiret, Encycl. méth. suppl. II, 251 (1811)	W.A.	—	—	—	—	—	—	B.fl. VII,380	M.fr. IX,30.			
S. paludosus; Poiret, Encycl. méth. suppl. II, 251 (1811)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII,382	M.fr. IX,35.			
S. pauciflorus; syst. Cens. of Austr. pl. 128 (1882)	—	—	—	V.	—	—	—	B.fl. VII,383	M.fr. IX,23.			
S. Tricostularia; F. v. M., syst. Cens. of Austr. pl. 128 (1882)	W.A.	—	—	—	—	—	—	B.fl. VII,383	M.fr. IX,37.			
S. Neesii; F. v. M., syst. Cens. of Austr. pl. 128 (1882) from Lehmann (1846)	W.A.	—	—	—	—	—	—	B.fl. VII,383	M.fr. IX,37.			
S. fimbristyloides; F. v. M., syst. Cens. of Austr. pl. 128 (1882)	—	—	—	—	—	—	N.A.	B.fl. VII,384	M.fr. IX,34.			
LEPIDOSPORA; F. v. M., fragm. phytograph. Austral. IX, 34 (1875).												
L. tenuissima; F. v. M., fragm. IX, 34 (1875)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII,365	M.fr. IX,34; XII,26.			
LEPIDOSPERMA; Labillardière, Nov. Holl. pl. spec. I, 14 (1804).												
L. gladiatum; Labillardière, Nov. Holl. pl. spec. I, 15, t. 12 (1804)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl. VII,387	M.fr. XII,26.			
L. effusum; Bentham, Fl. Austr. VII, 387 (1878)	W.A.	—	—	—	—	—	—	B.fl. VII,387				
L. rupestre; Bentham, Fl. Austr. VII, 388 (1878)	W.A.	—	—	—	—	—	—	B.fl. VII,388				
L. elatius; Labillardière, Nov. Holl. pl. spec. I, 15, t. 11 (1804)	—	S.A.	T.	V.	—	—	—	B.fl. VII,388	M.fr. IX,25.			
L. tetraquetrum; Nees in Lehm. pl. Preiss. II, 90 (1846)	W.A.	—	—	—	—	—	—	B.fl. VII,388	M.fr. IX,24.			
L. Oldfieldii; J. Hooker, Fl. Tasman. II, 91, t. 146 (1860)	—	—	T.	—	—	—	—	B.fl. VII,389				
L. exaltatum; R. Brown, prodr. 234 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII,389				

L. longitudinale; Labillardière, Nov. Holl. pl. spec. I, 16, t. 13 (1804) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,389	M.fr.IX,25.		
L. concavum; R. Brown, prodr. 234 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,390	M.fr.IX,24 ...	...	N.Z.
L. angustatum; R. Brown, prodr. 235 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,391	M.fr.IX,26,80.		
L. Drummondii; Bentham, Fl. Austr. VII, 391 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,391			
L. Brunonianum; Nees in Lehm. pl. Preiss. II, 92 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,392			
L. tuberculatum; Nees in Lehm. pl. Preiss. II, 90 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,392	M.fr.IX,26.		
L. resinosum; F. v. M., fragm. IX, 26 (1875) implied ...	W.A.	—	—	—	—	—	—	B.fl.VII,392	M.fr.IX,26.		
L. viscidum; R. Brown, prodr. 234 (1810) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,393			
L. costale; Nees in Lehm. pl. Preiss. II, 92 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,393			
L. laterale; R. Brown, prodr. 234 (1810) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,393			
L. congestum; R. Brown, prodr. 234 (1810) ...	—	S.A.	—	—	—	—	—	B.fl.VII,394			
L. globosum; Labillardière, Nov. Holl. pl. spec. I, 16, t. 14 (1804) ...	—	S.A.	T.	V.	—	—	—	B.fl.VII,394			
L. lineare; R. Brown, prodr. 235 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,395	M.fr.IX,26.		
L. aphyllum; R. Brown, prodr. 235 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,395			
L. gracile; R. Brown, prodr. 235 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,395			
L. semiteres; F. v. M. in Garcke, Linnaea XXXVIII, 327 (1874)...	—	S.A.	—	V.	—	—	—	B.fl.VII,396			
L. canescens; Boeckeler in Garcke, Linnaea XXXVIII, 330 (1874) ...	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,396	M.fr.IX,24.		
L. pubisquameum; Steudel, syn. pl. glum. II, 158 (1855) ...	W.A.	—	—	—	—	—	—	B.fl.VII,396	M.fr.IX,27.		
L. scabrum; Nees in Lehm. pl. Preiss. II, 92 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,397	M.fr.IX,27.		
L. tenue; Bentham, Fl. Austr. VII, 397 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,397			
L. leptostachyum; Bentham, Fl. Austr. VII, 397 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,397			
L. leptophyllum; Bentham, Fl. Austr. VII, 398 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,398			
L. tortuosum; F. v. M., general Report 9 (1855) ...	—	—	—	V.	—	—	—	B.fl.VII,398	M.fr.IX,23.		
L. flexuosum; R. Brown, prodr. 235 (1810) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,398			
L. filiforme; Labillardière, Nov. Holl. pl. spec. I, 17, t. 15 (1804) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,398	M.fr.IX,27; XII,26.		
L. striatum; R. Brown, prodr. 235 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,399	M.fr.IX,26,27.		
L. Neesii; Kunth, enum. II, 319 (1837) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,399	M.fr.IX,24,27.		
L. carphoides; F. v. M. in Benth. Fl. Austr. VII, 400 (1878) ...	W.A.	S.A.	—	V.	—	—	—	B.fl.VII,400			

GAHNIA; R. et G. Forster, char. gen. 51 (1776). (Cladium, Lampocarya, Melachne, Epiandra, Didymonema, Baumea, Chapelliera, Morelotia.)

G. Mariscus; F. v. M., Key to Vict. pl. I, 456 (1888) from Linné (1753) ...	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,402	M.fr.IX,14,57; XII,25.	E.,As.,Afr.,Am.,P.	
G. insularis; Cladium; Bentham, Fl. Austr. 403 (1878) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,403			
G. articulata; F. v. M., Key to Vict. pl. I, 453 (1888) from R. Brown (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,403	M.fr.IX,14,57 ...	N.Z., P.	
G. arthrophylla; Cladium; F. v. M., fragm. IX, 14 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,403	M.fr.IX,14.		
G. glomerata; F. v. M., Key to Vict. pl. I, 455 (1888) from R. Brown (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,404	M.fr.IX,15,57 ...	N.Z.	
G. Preissii; Baumea; Nees in Lehm. pl. Preiss. II, 75 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,405	M.fr.IX,13,15,56.		
G. laxa; Chapelliera; Nees in Lehm. pl. Preiss. II, 76 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,405			
G. riparia; Cladium; Bentham, Fl. Austr. VII, 405 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,405			
G. teretifolia; Cladium; R. Brown, prodr. 237 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,406	M.fr.IX,15,56; XII,25.	N.Z.	
G. tetraquetra; F. v. M., Key to Vict. pl. I, 454 (1888) from J. Hooker (1860) ..	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,406	... ...	...	N.Z.
G. schoenoides; F. v. M., Key to Vict. pl. I, 454 (1888) from R. Brown (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,407	M.fr.XII,26.		
G. Gunnii; F. v. M., Key to Vict. pl. I, 455 (1888) from J. Hooker (1860) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,407	M.fr.IX,15,57.		
G. juncea; F. v. M., Key to Vict. pl. I, 453 (1888) from R. Brown (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,408	M.fr.IX,16,57 ...	N.Z.	
G. vaginalis; Cladium; Bentham, Fl. Austr. VII, 409 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,408			
G. elynanthoides; Cladium; F. v. M., fragm. IX, 31 (1875) ...	W.A.	—	—	—	—	—	—	B.fl.VII,409	M.fr.IX,31.		
G. Filum; F. v. M., Key to Vict. pl. I, 456 (1888) from Labillardière (1804) ...	—	S.A.	T.	V.	—	—	—	B.fl.VII,409	M.fr.IX,14,57; XII,26.		
G. trifida; Labillardière, Nov. Holl. pl. spec. I, 89, t. 116 (1804)...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,413	M.fr.XII,26.		
G. decomposita; Bentham, Fl. Austr. VII, 417 (1876) from R. Brown (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,417	M.fr.IX,18.		
G microstachya; Bentham, Fl. Austr. VII, 414 (1878) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,414			
G. polyphylla; Bentham, Fl. Austr. VII, 415 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,415			
G. ancistrophylla; F. v. M. in Benth. Fl. Austr. VII, 415 (1878) ...	W.A.	—	—	—	—	—	—	B.fl.VII,415			

G. lanigera; Bentham, Fl. Austr. VII, 415 (1878) from R. Brown (1810) ...	W.A.	S.A.	—	—	—	—	—	B.fl.VII,415	M.fr.IX,14,57.		
G. aristata; F. v. M. in Benth. Fl. Austr. 416 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,416			
G. deusta; Bentham, Fl. Austr. VII, 416 (1878) from R. Brown (1810)	W.A.	S.A.	—	—	—	—	—	B.fl.VII,416	M.fr.IX,14.		
G. Radula; Bentham, Fl. Austr. VII, 417 (1878) from R. Brown (1810)... ...	—	S.A.	T.	V.	—	—	—	B.fl.VII,414	M.fr.IX,13,56.		
G. Sieberi; Boeckeler in Garcke, Linn. XXXVIII, 343 (1874) from Schrader (1830)	—	—	—	—	N.S.W.	—	—	B.fl.VII,414	M.fr.IX,14,57.		
G. tetragonocarpa; Boeckeler in Linnaea XXXVIII, 347 (1874)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,418			
G. melanocarpa; R. Brown, prodr. 239 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,414	M.fr.IX,13,56; XII,26.		
G. psittacorum; Labillardière, Nov. Holl. pl. spec. I, 89, t. 115 (1804)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,418	M.fr.IX,13,56; XII,25.		
G. aspera; Sprengel, syst. veg. II, 114 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,412	M.fr.IX,12,56	...	P.
G. xanthocarpa; J. Hooker, Handb. of the N.Z. Flor. 306 (1864)...	—	—	—	—	N.S.W.	—	—	B.fl.VII,418	M.fr.IX,13,56	...	N.Z.
CAUSTIS; R. Brown, prodr. 239 (1810). (Eurostorrhiza.)											
C. pentandra; R. Brown, prodr. 240 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,420	M.fr.IX,19,58; XII,26.		
C. flexuosa; R. Brown, prodr. 239 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,421	M.fr.IX,58.		
C. recurvata; Sprengel, syst. cur. post. 26 (1827)	—	—	—	—	N.S.W.	—	—	B.fl.VII,421	M.fr.IX,19.		
C. restiacea; F. v. M. in Benth. Fl. Austr. VII, 421 (1878)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,421	M.fr.IX,19.		
C. dioica; R. Brown, prodr. 239 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,422	M.fr.IX,19,58.		
ARTHROSTYLIS; R. Brown, prodr. 229 (1810). (Arthrostyles.)											
A. aphylla; R. Brown, prodr. 229 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,423	M.fr.IX,9,55; XII,25.		
REEDIA; F. v. M., fragm. I, 240, t. 10 (1859).											
R. spathacea; F. v. M., fragm. I, 240, t. 10 (1859)	W.A.	—	—	—	—	—	—	B.fl.VII,423	M.fr.IX,15.		
EUANDRA; R. Brown, prodr. 239 (1810). (Evandra.)											
E. aristata; R. Brown, prodr. 239 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,424	M.fr.IX,18.		
E. pauciflora; R. Brown, prodr. 239 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,425	M.fr.IX,18.		
SCLERIA; Bergius in Svensk. Vet. Acad. Handl. 149, t. 4 (1765). (Diplacrum, Hypoporum, Sphaeropus.)											
S. caricina; Bentham, Fl. Austr. VII (1878)	—	—	—	—	—	Q.	—	B.fl.VII,426	M.fr.IX,58...	...	As.
S. pygmaea; R. Brown, prodr. 240 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,427	M.fr.IX,22,58.		
S. rugosa; R. Brown, prodr. 240 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,428	M.fr.IX,22,58.		
S. laxa; R. Brown, prodr. 240 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,428	M.fr.IX,21.		
S. Brownii; Kunth, enum. II, 349 (1837)	—	—	—	—	—	Q.	N.A.	B.fl.VII,429	M.fr.IX,21,58.		
S. lithosperma; Willdenow, spec. plant. IV, 316 (1805)	—	—	—	—	—	Q.	N.A.	B.fl.VII,429	M.fr.IX,21,58	...	As., P.
S. tessellata; Willdenow, spec. plant. IV, 315 (1805)	—	—	—	—	—	Q.	—	B.fl.VII,430	... ...	...	As.
S. margaritifera; Willdenow, spec. plant. IV, 312 (1805)	—	—	—	—	—	Q.	—	B.fl.VII,430	M.fr.IX,21,58	...	As., P.
S. Graeffeana; Boeckeler in Regensb. Flora 121 (1875)	—	—	—	—	—	Q.	N.A.	B.fl.VII,431			
S. hebecarpa; Nees in Wight, Contrib. to Bot. of Ind. 117 (1834)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,431	M.fr.IX,21,58	...	As.
S. Chinensis; Kunth, enum. II, 357 (1837)	—	—	—	—	—	Q.	—	B.fl.VII,431	M.fr.IX,20,58	...	As.
S. oryzoides; Presl, reliq. Haenk. I, 201 (1830)	—	—	—	—	—	—	N.A.	B.fl.VII,432	... ...	...	As., P.
S. sphacelata; F. v. M., fragm. IX, 20 (1875)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,432	M.fr.IX,20.		
UNCINIA; Persoon, synops. plant. I, 534 (1807).											
U. tenella; R. Brown, prodr. 241 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,433	M.fr.VIII,151.		
U. compacta; R. Brown, prodr. 241 (1810)	—	—	T.	V.	—	—	—	B.fl.VII,434	M.fr.VIII,152.		
U. riparia; R. Brown, prodr. 241 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,434	M.fr.VIII,152.		
U. debilior; F. v. M., fragm. VIII, 151 (1874)	—	—	—	—	N.S.W.	—	—	B.fl.VII,435	M.fr.VIII,151.		
CAREX; Ruppius, Fl. Jenens. 306 (1718).											
C. cephalotes; F. v. M. in Transact. phil. Soc. Vict. I, 110 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,437	M.fr.VIII,251.		
C. acicularis; Boott in J. Hook. Fl. N. Zel. I, 280, t. 63 (1853)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,437	M.fr.VIII,251	...	N.Z.
C capillacea; Boott, Illustr. of the gen. Carex I, 44, t. 110 (1858)	—	—	—	—	N.S.W.	—	—	B.fl.VII,438	M.fr.IX,191	...	As.
C. inversa; R. Brown, prodr. 242 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,438	M.fr.VIII,252; XII,26.		N.Z.

C. canescens; Linné, spec. plant. 974 (1753) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,439	M.fr.VIII,253	...	E., As., Am.
C. echinata; Murray, prodr. design. stirp. Goett. 76 (1770) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,439	M.fr.VIII,252	...	E., As., Am.
C. hypandra; F. v. M., fragm. VIII, 259 (1874) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,439	M.fr.VIII,259.		
C. chlorantha; R. Brown, prodr. 242 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,440	M.fr.VIII,256.		
C. paniculata; Linné, amoen. acad. IV, 294 (1759) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,440	M.fr.VIII,256	...	E., As., N.Z.
C. declinata; Boott, Ill. of the gen. Carex IV, 17, t. 580 (1867) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,441	M.fr.VIII,257; XII,26.		
C. tereticaulis; F. v. M., fragm. VIII, 256 (1874) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,441	M.fr.VIII,256; XII,26.		
C. haematostoma; Nees in Wight's contrib. to the Bot. of Ind. 125 (1834) ...	—	—	T.	—	—	—	—	—	...	...	As.
C. fissilis; Boott, Ill. of the gen. Carex. II, 86, t. 245 (1860) ...	—	—	—	—	—	Q.	—	B.fl.VII,441	M.fr.VIII,249; IX,80.		P.
C. brunnea; Thunberg, fl. Japon. 38 (1784) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,442	M.fr.VIII,250	...	As., Afr., P.
C. contracta; F. v. M., fragm. VIII, 258 (1874) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,442	M.fr.VIII,258.		
C. caespitosa; Linné, spec. plant. 978 (1853) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,442	M.fr.VIII,257	...	E., As., Afr., Am.
C. acuta; Linné, spec. plant. 978 (1753) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,443	M.fr.VIII,259; XII,26.		E., As., Am.
C. lobolepis; F. v. M., fragm. VIII, 258 (1874) ...	—	—	—	—	N.S.W.	—	—	B.fl.VII,443	M.fr.VIII,258.		
C. flava; Linné, spec. plant. 975 (1753) ...	—	—	T.	—	—	—	—	B.fl.VII,444	M.fr.VIII,250	...	E., As., Afr., Am.
C. Buxbaumii; Wahlenberg in Act. Holmiens. 139 (1803) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,444	M.fr.VIII,252	...	E., As., Am.
C. pumila; Thunberg, Fl. Japon. 38 (1784) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,445	M.fr.VIII,251; XII,26.		As., Am., N.Z.
C. breviculmis; R. Brown, prodr. 242 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,445	M.fr.VIII,255; XII,26.		As., N.Z.
C. Neesiana; Endlicher, prodr. fl. ins. Norf. (1837) ...	—	—	—	—	N.S.W.	—	—	—	M.fr.VIII,259.		
C. Preissii; Nees in Lehm. pl. Preiss. II, 94 (1846) ...	W.A.	—	—	—	—	—	—	B.fl.VII,446	M.fr.VIII,251.		
C. Gunnisna; Boott in Transact. Linn. Soc. XX, 143 (1851) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,446	M.fr.VIII,251.		
C. Bichenoviana; Boott in J. Hook. Fl. Tasman. II, 101 (1860) ...	—	—	T.	—	—	—	—	B.fl.VII,446			
C. maculata; Boott in Transact. Linn. Soc. XX, 128 (1851) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,447	M.fr.VIII,258	...	As.
C. lacistoma; R. Brown, prodr. 242 (1810) ...	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,447	M.fr.IX,250; XII,26.		
C. alsophila; F. v. M., fragm. VIII, 257 (1874) ...	—	—	—	V.	—	—	—	B.fl.VII,447	M.fr.VIII,257.		
C. longibrachiata; Boeckeler in Garcke, Linnaea XXXVIII, 288 (1881) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,448	M.fr.VIII,250.		
C. Pseudo-Cyperus; Linné, spec. plant. 978 (1753) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,448	M.fr.VIII,249		E., As., Afr., Am., N.Z.

GRAMINEAE.

HALLER, enum. stirp. Helv. I, 203 (1742).—Genera 76, Species 345.

ERIOCHLOA; Humboldt, Bonpland & Kunth, nov. gen. et sp. pl. I, 94, t. 30 (1815). (Helopus.)

E. polystachya; Humboldt et Kunth, nov. gen. I, 95, t. 31 (1815) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,462	M.fr.VI,85; VIII,126.		As., P.

PASPALUM; Linné, syst. nat. ed. decim. 855 (1759).

P. scrobiculatum; Linné, mantiss. 29 (1767) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,460	M.fr.VIII,156	...	As., Afr., P.
P. distichum; Linné, syst. nat. ed. decim. 855 (1759) ...	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.VII,460	M.fr.VIII,156	...	As., Afr., Am.
P. brevifolium; Fluegge, gram. monogr. 150 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,461	M.fr.VIII,195	...	As.
P. minutiflorum; Steudel, sym. glum. I, 17 (1855) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,461	...	...	As.

PANICUM; Tournefort, inst. 515, t. 298 (1700) from Plinius. (Isachne, Digitaria, Echinochloa, Coridochloa, Chamaeraphis, Paractaenum.)

P. coenicolum; F. v. M. in Transact. Vict. Inst. 45 (1854) ...	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,467			
P. divaricatissimum; R. Brown, prodr. 192 (1810) ...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,467	M.fr.VIII,154; X,76.		
P. macractinium; Bentham, Fl. Austr. VII, 468 (1878) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,468			
P. papposum; R. Brown, prodr. 192 (1810) ...	—	—	—	—	—	—	N.A.	B.fl.VII,468	M.fr.VIII,155,196.		
P. sanguinale; Linné, spec. plant. 57 (1753) ...	W.A.	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,469	M.fr.VIII,153	...	As., Afr., Am., P.
P. ctenanthum; F. v. M., fragm. VIII, 153 (1874) ...	—	—	—	—	—	—	N.A.	B.fl.VII,469	M.fr.VIII,153.		
P. stenostachyum; Bentham, Fl. Austr. VII, 470 (1878) ...	—	—	—	—	—	—	N.A.	B.fl.VII,470			
P. tenuissimum; Bentham, Fl. Austr. VII, 470 (1878) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,470			
P. parviflorum; R. Brown, prodr. 192 (1810) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,470	M.fr.VIII,196; XI,129.		
P. Baileyi; Bentham, Fl. Austr. VII, 470 (1878) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,471			

P. gibbosum; R. Brown, prodr. 193 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 471	M.fr. VIII, 155; XI, 129.	
P. leucophaeum; Humboldt & Kunth, nov. gen. et spec. I, 97 (1815)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 472	M.fr. VIII, 155	Afr., Am.
P. semialatum; R. Brown, prodr. 192 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 472	M.fr. VIII, 196	As.
P. rarum; R. Brown, prodr. 189 (1810)	—	—	—	—	—	—	N.A.	B.fl. VII, 473		
P. argenteum; R. Brown, prodr. 190 (1810)	—	S.A.	—	—	—	—	N.A.	B.fl. VII, 473		
P. holosericeum; R. Brown, prodr. 190 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 473	M.fr. VIII, 155.	
P. flavidum; Retzius, observ. IV, 15 (1787)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 474	M.fr. VIII, 189	As.
P. gracile; R. Brown, prodr. 190 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 475	M.fr. VIII, 189.	
P. tubulatum; Haeckel in Engler's Jahrb. VI, 234 (1885)	—	—	—	—	—	—	N.A.	—		
P. prostratum; Lamarck, illustr. I, 171 (1791)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 476	M.fr. VIII, 191	As., Afr., Am.
P. helopus; Trinius in Spreng. Neue Entd. II, 84 (1821)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 476	... ...	As., Afr.
P. Gilesii; Bentham, Fl. Austr. VII, 477 (1878)	—	S.A.	—	—	—	Q.	N.A.	B.fl. VII, 477	M.fr. XI, 129.	
P. piligerum; F. v. M. in Benth. Fl. Austr. VII, 477 (1878)	—	—	—	—	—	—	N.A.	B.fl. VII, 477		
P. polyphyllum; R. Brown, prodr. 190 (1810)	—	—	—	—	—	—	N.A.	B.fl. VII, 477	M.fr. VIII, 194.	
P. distachyon; Linné, mantiss. alt. pl. 138 (1771)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 478	M.fr. VIII, 194	As.
P. reversum; F. v. M., fragm. VIII, 152 (1874)	W.A.	S.A.	—	—	N.S.W.	Q.	—	B.fl. VII, 478	M.fr. X, 76.	
P. colonum; Linné, syst. nat. ed. decim. 5 (1758)	—	—	—	—	—	—	N.A.	B.fl. VII, 478	... ...	E., As., Afr., Am.
P. Crus Galli; Linné, spec. plant. 56 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 479	M.fr. VIII, 198	As., Afr., Am.
P. myosuroides; R. Brown, prodr. 189 (1810)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 480	... ...	As., Afr.
P. Indicum; Linné, mantiss. pl. alt. 184 (1771)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 480	M.fr. VIII, 197; XI, 129.	As., Afr.
P. Myurus; Lamarck, illustr. I, 172 (1791)	—	—	—	—	—	Q.	—	B.fl. VII, 480	M.fr. X, 114 ...	As., Afr., Am.
P. foliosum; R. Brown, prodr. 191 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 481	M.fr. VIII, 194.	
P. adspersum; Trinius, spec. gram. t. 169 (1830)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl. VII, 481	... ...	Am.
P. inaequale; F. v. M., fragm. VIII, 189 (1874)	—	—	—	—	—	Q.	—	B.fl. VII, 482	M.fr. XI, 129.	
P. uncinulatum; R. Brown, prodr. 191 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 482	M.fr. VIII, 195; X, 76; XI, 129.	
P. majusculum; F. v. M. in Benth. Fl. Austr. VII, 482 (1878)	—	—	—	—	—	—	N.A.	B.fl. VII, 482		
P. pauciflorum; R. Brown, prodr. 483 (1810)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl. VII, 483	M.fr. VIII, 193.	
P. semitonsum; F. v. M. in Benth. Fl. Austr. VII, 483 (1878)	—	—	—	—	—	—	N.A.	B.fl. VII, 483		
P. antidotale; Retzius, observ. bot. IV, 17 (1787)	—	—	—	—	—	—	N.A.	B.fl. VII, 483		
P. repens; Linné, spec. pl. ed. sec. 87 (1762)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 484	... ...	E., As., Afr., Am., P.
P. Buncei; F. v. M. in Benth. Fl. Austr. VII, 487 (1878)	—	—	—	—	—	Q.	—	B.fl. VII, 487		
P. capillipes; Bentham, Fl. Austr. VII, 484 (1878)	—	—	—	—	—	—	N.A.	B.fl. VII, 484	M.fr. VIII, 193.	
P. pygmaeum; R. Brown, prodr. 191 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 484	M.fr. VIII, 193.	
P. brevifolium; Linné, spec. plant. 59 (1753)	—	—	—	—	—	Q.	N.A.	B.fl. VII, 485	M.fr. VIII, 193; XI, 129.	
P. hermaphroditum; Steudel, syn. glum. I, 67 (1855)	—	—	—	—	—	Q.	—	B.fl. VII, 485	... ...	As.
P. marginatum; R. Brown, prodr. 190 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII, 485	M.fr. VIII, 190.	
P. lachnophyllum; Bentham, Fl. Austr. VII, 486 (1878)	—	—	—	—	—	Q.	—	B.fl. VII, 486		
P. obseptum; Trinius, gram. panic. diss. alt. 149 (1826)	—	—	—	—	N.S.W.	—	—	B.fl. VII, 486		
P. bicolor; R. Brown, prodr. 191 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl. VII, 487		
P. melananthum; F. v. M. in Transact. Vict. Inst. 47 (1854)	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII, 488	M.fr. VIII, 192.	
P. effusum; R. Brown, prodr. 191 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 488	M.fr. VIII, 191.	
P. Mitchelli; Bentham, Fl. Austr. VII, 489 (1878)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 489	M.fr. X, 76.	
P. decompositum; R. Brown, prodr. 191 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 489	M.fr. VI, 86; VIII, 191.	As.
P. trachyrhachis; Bentham, Fl. Austr. VII, 490 (1878)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl. VII, 490	M.fr. VIII, 192; XI, 129.	
P. prolutum; F. v. M. in Transact. Vict. Inst. 46 (1856)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl. VII, 490	M.fr. VIII, 192.	
P. spinescens; R. Brown, prodr. 193 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl. VII, 499	M.fr. VIII, 197; XI, 129.	As.
P. paradoxum; R. Brown, prodr. 193 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII, 499	M.fr. VIII, 197.	
P. Chamaeraphis; Trinius in Mem. acad. Petersb. ser. 6, III, 217 (1835)	—	—	—	—	—	—	N.A.	B.fl. VII, 500		
P. atro-virens; Trinius in Spreng. neue Entdeckung. II, 89 (1821)	—	—	—	V.	N.S.W.	Q.	—	B.fl. VII, 625	M.fr. VIII, 193; XI, 129.	As.
P. Myosotis; Steudel, syn. glum. I, 96 (1855)	—	—	—	—	—	Q.	—	B.fl. VII, 625	M.fr. VIII, 193	As.

OPLISMENUS ; Palisot, Flore d'Oware et de Benin. II, 14 (1807). (Orthopogon.)

O. compositus ; Palisot, Agrostogr. 54 (1812) ... ... — — — V. N.S.W. Q. — B.fl.VII,491 M.fr.VIII,199 ... As., P., N.Z.

SETARIA ; Palisot, Essai d'une nouv. Agrostogr. 51, t. 13 (1812).

S. glauca ; Palisot, Agrostogr. 51, t. 13 (1812) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,492 M.fr.VIII,110. E., As., Afr., Am., P.
S. macrostachya ; Humboldt & Kunth, nov. gen. et spec. I, 110 (1815) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,493 M.fr.VIII,110; XI,129. As., Am.
S. viridis ; Palisot, Agrostogr. 51 (1812) ... ... W.A. S.A. — — — — N.A. B.fl.VII,494 ... ... E., As., Afr.
S. verticillata ; Palisot, Agrostogr. 51 (1812) ... ... W.A. — — — — — N.A. B.fl.VII,494 M.fr.VIII,110 ... E., As., Afr.

PENNISETUM ; L. C. Richard in Persoon, synops. plant. I, 72 (1805). (Gymnothrix, Plagiosetum.)

P. compressum ; R. Brown, prodr. 195 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,495 M.fr.VIII,109. ... As., P.
P. Arnhemicum ; F. v. M., fragm. VIII, 109 (1874) ... ... — — — — — — N.A. B.fl.VII,496 M.fr.VIII,109.
P. refractum ; F. v. M., fragm. VIII, 109 (1874) ... ... — S.A. — — — Q. N.A. B.fl.VII,495 M.fr.III,147; IX,194.

CENCHRUS ; Linné, coroll. gen. 20 (1737).

C. Australis ; R. Brown, prodr. 196 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,497 M.fr.VIII,107; XI,129.
C. inflexus ; R. Brown, prodr. 195 (1810) ... ... — — — — — — N.A. B.fl.VII,497 M.fr.VIII,107.
C. elymoides ; F. v. M., fragm. VIII, 107 (1874) ... ... — — — — — Q. N.A. B.fl.VII,497 M.fr.VIII,107.

XEROCHLOA ; R. Brown, prodr. fl. Nov. Holl. 196 (1810).

X. imberbis ; R. Brown, prodr. 197 (1810) ... ... — — — — — Q. N.A. B.fl.VII,501 M.fr.VIII,117.
X. barbata ; R. Brown, prodr. 197 (1810) ... ... — — — — — Q. N.A. B.fl.VII,501 M.fr.VIII,117.
X. laniflora ; Bentham, Fl. Austr. VII, 502 (1878) ... ... — — — — — — N.A. B.fl.VII,502

STENOTAPHRUM ; Trinius, fundam. Agrostogr. 175 (1820). (Diastemanthe.)

S. dimidiatum ; Trinius, fund. Agrost. 175 (1820) ... ... — — — — — Q. N.A. B.fl.VII,500 M.fr.VIII,156 ... As., Afr., Am.

THUAREA ; Persoon, synops. plant. I, 110 (1805). (Thouarsea.)

T. sarmentosa ; Persoon, synops. I, 110 (1805) ... ... — — — — — Q. N.A. B.fl.VII,502 M.fr.VI,86 ... As., Afr., P.

SPINIFEX ; Linné, syst. nat. ed. XII, 757 (1767).

S. hirsutus ; Labillardière, Nov. Holl. pl. spec. II, 81, t. 230 (1806) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,503 M.fr.VIII,138 ... N.Z., P.
S. longifolius ; R. Brown, prodr. 198 (1810) ... ... W.A. — — — — Q. N.A. B.fl.VII,504 M.fr.VIII,139.
S. paradoxus ; Bentham in Hook. icon. plant. t. 1243 (1877) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,504 M.fr.VI,86; VIII,117,199; IX,195.

LEPTASPIS ; R. Brown, prodr. fl. Nov. Holl. 211 (1810).

L. Banksii ; R. Brown, prodr. 211 (1810) ... ... — — — — — Q. — B.fl.VII,548 M.fr.VIII,116. ... P.

PEROTIS ; Aiton, hort. Kew. I, 85 (1789). (Xystidium.)

P. rara ; R. Brown, prodr. 172 (1810) ... ... — S.A. — — N.S.W. Q. N.A. B.fl.VII,509 M.fr.VIII,115 ... As., P.

POLYPOGON ; Desfontaines, flor. Atlantic. I, 66 (1798).

P. fugax ; Nees in Steudel, syn. glum. I, 184 (1855) ... ... W.A. — — — — — — B.fl.VII,547 .. .. As.
P. tenellus ; R. Brown, prodr. 173 (1810) ... ... W.A. — — — — — — B.fl.VII,547 M.fr.VIII,114.

ARUNDINELLA ; Raddi, Agrostogr. Brasil. 37 (1823).

A. Nepalensis ; Trinius, spec. gram. t. 268 (1833) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,545 M.fr.VIII,139 ... As., Afr.
A. Schultzii ; Bentham, Fl. Austr. VII, 545 (1878) ... ... — — — — — — N.A. B.fl.VII,545

TRAGUS ; Haller, hist. stirp. Helvet. n. 1413 (1768). (Lappago.)

T. racemosus ; Haller, stirp. Helv. n. 1413 (1768) ... ... — S.A. — V. N.S.W. Q. N.A. B.fl.VII,507 M.fr.VIII,107 ... E., As., Afr., Am.

NEURACHNE ; R. Brown, prodr. fl. Nov. Holl. 196 (1810).

N. alopecuroides ; R. Brown, prodr. 196 (1810) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,507 M.fr.VI,86; VIII,200; XI,129.
N. Mitchelliana ; Nees in Hook. Lond. Journ. of Bot. II, 410 (1843) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,508 M.fr.VI,86; VIII,200.
N. Munroi ; F. v. M., fragm. VIII, 200 (1874) ... ... — S.A. — V. N.S.W. — — B.fl.VII,508 M.fr.V,204; XI,129.

ZOYSIA; Willdenow, Neue Schrift. nat. Freunde zu Berl. III, 440 (1801).

Z. pungens; Willdenow, Neue Schrift. nat. Fr. Berl. III, 440 (1801)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,506	M.fr.VIII,116	...	As., N.Z.

IMPERATA; Cyrillo, plant. rar. Neap. II, t. 11 (1792).

I. arundinacea; Cyrillo, plant. rar. Neap. II, t. 11 (1792)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,536	M.fr.VIII,126	...	E., As., Afr., Am., P.

ERIANTHUS; L. C. Richard in Michaux, fl. bor. Amer. I, 54 (1803). (Pollinia, Eulalia, Pogonatherum partly, Leptatherum, Saccharum partly.)

E. articulatus; F. v. M., fragm. VIII, 118 (1874)	—	—	—	—	—	Q.	N.A.	B.fl.VII,525	M.fr.X,76	...	As.
E. irritans; Kunth, Graminées I, 160 (1832)	—	—	—	—	—	Q.	N.A.	B.fl.VII,526	M.fr.VIII,118.		
E. fulvus; Kunth, Graminées I, 160 (1832)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,526	M.fr.VIII,118.		
E. Roxburghii; F. v. M., fragm. VIII, 117 (1874)	—	—	—	—	—	Q.	—	B.fl.VII,527	M.fr.VIII,117	...	As.
E. Mackinlayi; F. v. M., Cens. 131 (1882)	—	—	—	—	—	—	N.A.	B.fl.VII,527	M.fr.VIII,118.		

POGONATHERUM; Palisot, Agrostogr. 176 (1812).

P. saccharoideum; Palisot, Agrostogr. 56, t. 11 (1812)	—	—	—	—	—	Q.	—	—	...	...	...	As.

DIMERIA; R. Brown, prodr. fl. Nov. Holl. 204 (1810).

D. acinaciformis; R. Brown, prodr. 204 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,523				
D. ornithopoda; Trinius, fundam. agrostogr. 167 (1820)	—	—	—	—	—	Q.	N.A.	B.fl.VII,523	M.fr.VII,104; XI,121.		As.

ARTHRAXON; Palisot, Agrostogr. III, t. 11, f. 6 (1812). (Batratherum.)

A. ciliaris; Palisot, Agrostogr. III, t. 11 (1812)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,524	M.fr.VIII,119; X,76.		As.

ELIONURUS; Humboldt & Bonpland in Willd. sp. pl. IV, 941 (1805).

E. citreus; Munro in Benth. Fl. Austr. VII, 510 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,510	M.fr.XI,129.		

LEPTURUS; R. Brown, prodr. fl. Nov. Holl. 207 (1810). (Monerma.)

L. incurvatus; Trinius, fundam. Agrostogr. 123 (1820)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,668	M.fr.VIII,117	...	E., As., Afr., N.Z.
L. cylindricus; Trinius, fundam. Agrostogr. 123 (1820)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,668	...	...	E., As., Afr.
L. repens; R. Brown, prodr. 207 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,668	M.fr.VI,86	...	As., P.

HEMARTHRIA; R. Brown, prodr. fl. Nov. Holl. 207 (1810). (Rottboelliae subgenus.)

H. compressa; R. Brown, prodr. 207 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,511	M.fr.XI,129.		

ROTTBOELLIA; Linné, fil. gram. gen. 22 (1779). (Coelorrhachis.)

R. formosa; R. Brown, prodr. 206 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,513	M.fr.VI,85; XI,129.		
R. exaltata; Linné, fil. suppl. 114 (1781)	—	—	—	—	—	Q.	N.A.	B.fl.VII,513	...	...	As.
R. ophiuroides; Bentham, Fl. Austr. VII, 514 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.VII,514	M.fr.VIII,123	...	As., P.

OPHIUROS; K. F. Gaertner, de fruct. III, 3, t. 181 (1805). (Ophiurus.)

O. corymbosus; K. F. Gaertner, de fruct. III, 3, t. 181 (1805)	—	—	—	—	—	Q.	N.A.	B.fl.VII,512	M.fr.VI,85	...	As., Afr.

MANISURIS; Linné, syst. nat. ed. duodec. 762 (1767).

M. granularis; Swartz, Fl. Ind. occid. I, 186 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.VII,514	M.fr.VII,116; XI,129.		As., Afr., Am.

ISCHAEMUM; Linné, syst. nat. 8 (1735); Linné, gen. pl. ed. sec. 525 (1742). (Hologamium, Spodiopogon partly.)

I. truncatiglume; F. v. M. in Benth. Fl. Austr. VII, 518 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,518				
I. arundinaceum; F. v. M. in Benth. Fl. Austr. VII, 519 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,519				
I. triticeum; R. Brown, prodr. 205 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,519				
I. Australe; R. Brown, prodr. 205 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,519				
I. muticum; Linné, spec. plant. 1049 (1753)	—	—	—	—	—	Q.	—	B.fl.VII,520	M.fr.VIII,120	...	As., P.
I. ciliare; Retzius, observ. bot. VI, 36 (1791)	—	—	—	—	N.S.W.	—	—	B.fl.VII,520	M.fr.VIII,119	...	As.
I. decumbens; Bentham, Fl. Austr. VII, 521 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,521			
I. pectinatum; Trinius in Mem. de l'Acad. de Petersb. six. sèr. II, 296 (1833)...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,521	M.fr.VIII,118	...	As.
I. fragile; R. Brown, prodr. 205 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,522	M.fr VIII,123; XI,130.		
I. laxum; R. Brown, prodr. 205 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,522	M.fr.VIII,119	...	As., Afr.

ANDROPOGON; Royen. prodr. exhib. pl. hort. acad. Lugd. 52 (1740). (Sorghum, Chrysopogon, Heteropogon, Schizachyrium, Trachypogon, Holcus partly, Spodiopogon partly.)

A. erianthoides; F. v. M., fragm. X, 75 (1876)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,529	M.fr.X,75.				
A. sericens; R. Brown, prodr. 201 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,529	...	..	...	As., P.	
A. affinis; R. Brown, prodr. 201 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,530					
A. pertusus; Willdenow, spec. plant. IV, 922 (1805)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,530	M.fr.VIII,122; X,76.			As.	
A. annulatus; Forskael, Fl. Aegypt. Arab. 173 (1775)	—	S.A.	—	V.	—	Q.	N.A.	B.fl.VII,531	M.fr.VIII,123	...		As., Afr.	
A. Ischaemum; Linné, spec. plant. 1047 (1753)	W.A.	—	—	—	—	—	—	B.fl.VII,531	M.fr.VIII,122	...		E., As., Afr.	
A. punctatus; Roxburgh, fl. Indic. I, 268 (1820)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,531	..	...	...	As.	
A. procerus; R. Brown, prodr. 202 (1810)	W.A.	—	—	—	—	—	N.A.	B.fl.VII,532	M.fr.VIII,124.				
A. exaltatus; R. Brown, prodr. 202 (1810)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.VII,532	M.fr.VIII,124.				
A. bombycinus; R. Brown, prodr. 202 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,533	M.fr.XI,20.				
A. Schoenanthus; Linné, mater. med. 31 (1749)	—	—	—	—	—	Q.	—	B.fl.VII,534	M.fr.VIII,124	...		As., P.	
A. refractus; R. Brown, prodr. 202 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,534	M.fr.VIII,124; X,76.			P.	
A. filipendulinus; Hochstetter in Regensb. Fl. 175 (1846)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,534	...	...	...	Afr.	
A. brevifolius; Swartz, fl. Ind. occid. I, 209 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.VII,535					
A. contortus; Linné, spec. plant. 1045 (1753)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,517	M.fr.VIII,120	...		As., Afr., Am., P.	
A. triticeus; R. Brown, prodr. 201 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,517	M.fr.VIII,120	...		As.	
A. Gryllus; Linné, amoen. acad. IV, 332 (1759)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,537	M.fr.VIII,121; XI,130.			As., Afr., P.	
A. micranthus; Kunth. gram. I, 165 (1835)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,538	M.fr.VIII,122; X,76; XI,130.			As., P.	
A. aciculatus; Retzius, observ. bot. V, 22 (1789)	—	—	—	—	—	Q.	N.A.	B.fl.VII,538	M.fr.VIII,122	...		As., P.	
A. elongatus; Sprengel, syst. I, 289 (1825)	—	—	—	—	—	Q.	N.A.	B.fl.VII,539	M.fr.VIII,121.				
A. Halepensis; Sibthorp & Smith, Fl. Graeca I, 52, t. 68 (1806)	W.A.	—	—	—	N.S.W.	Q.	—	B.fl.VII,540	M.fr.VIII,119	...		E., As., P.	
A. Australis; Sprengel. syst. I, 287 (1825)	W.A.	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,540					
A. serratus; Thunberg, fl. Japon. 41 (1784)	—	—	—	—	—	Q.	N.A.	B.fl.VII,541	...	...	...	As.	
A. intrans; F. v. M., Cens. of Austr. pl. 132 (1882)	—	—	—	—	—	—	N.A.	B.fl.VII,541					

ANTHISTIRIA; Linné, fil. gram. gen. 35 (1779). (Themeda 1775, Iseilema.)

A. ciliata; Linné, fil. gram. gen. 35 (1779)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,542	M.fr.V,207; VIII,139.			As., Afr.	
A. frondosa; R. Brown, prodr. 200 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,542	M.fr.VIII,139; XI,130.				
A. avenacea; F. v. M., fragm. V, 206 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,543	M.fr.VI,253; VIII,139.				
A. membranacea; Lindley in Mitch. Trop. Austr. 88 (1848)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,543	M.fr.V,207; VIII,139.				

APLUDA; Linné, spec. plant. 82 (1753).

A. mutica; Linné, spec. plant. 82 (1753)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,544	M.fr.VIII,140	...		As., Afr., P.	

CHIONACHNE; R. Brown in Horsfield's pl. Jav. rar. 15 (1838).

C. barbata; R. Brown in Bennett, pl. Javan. rar. 18 (1838)	—	—	—	—	—	Q.	—	B.fl.VII,515	...	...	...	As.	
C. cyathopoda; F. v. M. in Benth. Fl. Austr. VII, 516 (1878)	—	—	—	—	—	Q.	N.A.	B.fl.VII,515	M.fr.VIII,116.				

ALOPECURUS; Linné, syst. nat. 8 (1735); gen. plant. 18 (1737). (Perhaps immigrated.)

A. geniculatus; Linné, spec. plant. 60 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,555	M.fr.VIII,138	...		As., Am., N.Z.	

LEERSIA; Solander in Swartz, nov. gen. et spec. plant. 1 et 21 (1788). (Asprella, Asperella.)

L. hexandra; Swartz, nov. gen. et spec. pl. 21 (1788)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,549	M.fr.VIII,115	...		E., As., Afr., Am.	

ORYZA; Tournefort, inst. 513, t. 296 (1700), from Theophrastos & Dioscorides.

O. sativa; Linné, spec. plant. 333 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,550	M.fr.VIII,115; XI,130.			As.	

POTAMOPHILA; R. Brown, prodr. fl. Nov. Holl. 211 (1810).

P. parviflora; R. Brown, prodr. 211 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,550	M.fr.VIII,126.				

EHRHARTA; Thunberg in Svensk. Vetensk. Acad. Handl. 216, t. 8 (1779). (Tetrarrhena, Microlaena, Diplax.)

E. distichophylla; Labillardière, Nov. Holl. pl. spec. I, 90, t. 117 (1804)	—	—	T.	V.	—	—	—	B.fl.VII,554	M.fr.VII,90.				
E. juncea; Sprengel, syst. II, 114 (1825)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,554	M.fr.VII,90.				

E. laevis; Sprengel, syst. II, 115 (1825)	W.A.	—	—	—	—	—	—	B.fl.VII,554	M.fr.VII,90.					
E. acuminata; Sprengel, syst. II, 114 (1825)	—	—	T.	V.	—	—	—	B.fl.VII,555	M.fr.VII,90; XI,130.					
E. stipoides; Labillardière, Nov. Holl. pl. spec. I, 91, t. 118 (1804)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,552	M.fr.VII,90	...	N.Z.			
E. Diarrhena; F. v. M., fragm. VII, 89 (1870)	—	—	T.	—	—	—	—	B.fl.VII,553	M.fr.VII,89.					
HIEROCHLOE; J. G. Gmelin, Fl. Sibiric. I, 100 (1747). (Hierochloa, Disarrhenum.)														
H. redolens; R. Brown, prodr. 209 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,558	M.fr.VIII,137	...	Am., N.Z.			
H. rariflora; J. Hooker, Fl. Antarct. I, 93 (1844)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,559	M.fr.VIII,138.					
ARISTIDA; Linné, spec. plant. 82 (1753). (Chaetaria, Arthratherum.)														
A. hygrometrica; R. Brown, prodr. 174 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,561						
A. stipoides; R. Brown, prodr. 174 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,561	M.fr.VIII,111.					
A. arenaria; Gaudichaud in Freyc. voy. Bot. 407 (1826)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,561						
A. Behriana; F. v. M. in Transact. Vict. Inst. 44 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,562	M.fr.XI,130.					
A. leptopoda; Bentham, Fl. Austr. VII, 562 (1878)...	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,562	M.fr.XI,130.					
A. vagans; Cavanilles, icon. et descr. pl. V, 45, t. 471 (1799)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,562	M.fr.VIII,111.					
A. ramosa; R. Brown, prodr. 173 (1810)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,563						
A. calycina; R. Brown, prodr. 173 (1810)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,563	M.fr.XI,130.					
A. depressa; Retzius, observ. bot. IV, 22 (1787)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,563	...	...	...	As., Afr.		
STIPA; Linné, spec. plant. 78 (1737). (Streptachne, Dichelachne, Urachne partly, Muehlenbergia partly.)														
S. elegantissima; Labillardière, Nov. Holl. pl. spec. I, 23, t. 29 (1804)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,565	M.fr.VIII,103.					
S. Tuckeri; F. v. M., fragm. XI, 128 (1881)	—	—	—	V.	N.S.W.	—	—	—	M.fr.XI,129.					
S. flavescens; Labillardière, Nov. Holl. pl. spec. I, 24, t. 30 (1804)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,566	M.fr.VIII,104.					
S. teretifolia; Steudel, syn. glum. I, 128 (1855)	W.A.	S.A.	T.	V.	—	—	—	B.fl.VII,567	M.fr.VIII,104	...	N.Z.			
S. compressa; R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,567						
S. Drummondii; Steudel, syn. glum. I, 128 (1855)	W.A.	—	—	—	—	—	—	B.fl.VII,567						
S. pycnostachya; Bentham, Fl. Austr. VII, 568 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,568						
S. setacea; R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,568						
S. semibarbata; R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,568	M.fr.VIII,104; XI,129.					
S. hemipogon; Bentham, Fl. Austr. VII, 569 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,569	M.fr.VIII,104.					
S. pubescens; R. Brown, prodr. 174 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,569	M.fr.VIII,104.					
S. aristiglumis; F. v. M. in Transact. Vict. Inst. 43 (1854)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,570	M.fr.VIII,103.					
S. Eriopus; Bentham, Fl. Austr. VII, 570 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,570						
S. trichophylla; Bentham, Fl. Austr. VII, 570 (1878)	W.A.	—	—	—	—	—	—	B.fl.VII,570						
S. scabra; Lindley in Mitch. Trop. Austr. 31 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,570						
S. Streptachne; F. v. M. in journ. R.S. of N.S.W. 237 (1881)	—	—	—	—	—	Q.	—	B.fl.VII,572						
S. micrantha; Cavanilles, icon. et descr. pl. V, 42, t. 467 (1799)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,574	M.fr.VIII,105	...	N.Z.			
S. verticillata; Nees in Spreng. syst. veg. cur. poster. 30 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,566	M.fr.VIII,105.					
S. Dichelachne; Steudel, nomencl. botan. edit. sec. I, 502 (1841)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,574	M.fr.VIII,105; XI,129.		N.Z.			
PENTAPOGON; R. Brown, prodr. fl. Nov. Holl. 173 (1810).														
P. Billardieri; R. Brown, prodr. 173 (1810)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,572	M.fr.VIII,106.					
ECHINOPOGON; Palisot, Essai d'une nouv. Agrostogr. 42, t. 9, f. 5 (1812). (Cinna partly.)														
E. ovatus; Palisot, Agrostogr. 42, t. 9 (1812)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,599	M.fr.VIII,106	...	N.Z.			
DIPLOPOGON; R. Brown, prodr. fl. Nov. Holl. 176 (1810). (Dipogonia.)														
D. setaceus; R. Brown, prodr. 176 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,573	M.fr.VI,86; VIII,106.					
AMPHIPOGON; R. Brown, prodr. 175 (1810). (Aegopogon partly, Pentacraspedion, Gamelythrum.)														
A. debilis; R. Brown, prodr. 175 (1810)	W.A.	—	—	—	—	—	—	B.fl.VII,597	M.fr.VIII,201.					
A. strictus; R. Brown, prodr. 175 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,597	M.fr.VIII,201.					

A. laguroides ; R. Brown, prodr. 175 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,598	M.fr.VI,86; VIII,201.			
A. cygnorum ; Nees in Lehm. pl. Preiss. II, 100 (1846)	W.A.	—	—	—	—	—	—	B.fl.VII,599				
A. turbinatus ; R. Brown, prodr. 175 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,599	M.fr.VI,86; VIII,201.			
PAPPOPHORUM ; Schreber, gen. pl. II, 787 (1791). (Enneapogon.)												
P. commune ; F. v. M., plants of Gregory's Exp. 10 (1859)...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,600	M.fr.VIII,200	...	P.	
SPOROBOLUS ; R. Brown, prodr. fl. Nov. Holl. 169 (1810). (Vilfa partly.)												
S. Virginicus ; Humboldt & Kunth, gram. I, 67 (1832) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,621	M.fr.VI,84 ...	...	As., Afr., Am.	
S. Indicus ; R. Brown, prodr. 170 (1810) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,622	M.fr.VI,84; X,76.		As., Afr., Am., N.Z.	
S. diander ; Palisot, Agrostogr. 26 (1812) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,622	M.fr.VI,84; VIII,140.		As.	
S. pulchellus ; R. Brown, prodr. 170 (1810) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,623	M.fr.VIII,140.			
S. Lindleyi ; Bentham, Fl. Austr. VII, 623 (1878) ...	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,623	M.fr.XI,130.			
S. actinocladus ; F. v. M., fragm. VIII, 140 (1874) ...	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,623	M.fr.VI,84; IX,194; XI,130.			
AGROSTIS ; Linné, syst. nat. 8 (1135) ; gen. plant. 19 (1737). (Deyeuxia, Didymochaeta, Trichodium, Bromidium, Lachnagrostis partly, Apera partly.)												
A. Muelleri ; Bentham, Fl. Austr. VII, 576 (1878) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,576	M.fr.VI,86 ...	...	N.Z.	
A. scabra ; Willdenow, spec. plant. I, 378 (1797) ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,576	...	...	Am.	
A. venusta ; Trinius in Mém. de l'Acad. de Petersb. VI, 340 (1850) ...	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,576				
A. aequata ; Nees in Hook. Lond. Journ. of Bot. II, 412 (1843) ...	—	—	T.	—	—	—	—	B.fl.VII,579				
A. Solandri ; F. v. M., Veget. of the Chath. Isl. 60 (1864)...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,579	M.fr.XI,27,130	...	N.Z.	
A. montana ; R. Brown, prodr. 171 (1810) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,581				
A. quadriseta ; R. Brown, prodr. 171 (1810) ...	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,581	..	...	N.Z.	
A. cylindrica ; R. Brown, prodr. 171 (1810) ...	W.A.	—	—	—	—	—	—	B.fl.VII,582				
A. densa ; F. v. M., Cens. 133 (1882) from Bentham (1878) ...	—	S.A.	—	V.	—	—	—	B.fl.VII,582				
A. frigida ; F. v. M., first gen. rep. 20 (1853)...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,583				
A. rudis ; Roemer & Schultes, syst. veg. II, 360 (1817) ...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,583	M.fr.VI,85.			
A. nivalis ; F. v. M. in Transact. Vict. Inst. 43 (1854) ...	—	—	—	V.	N.S.W.	—	—	B.fl.VII,584				
A. Gunniana ; F. v. M., Cens. 133 (1882) from Bentham (1878) ...	—	—	T.	—	—	—	—	B.fl.VII,584				
A. breviglumis ; F. v. M., Cens. 133 (1882) from Bentham (1878)...	—	—	—	—	N.S.W.	—	—	B.fl.VII,584				
COELACHNE ; R. Brown, prodr. fl. Nov. Holl. 187 (1810).												
C. pulchella ; R. Brown, prodr. 187 (1810) ...	—	—	—	—	—	Q.	—	B.fl.VII,626	...	...	As.	
MICRAIRA ; F. v. M., fragm. V, 208 (1866).												
M. subulifolia ; F. v. M., fragm. V, 208 (1866) ...	—	—	—	—	—	Q.	—	B.fl.VII,624	M.fr.V,208.			
AIRA ; Linné, gen. plant. 335 (1737). (Aera, Deschampsia.)												
A. caespitosa ; Linné, spec. plant. 64 (1753) ...	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,587	M.fr.VIII,138	...	E., As., Am., N.Z.	
TRISETUM ; Persoon, synops. plant. I, 97 (1805). (Airae subgenus.)												
T. subspicatum ; Palisot, Agrostogr. 88 (1812) ...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,588	M.fr.VIII,136	...	E., As., Am., N.Z.	
ERIACHNE ; R. Brown, prodr. fl. Nov. Holl. 183 (1810.)												
E. stipacea ; F. v. M., fragm. V, 206 (1866) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,626	M.fr.VIII,137; X,76.			
E. Armitii ; F. v. M. in Benth. Fl. Austr. VII, 627 (1878) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,627	...	...	As.	
E. squarrosa ; R. Brown, prodr. 183 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,628				
E. glauca ; R. Brown, prodr. 184 (1810) ...	W.A.	—	—	—	—	Q.	N.A.	B.fl.VII,628				
E. rara ; R. Brown, prodr. 183 (1810) ...	—	—	—	—	—	Q.	—	B.fl.VII,628				
E. agrostidea ; F. v. M., fragm. VII, 82 (1870) ...	—	—	—	—	—	—	N.A.	B.fl.VII,628	M.fr.VIII,137.			
E. ciliata ; R. Brown, prodr. 184 (1810) ...	—	—	—	—	—	Q.	N.A.	B.fl.VII,629				
E. setacea ; Bentham, Fl. Austr. VII, 629 (1878) ...	—	—	—	—	—	—	N.A.	B.fl.VII,629				
E. avenacea ; R. Brown, prodr. 184 (1810) ...	—	—	—	—	—	—	N.A.	B.fl.VII,629				

E. aristidea; F. v. M., fragm. V, 205 (1866)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,629	M.fr.V,205.	
E. pallescens; R. Brown, prodr. 184 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,630		As.
E. festucacea; F. v. M., fragm. V, 205 (1866)	—	—	—	—	—	—	N.A.	B.fl.VII,630	M.fr.V,205.	
E. ovata; Nees in Hook. Lond. Journ. II, 418 (1843)	W.A.	S.A.	—	—	—	—	N.A.	B.fl.VII,630	M.fr.VIII,137.	
E. meliacea; F. v. M., fragm. V, 205 (1866)	—	—	—	—	—	—	N.A.	B.fl.VII,631	M.fr.V,205.	
E. pallida; F. v. M. in Benth. Fl. Austr. VII, 631 (1878)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,631		
E. scleranthoides; F. v. M., fragm. VIII, 233 (1874)	—	S.A.	—	—	—	—	—	B.fl.VII,631		
E. mucronata; R. Brown, prodr. 184 (1810)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,632		
E. obtusa; R. Brown, prodr. 184 (1810)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,632	M.fr.VIII,137; XI,130.	
E. capillaris; R. Brown, prodr. 184 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,632		
ANISOPOGON; R. Brown, prodr. fl. Nov. Holl. 176 (1810).										
A. avenaceus; R. Brown, prodr. 176 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,590	M.fr.VII,106.	
DANTHONIA; De Candolle, Fl. Franç. III, 32 (1805). (Amphibromus, Monacather, Plinthacanthus, Danthoinia.)										
D. paradoxa; R. Brown, prodr. 177 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,591		
D. bipartita; F. v. M., fragm. I, 160 (1859)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,592	M.fr.VIII,136.	
D. carphoides; F. v. M. in Benth. Fl. Austr. VII, 592 (1878)	—	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,592	M.fr.XI,130.	
D. penicillata; F. v. M., fragm. VIII, 135 (1873)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,592	M.fr.VIII,135	P., N.Z.
D. robusta; F. v. M. in Transact. Vict. Inst. 44 (1854)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,593	M.fr.VIII,136.	
D. pauciflora; R. Brown, prodr. 177 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,596		N.Z.
D. nervosa; J. Hooker, Fl. Tasman. II, 121, t. 163 (1860)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,589	M.fr.VIII,135.	
ASTREBLA; F. v. M. fragm. phytogr. Austr. X, 76 (1876).										
A. pectinata; F. v. M., fragm. X, 76 (1876)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,602	M.fr.VIII,134.	
A. triticoides; F. v. M., fragm. X, 76 (1876)	—	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,602	M.fr.VIII,134.	
MICROCHLOA; R. Brown, prodr. fl. Nov. Holl. 208 (1810).										
M. setacea; R. Brown, prodr. 208 (1810)	—	—	—	—	—	—	N.A.	B.fl.VII,608		As., Afr., Am.
CYNODON; L. C. Richard in Persoon, synops. I, 85 (1805). (Fibrichia 1802.)										
C. Dactylon; L. C. Richard in Persoon, synops. I, 85 (1805)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,609	M.fr.VIII,113	E., As., Afr.
C. tenellus; R. Brown, prodr. 187 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,609	M.fr.VIII,113	As.
C. convergens; F. v. M., fragm. VIII, 113 (1874)	—	—	—	—	—	Q.	N.A.	B.fl.VII,610	M.fr.X,76.	
C. ciliaris; Bentham, Fl. Austr. VII, 610 (1878)	—	S.A.	—	—	—	—	—	B.fl.VII,610		
CHLORIS; Swartz, nov. gen. et spec. plant. 25 (1788).										
C. unispicea; F. v. M., fragm. VII, 118 (1871)	—	—	—	—	—	Q.	—	B.fl.VII,611	M.fr.VII,118.	
C. pumilio; R. Brown, prodr. 186 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,611		
C. pectinata; Bentham, Fl. Austr. VII, 612 (1878)	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,612		
C. divaricata; R. Brown, prodr. 186 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,612	M.fr.VI,85.	
C. acicularis; Lindley in Mitch. Trop. Austr. 33 (1848)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,612	M.fr.VI,85.	
C. truncata; R. Brown, prodr. 186 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,612	M.fr.VI,85; X,130.	
C. ventricosa; R. Brown, prodr. 186 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,613	M.fr.VI,85.	
C. barbata; Swartz, nov. gen. et spec. plant. 25 (1788)	—	S.A.	—	—	—	—	N.A.	B.fl.VII,613	M.fr.VI,85	As., Afr.
C. pallida; Hackel in Engler's bot. Jahrb. VI, 248 (1885)	W.A.	—	—	—	—	—	N.A.	—		
C. scariosa; F. v. M., fragm. VI, 85 (1868)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.VII,614	M.fr.XI,130.	
ELEUSINE; J. Gaertner, de fruct. I, 7 (1788). (Leptochloa, Dactyloctenium, Acrachne.)										
E. cruciata; Lamarck, tabl. enclop. méth. I, 203, t. 48 (1791)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,615	M.fr.VIII,111	As., Afr., Am., P.
E. Indica; J. Gaertner, de fruct. I, 7 (1788)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,615	M.fr.VIII,112	As., Afr., Am., P.
E. verticillata; Roxburgh, Fl. Ind. ed. Car. & Wall. I, 346 (1820)	—	—	—	—	—	Q.	N.A.	B.fl.VII,616	M.fr.VIII,112	As., Afr.
E. digitata; Sprengel, syst. cur. post. 36 (1827)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,617	M.fr.I,216; VIII,112; XI,130.	
E. Chinensis; F. v. M., Cens. 134 (1882) from Koenig (1821)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,617	M.fr.VIII,132	As.
E. polystachya; F. v. M.; Cynodon; R. Brown, prodr. 187 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,617		

CENTOTHECA; Desvaux in Palisot, Agrost. 69, t. 14 (1812).

C. lappacea; Desvaux in Palisot, Agrost. 69, t. 14 (1812) ... . — — — — — Q. — B.fl.VII,640 M.fr.VI,85; VIII,116; X,76. As., Afr., P.

POA; Linné, gen. plant. 20 (1737). (Glyceria partly.)

P. Labillardieri; Steudel, syn. glum. I, 262 (1855) ... ... W.A. S.A. T. V. — — N.A. B.fl.VII,651
P. homomalla; Nees in Lehm. pl. Preiss. II, 104 (1846) ... ... W.A. — — — — — — B.fl.VII,651
P. caespitosa; G. Forster, fl. ins. Austr. prodr. 89 (1786) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,651 M.fr.VIII,133; XI,13. N.Z.
P. Maxwelli; Bentham, Fl. Austr. VII, 653 (1878) ... ... W.A. — — — — — — B.fl.VII,652
P. nodosa; Nees in Lehm. pl. Preiss. II, 105 (1846) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,653 M.fr.VIII,132.
P. saxicola; R. Brown, prodr. 654 (1810) ... ... — — T. — — — — B.fl.VII,654
P. lepida; F. v. M., fragm. VIII, 130 (1874) ... ... W.A. S.A. — V. N.S.W. — — B.fl.VII,654 M.fr.XI,130.
P. Fordeana; F. v. M., fragm. VIII, 130 (1874) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,657 M.fr.VIII,130.
P. fluitans; Scopoli, flor. Carniol. 106 (1772) ... ... W.A. S.A. T. V. N.S.W. — — B.fl.VII,657 M.fr.VIII,129 ... E., As., Afr., Am.
P. latispicea; F. v. M., Cens. of Austr. pl. 134 (1882) ... ... — — — — N.S.W. Q. — B.fl.VII,658 M.fr.VIII,127.
P. syrtica; F. v. M. in Transact. Vict. Inst. 45 (1854) ... ... W.A. S.A. T. V. — — — B.fl.VII,658 M.fr.VIII,130 ... N.Z.
P. dives; F. v. M., Cens. of Austr. pl. 134 (1882) ... ... — — — V. — — — B.fl.VII,659 M.fr.VI,147; VIII,129; XI,130; III,147.
P. ramigera; F. v. M. in Transact. Vict. Inst. 45 (1854) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,659 M.fr.VI,85; VIII,131; XI,130.

FESTUCA; Dillenius, nov. gen. 90, t. 3 (1719). (Glyceria partly, Vulpia partly, Schedonorus, Schenodorus, Brizopyrum.)

F. duriuscula; Linné, spec. plant. 74 (1753) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,663 M.fr.VI,85 ... ... E., As., Am., P.
F. scirpoidea; F. v. M., fragm. VIII, 129 (1874) ... ... W.A. — — — — — — B.fl.VII,655 M.fr.XI,130.
F. litoralis; Labillardière, Nov. Holl. plant. spec. I, 22, t. 27 (1804) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,656 M.fr.VIII,129 ... N.Z.
F. Hookeriana; F. v. M. in J. Hooker, Fl. Tasman. II, 122, t. 115 (1860) ... — — T. V. N.S.W. — — B.fl.VII,656 M.fr.VIII,131.

DIPLACHNE; Palisot, Essai d'une nouv. Agrostogr. 9, t. 16 (1812). (Uralepis.)

D. loliiformis; F. v. M. in Benth. Fl. Austr. VII, 618 (1878) ... ... — S.A. — V. N.S.W. Q. — B.fl.VII,616 M.fr.VIII,128; XI,130. As.
D. Muelleri; Bentham, Fl. Austr. VII, 619 (1878) ... ... W.A. S.A. — — — — N.A. B.fl.VII,619
D. fusca; Palisot, Agrost. 163 (1812) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,619 M.fr.VIII,127 ... As., Afr.
D. parviflora; Bentham, Fl. Austr. VII, 620 (1878) ... ... — — — — — — N.A. B.fl.VII,620 M.fr.VIII,129.

TRIODIA; R. Brown, prodr. fl. Nov. Holl. 182 (1810). (Sieglingia 1799.)

T. Mitchelli; Bentham, Fl. Austr. VII, 606 (1878) ... ... W.A. S.A. — — N.S.W. Q. N.A. B.fl.VII,606
T. pungens; R. Brown, prodr. 182 (1810) ... ... — S.A. — — — Q. N.A. B.fl.VII,606 M.fr.VIII,129.
T. Cunninghamii; Bentham, Fl. Austr. VII, 606 (1878) ... ... — — — — — Q. N.A. B.fl.VII,606
T. irritans; R. Brown, prodr. 182 (1810) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,607 M.fr.VIII,129.
T. procera; R. Brown, prodr. 182 (1810) ... ... — — — — — — N.A. B.fl.VII,607
T. microstachya; R. Brown, prodr. 182 (1810) ... ... — — — — — — N.A. B.fl.VII,607
T. microdon; F. v. M., syst. Cens. of Austr. pl. 135 (1882) from Bentham (1878) — — — — N.S.W. — — B.fl.VII,605

DISTICHLIS; Rafinesque in Journ. de Phys. LXXXIX, 104 (1819). (Uniola partly, Brizopyrum partly.)

D. maritima; Rafinesque in Journ. de Phys. LXXXIX, 104 (1819) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,637 M.fr.VIII,129 ... Am.

BROMUS; Dillenius in Linné, syst. nat. 8 (1735); Linné, gen. plant. 15 (1737).

B. arenarius; Labillardière, Nov. Holl. pl. spec. I, 23, t. 28 (1804) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,660 M.fr.VI,126; XI,20 ... N.Z.

ERAGROSTIS; Palisot, Essai d'une nouv. Agrostogr. 70, t. 14 (1812). (Poa partly.)

E. tenella; Palisot in Roem. et Schult. syst. veg. II, 576 (1817) ... ... W.A. S.A. — V. N.S.W. Q. N.A. B.fl.VII,643 M.fr.VIII,132; XI,130 As.
E. nigra; Nees in Steud. nomencl. bot. I, 563 (1841) ... ... — — — — N.S.W. Q. — B.fl.VII,643 ... ... As.
E. imbecilla; Bentham, Fl. Austr. VII, 643 (1878) ... ... — — — — — Q. — B.fl.VII,643 ... ... N.Z.
E. trichophylla; Bentham, Fl. Austr. VII, 644 (1878) ... ... — S.A. — — — — — B.fl.VII,644
E. leptocarpa; Bentham, Fl. Austr. VII, 644 (1878) ... ... — S.A. — — — Q. N.A. B.fl.VII,644
E. megalosperma; F. v. M. in Benth. Fl. Austr. VII, 644 (1878) ... ... — — — — N.S.W. Q. — B.fl.VII,644 [P.
E. pilosa; Palisot, Agrost. 71 (1812) ... ... W.A. S.A. — V. N.S.W. Q. — B.fl.VII,645 M.fr.VIII,133; XI,130. E., As., Afr., Am.,

E. leptostachya; Steudel, syn. glum. I, 279 (1855)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,645						
E. Schultzii; Bentham, Fl. Austr. VII, 646 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,646						
E. diandra; Steudel, syn. glum. I, 279 (1855)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,646	M.fr.XI,130.					
E. Brownii; Nees in Steud. nomencl. bot. I, 562 (1841)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,646	...	..	..	As.		
E. concinna; Steudel, syn. glum. I, 279 (1855)	—	S.A.	—	—	—	Q.	N.A.	B.fl.VII,647						
E. speciosa; Steudel, syn. glum. I, 279 (1855)	W.A.	S.A.	—	—	—	Q.	N.A.	B.fl.VII,648						
E. laniflora; Bentham, Fl. Austr. VII, 648 (1878)	—	S.A.	—	—	N.S.W.	Q.	—	B.fl.VII,648						
E. eriopoda; Bentham, Fl. Austr. VII, 648 (1878)	W.A.	S.A.	—	—	N.S.W.	Q.	N.A.	B.fl.VII,648						
E. setifolia; Nees in Hook. Lond. Journ. II, 419 (1843)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,648						
E. lacunaria; F. v. M., first gen. Rep. 20 (1853)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,649						
E. falcata; Gaudichaud in Voy. Freyc. Bot. 408, t. 25 (1826)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,649						
E. stenostachya; Steudel, syn. glum. I, 279 (1855)	—	—	—	—	—	Q.	N.A.	B.fl.VII,650						
HETERACHNE; Bentham in J. Hook. icon. plant. XIII, 39 (1877).														
H. Brownii; Bentham in J. Hook. icon. XIII, 40 (1877)	—	—	—	—	—	Q.	N.A.	B.fl.VII,635	M.fr.VIII,132.					
H. Gulliveri; Bentham in J. Hook. icon. XIII, 39, t. 1250 (1877)	—	—	—	—	—	Q.	N.A.	B.fl.VII,635						
ECTROSIA; R. Brown, prodr. fl. Nov. Holl. 185 (1810).														
E. Schultzii; Bentham, Fl. Austr. VII, 633 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,633						
E. leporina; R. Brown, prodr. 186 (1810)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,633	M.fr.VIII,109.					
E. agrostoides; Bentham, Fl. Aust. VII, 634 (1878)	—	—	—	—	—	—	N.A.	B.fl.VII,634						
E. Gulliveri; F. v. M., fragm. VIII, 201 (1874)	—	—	—	—	—	Q.	N.A.	B.fl.VII,634	M.fr.VIII,201.					
ELYTHROPHORUS; Palisot, Essai d'une nouv. Agrostogr. 67, t. 14 (1812).														
E. articulatus; Palisot, Agrost. 67, t. 14 (1812)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,638	M.fr.VII,109	...	As., Afr.			
TRIRHAPHIS; R. Brown, prodr. fl. Nov. Holl. 185 (1810.) (Triraphis.)														
T. mollis; R. Brown, prodr. 185 (1810)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,603	M.fr.VIII,108.					
T. pungens; R. Brown, prodr. 185 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,604	M.fr.VIII,108,125.					
T. bromoides; F. v. M., fragm. VIII, 108 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,604	M.fr.VIII,108.					
T. danthonioides; F. v. M., fragm. VIII, 125 (1874)	W.A.	—	—	—	—	—	—	B.fl.VII,604	M.fr.VIII,108.					
AGROPYRON; J. Gaertner, nov. comm. Petrop. XIV, 539 (1770). (Agropyrum, Triticum partly, Anthosachne, Vulpia partly.)														
A. scabrum; Palisot, Agrost. 102 (1812)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,665	M.fr.VI,85; XI,27,130.	N.Z.				
A. velutinum; Nees in Hook. Lond. Journ. of Bot. II, 417 (1843)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,665						
A. pectinatum; Palisot, Agrost. 102 (1812)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,666	M.fr.VI,85.					
ARUNDO; Tournefort, inst. 526 (1700), from Varro. (Phragmites.)														
A. Phragmites; Dodoens, stirp. hist. pemptad. 602 (1583)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,637	...	...	E., As., Afr., Am., P., N.Z.			
A. Roxburghii; F. v. M., syst. Cens. of Austr. pl. 135 (1882) from Kunth (1843)	—	—	—	—	—	Q.	N.A.	—	M.fr.XI,139.	As.				
BAMBUSA; Schreber, gen. plantar. I, 236 (1789).														
B. Arnhemica; F. v. M. in Austral. journ. of Pharm. new ser. I, 447 (1886)	—	—	—	—	—	—	N.A.	—	M.fr.VI,85; XI,129.					

ACOTYLEDONEAE.

A. L. de Jussieu, genera plantarum 1 (1789).

ACOTYLEDONEAE VASCULARES.

Meissner, plantarum vascularium genera, 429 (1843).

RHIZOSPERMAE.

G. Weber, primit. Fl. Holsat. 74 (1780).—Genera 5, Species 11.

AZOLLA; Lamarck, Encycl. méthod. I, 343 (1783).

A. pinnata; R. Brown, prodr. 167 (1810)	—	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,679	M.fr.V,140	...	As., Afr.
A. filiculoides; Lamarck, Encycl. méth. I, 343 (1783)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,680	M.fr.V,140; XI,127.		Am., N.Z.

SALVINIA; Micheli, nov. plant. gen. 107, t. 58 (1729).

S. cucullata; Roxburgh, fl. Indic. IV, 547 (1843)	W.A.	—	—	—	—	—	—	—	...	...	...	As.

MARSILEA; Linné, syst. nat. 8 (1735); gen. plant. 326 (1737). (Marsiglia.)

M. quadrifolia; Linné, spec. plant. 1099 (1753)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,683	M.fr.V,140	...	E., As., Afr.

PILULARIA; Vaillant, Bot. Paris. 159, t. 15 (1727).

P. globulifera; Linné, spec. plant. 1100 (1753)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,684	M.fr.V,140	...	E., As., Afr., N.Z

ISOETES; Linné, skanska resa 420 (1751).

I. tripus; A. Braun in Berl. Monatsb. 559 (1863)	W.A.	—	—	—	—	—	—	—				
I. Drummondii; A. Braun in Berl. Monatsb. 593 (1863)	W.A.	—	—	V.	N.S.W.	—	—	B.fl.VII,672	M.fr.V,140.			
I. elatior; F. v. M. & A. Braun in Linnaea XXV, 722 (1852)	—	—	T.	—	—	—	—	—	M.fr.V,140; XI,139.			
I. humilior; F. v. M. & A. Braun in Linnaea XXV, 722 (1852)	—	—	T.	—	—	—	—	—	M.fr.V,140; XI,139.			
I. Gunnii; A. Braun in Berl. Monatsb. 535 (1868)	—	—	T.	—	—	—	—	—	M.fr.XI,139.			
I. Muelleri; A. Braun in Berl. Monatsb. 541 (1868)...	—	—	—	—	—	Q.	—	—	M.fr.XI,139.			

LYCOPODINAE.

Swartz, synops. filic. XV (1806).—Genera 5, Species 21.

PSILOTUM; Swartz in Schrader, Journ. fuer die Bot. II, 109 (1800).

P. triquetrum; Swartz, synops. filic. 187 (1806)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,681	M.fr.V,112; X,118.		As., Afr., Am.
P. complanatum; Swartz, synops. filic. 187 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,682	M.fr.V,112	...	As., Afr., Am., P.

TMESIPTERIS; Bernhardi in Schrader, Journ. fuer die Bot. II, 131 (1800).

T. Tannensis; Bernhardi in Schrad. Journ. II, 131, t. 2 (1800)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,680	M.fr.V,112; X,118.		P., N.Z.

LYCOPODIUM; Ruppius, Fl. Jenens. 32 (1718), from Tabernaemontanus (1588).

L. Selago; Linné, spec. plant. 1102 (1753)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,674	M.fr.V,111; X,118.		E., As., Afr., Am., N.Z.
L. Phlegmaria; Linné, spec. plant. 1101 (1753)	—	—	—	—	—	Q.	—	B.fl.VII,674	M.fr.V,111; X,118.		As., Afr. [N.Z.
L. clavatum; Linné, spec. plant. 1101 (1753)...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,675	M.fr.V,111	...	E., As., Afr., Am., P.,
L. Carolinianum; Linné, spec. plant. 1104 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,675	M.fr.V,111; X,118.		As., Afr., Am., P., N.Z.
L. laterale; R. Brown, prodr. 165 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,675	M.fr.V,111; VI,124; X,118.		P., N.Z.
L. cernuum; Linné, spec. plant. 1103 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,676	M.fr.V,111; X,118.		As., Afr., Am., N.Z.
L. densum; Labillardière, Nov. Holl. plant. spec. II, 104, t. 251 (1806)... ...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,676	M.fr.V,111; X,118.		P., N.Z.
L. scariosum; G. Forster, florul. ins. Austr. prodr. 87 (1786)	—	—	T.	V.	—	—	—	B.fl.VII,676	M.fr.V,112.	...	Am., N.Z.
L. volubile; G. Forster, florul. ins. Austr. prodr. 86 (1786)	—	—	—	—	—	—	N.A.	B.fl.VII,677	M.fr.V,112.	...	As., Am., P., N.Z.

SELAGINELLA ; Palisot, prodr. des fam. de l'Aethéogamie 101 (1805). (Lycopodium partly.)

S. Preissiana ; Spring in Mém. de l'Ac. de Brux. 61 (1842) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,677 M.fr.V,112; VIII,275.
S. uliginosa ; Spring in Mém. de l'Ac. de Brux. 60 (1842) ... ... W.A. S.A. T. V. N.S.W. Q. — B.fl.VII,678 M.fr.V,112; VIII,275.
S. flabellata ; Spring in Mém. de l'Ac. de Brux. 174 (1849)... ... ... — — — — — Q. — B.fl.VII,678 M.fr.VIII,175,275. As., Am., P.
S. Australiensis ; Baker in Britten's Journ. of Bot. 144 (1883) ... ... — — — — — Q. — B.fl.VII,678 M.fr.V,112; VIII,274.
S. Belangeri ; Spring in Mém. de l'Ac. de Brux. 242 (1849) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,679 ... ... As., P.
S. Bakeriana ; Bailey, suppl. Queensl. Fl. 62 (1886)... ... ... — — — — — Q. — —
S. Brisbanensis ; Bailey, suppl. Queensl. Fl. 62 (1886) ... ... — — — — — Q. — —
S. Pumilio ; Spring in Mém. de l'Ac. de Brux. 241 (1849) ... ... — — — — — Q. — —

PHYLLOGLOSSUM ; Kunze in Mohl & Schlechtendal, Bot. Zeitung, 721 (1843).

P. Drummondii ; Kunze in der Bot. Zeit. 721 (1843)... ... ... W.A. — T. V. N.S.W. — — B.fl.VII,672 M.fr.V,112; X,118.

FILICES.

Linné, gen. plant. 322 (1737).—Genera 35, Species 212.

OPHIOGLOSSUM ; Tournefort, inst. rei herb. 548, t. 325 (1700), from Bock (1539). (Ophioderma.)

O. vulgatum ; C. Bauhin, pinax. theatr. bot. 354 (1623) ... ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,688 M.fr.V,112; VI,124. E., As., Afr., Am., P., [N.Z.
O. pendulum ; Linné, sp. pl. edit. sec. 1518 (1763) ... ... — — — — N.S.W. Q. — B.fl.VII,689 M.fr.V,113; VI,124. As., Afr.

BOTRYCHIUM ; Swartz in Schrader, Journ. fuer die Bot. II, 110 (1800).

B. Lunaria ; Swartz in Schrad. Journ. II, 110 (1800) ... ... — — T. V. N.S.W. — — B.fl.VII,690 M.fr.V,113 ... E., As., Am.
B. ternatum ; Swartz, synops. filic. 172 (1806) ... ... — — T. V. N.S.W. Q. — B.fl.VII,690 M.fr.V,113 ... As., Am., N.Z.

HELMINTOHSTACHYS ; Kaulfuss, enum. filic. Chamiss. 28 (1824).

H. Zeylanica ; Kaulfuss, enum. fil. 28 (1824) ... ... — — — — — Q. — B.fl.VII,690 M.fr.V,113; IX,190. As., P.

LYGODIUM ; Swartz in Schrader, Journ. fuer die Bot. II, 7 & 106 (1800). (Hydroglossum, Lygodictyon.)

L. scandens ; Swartz in Schrad. Journ. II, 106 (1800) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,691 M.fr.V,113; VII,84; X,118. As., Afr., P., [N.Z.
L. reticulatum ; Schkuhr. kryptog. Gewaechse 139, t. 139 (1809) ... ... — S.A. — — — Q. — B.fl.VII,692 M.fr.VII,83; VIII,157. P.
L. Japonicum ; Swartz, syn. filic. 154 (1806) ... ... — — — — — Q. N.A. B.fl.VII,692 M.fr.V,113; VII,84,156. As., P.
L. dichotomum ; Swartz, syn. filic. 154 (1806) ... ... — — — — — Q. — —

SCHIZAEA ; Smith in Mém. de l'Acad. Turin V, 149, t. 19 (1791). (Lopidium, Actinostachys.)

S. rupestris ; R. Brown, prodr. 162 (1810) ... ... — — — — N.S.W. — — B.fl.VII,693 M.fr.V,113.
S. fistulosa ; Labillardière, Nov. Holl. pl. spec. II, 103, t. 250 (1806) ... ... — S.A. T. V. N.S.W. — — B.fl.VII,693 M.fr.V,113 ... Afr., Am., P., N.Z.
S. dichotoma ; Smith in Mém. de l'Ac. Tur. V, 149 (1791) ... ... — — — — N.S.W. Q. N.A. B.fl.VII,694 M.fr.V,113; VIII,275; X,118 As., Afr., [Am., P., N.Z.
S. bifida ; Willdenow in act. Acad. Erfurt. 30, t. 3, f. 3 (1802) ... ... — S.A. T. V. N.S.W. Q. N.A. B.fl.VII,693 M.fr.V,113 ... N.Z.

ANGIOPTERIS ; G. Hoffmann in Comment. Goett. XII, 29 (1796).

A. evecta ; G. Hoffmann in Comment. Goett. XII, 29 (1796) ... ... — — — — N.S.W. Q. — B.fl.VII,694 M.fr.V,114; VII,120; X,118. As., Afr., P.

MARATTIA ; Swartz, nov. gen. et spec. plant. 8 & 128 (1788).

M. fraxinea ; Smith, plant. icon. hact. ined. t. 48 (1790) ... ... — — — — N.S.W. Q. — B.fl.VII,695 M.fr.V,114; X,118. As., Afr., Am., P., N.Z.

TRICHOMANES ; Linné, syst. nat. 8 (1835) ; hort. Cliffort. 476 (1737).

T. peltatum ; Baker in Journ. Linn. Soc. IX, 336, t. 8 (1865) ... ... — — — — — Q. — B.fl.VII,701 M.fr.X,117 ... P.
T. Vitiense ; Baker in Journ. Linn. Soc. IX, 338, t. 8 (1865) ... ... — — — — N.S.W. Q. — B.fl.VII,701 M.fr.VIII,32; IX,132. P.
T. Motleyi ; Bosch, Hymenoph. Javan. (1861) ... ... — — — — — Q. — — ... ... As., P.
T. cuspidatum ; Willdenow, spec. plant. V, 499 (1810) ... ... — — — — — Q. — — M.fr.XI,131&139. As., Afr.
T. parvulum ; Poiret, Encycl. méth. VIII, 44 (1810) ... ... — — — — N.S.W. Q. — B.fl.VII,701 M.fr.VII,155 ... As., Afr., P.
T. digitatum ; Swartz, synops. filic. 370 & 422 (1806) ... ... — — — — N.S.W. — — B.fl.VII,702 ... ... As., Afr., P.
T. venosum ; R. Brown, prodr. 159 (1810) ... ... — — T. V. N.S.W. Q. — B.fl.VII,702 M.fr.V,116;IX,191. N.Z.
T. Javanicum ; Blume, flor. Jav. filic. 224 (1829) ... ... — — — — — Q. — B.fl.VII,702 ... ... As., P.

T. rigidum ; Swartz, nov. gen. et spec. pl. 137 (1788)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,702	M.fr.V,115	...	As., Afr., Am., P., N.Z.
T. maximum ; Blume, enum. fil. Javan. 228 (1828)	—	—	—	—	—	Q.	—	...	...	...	As., P.
T. bipunctatum ; Poiret, Encycl. méth. VIII, 44 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,703	M.fr.X,117;XI,132.		As., P.
T. humile ; G. Forster, flornl. ins. Austr. prodr. 84 (1786)...	—	—	—	V.	N.S.W.	—	—	—	...	...	As., P., N.Z.
T. caudatum ; Brackenridge, Bot. of Wilk. U.S. Exped. fil. 256, t. 36 (1854) ...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,703	M.fr.V,116; VII,122		P.
T. meifolium ; Bory in Willd. sp. pl. V, 509 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,703	M.fr.VII,121; XI,132.		As., Afr., P.
T. parviflorum ; Poiret, Encycl. méth. VIII, 44 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,704	M.fr.VII,122	...	As., Afr.
T. Sayeri ; F. v. M. & Baker in Britten's Journ. of Bot. XXV, 163 (1887) ...	—	—	—	—	—	Q.	—	—			
HYMENOPHYLLUM ; Smith in Roemer, Archiv I, 56 (1797).											
H. Malingii ; J. Hooker, Handb. N.Z. Flora I, 357 (1864) implied	—	—	T.	—	—	—	—	—	...	...	N.Z.
H. marginatum ; Hooker & Greville, icon. filic. I, t. 34 (1829)	—	—	T.	--	N.S.W.	—	—	B.fl.VII,705			
H. nitens ; R. Brown, prodr. 159 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,705	...	...	P., N.Z.
H. Javanicum ; Sprengel, syst. veg. IV, 132 (1827)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,705	M.fr.V,116	...	As., P., N.Z.
H. Tunbridgense ; Smith in Roemer, Archiv. I, 56 (1797)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,706	M.fr.V,116		E., As., Afr., Am., N.Z., P.
H. multifidum ; Swartz, synops. filic. 149 & 378 (1806)	—	—	—	—	N.S.W.	—	—	B.fl.VII,707	...	...	P., N.Z.
H. bivalve ; Swartz, synops. filic. 146 & 372 (1806)	—	—	—	—	N.S.W.	—	—	—	...	...	P., N.Z.
CERATOPTERIS ; Brogniart in Bull. de la Soc. philom. 186 (1821). (Parkeria.)											
C. thalictroides ; Brogniart in Bull. Soc. philom. 186 (1821)	—	—	—	—	—	Q.	N.A.	B.fl.VII,695	M.fr.V,122; VIII,158.		As., Afr., Am., P.
GLEICHENIA ; Smith in Mém. de l'Acad. Turin V, 418 (1791). (Mertensia, Platyzoma, Stromatopteris.)											
G. platyzoma ; F. v. M., fragm. V, 114 (1866)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,696	M.fr.V,114; VIII,157; X,118.		
G. circinata ; Swartz, syn. filic. 165 & 394 (1806)	—	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,697	M.fr.V,115	...	As., P., N.Z.
G. dicarpa ; R. Brown, prodr. 161 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,698	M.fr.V,115	...	As., P., N.Z.
G. flabellata ; R. Brown, prodr. 161 (1810)	—	—	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,698	M.fr.V,114; VIII,157.		P., N.Z.
G. Hermanni ; R. Brown, prodr. 161 (1810)	—	—	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,698	M.fr.V,114; X,118.		As., Afr., Am., P., N.Z.
OSMUNDA ; Tournefort, inst. rei herb. 547, t. 324 (1700) from l'Obel (1576). (Todea, Leptopteris.)											
O. barbara ; Thunberg, prodr. plant. Capens. 171 (1800)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,699	M.fr.V,114	...	Afr., N.Z.
O. Fraseri ; F. v. M., Cens. 137 (1882) from Hooker & Greville (1829)	—	—	—	—	N.S.W.	—	—	B.fl.VII,700	M.fr.V,114; VII,156.		P.
O. Moorei ; F. v. M., Cens. 137 (1882) from Baker (1873)	—	—	—	—	N.S.W.	—	—	B.fl.VII,700	M.fr.VIII,157.		
CYATHEA ; Smith in Mém. de l'Acad. Turin V, 416 (1791). (Hemitelia, Amphicosmia).											
C. Lindsayana ; Hooker, syn. filic. 25 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,708	M.fr.VIII,177.		
C. arachnoidea ; Hooker, syn. filic. 24 (1866)...	—	—	—	—	—	Q.	—	B.fl.VII,708	M.fr.VI,200	...	As.
C. Macarthurii ; F. v. M., fragm. VII, 177 (1874)	—	—	—	—	N.S.W.	—	—	B.fl.VII,708	M.fr.VIII,176; X,120.		
C. medullaris ; Swartz, synops. filic. 140 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,708	M.fr.V,116; VIII,177.		As., P., N.Z.
C. Cunninghami ; J. Hooker, icon. plant. 985 (1854)	—	—	T.	V.	—	—	—	—	...	...	N.Z.
C. brevipinnea ; Baker in Bentham, fl. Austr. VII, 709 (1878)	—	—	—	—	N.S.W.	—	—	B.fl.VII,709	M.fr.XI,139.		
C. Moorei ; F. v. M., Cens. 137 (1882) from Baker (1872)	—	—	—	—	N.S.W.	—	—	B.fl.VII,709	M.fr.VIII,157.		
ALSOPHILA ; R. Brown, prodr. 158 (1810).											
A. Rebeccae ; F. v. M., fragm. V, 53 (1865)	—	—	—	—	—	Q.	—	B.fl.VII,710	M.fr.V,117,215; VIII,180.		
A. Loddigesii ; Kunze in Schlecht. Linnaea XXIII, 221 (1850)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,710.			
A. Australis ; R. Brown, prodr. 158 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,710	M.fr.V,52,116; VII,120; VIII,178.		
A. excelsa ; R. Brown, prodr. 158 (1810)	—	—	—	—	N.S.W.	Q.	—	—			
A. Leichhardtiana ; F. v. M., fragm. V, 53 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,711	M.fr.V,117; VI,200; VII,120.		
A. Robertsiana ; F. v. M., fragm. V, 54 (1865)	—	—	—	—	—	Q.	—	B.fl.VII,712	M.fr.V,117,213.		
DICKSONIA ; L'Héritier, sert. Angl. 31 (1788). (Cibotium, Patania, Dennstaedtia, Deparia, Balantium.)											
D. Billardieri ; F. v. M., fragm. VIII, 175 (1874)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,712	M.fr.V,117; VI,199; VII,120.		P., N.Z.
D. Youngiae ; C. Moore in Hooker & Baker synops. filic. 51 (1865)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,713	M.fr.VI,200.		

D. davallioides ; R. Brown, prodr. 158 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,713	M.fr.VIII,158	...	As., N.Z.
D. nephrodioides ; F. v. M., Cens. 137 (1882) from Baker (1872)	—	—	—	—	N.S.W.	—	—	B.fl.VII,714	M.fr.VIII,157; IX,78; X,104.		
DAVALLIA ; Smith in Mém. de l'Acad. de Turin V, 414 (1791). (Microlepia, Humata.)											
D. solida ; Swartz, synops. filic. 132 & 345 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,715	...	...	As., P.
D. elegans ; Swartz, synops. filic. 132 & 347 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,715	M.fr.V,118; VIII,157.		As., Afr.
D. pyxidata ; Cavanilles, descripcion de las plantas n. 694 (1802)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,716	M.fr.V,118	...	P.
D. pedata ; Smith in Mém. de l'Ac. de Turin V, 414 (1791)	—	—	—	—	—	Q.	—	B.fl.VII,716	M.fr.V,118	...	As., Afr.
D. dubia ; R. Brown, prodr. 157 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,716	M.fr.V,118.		
D. flaccida ; R. Brown, prodr. 157 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,717	M.fr.V,118; VII,156.		As., Afr., P.
D. tripinnata ; F. v. M. in Benth. Fl. Austr. VII, 717 (1878)	—	—	—	—	—	Q.	—	B.fl.VII,717			
VITTARIA ; Smith in Mém. de l'Acad. de Turin V, 414 (1791).											
V. elongata ; Swartz, synops. filic. 109 & 302 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,718	M.fr.V,122; VII,121.		As., Afr., P.
V. falcata ; Kunze in Feé, gen. fil. t. 6, f. 1 (1852)	—	—	—	—	—	Q.	—	—	...	...	As.
LINDSAYA ; Dryander in Mém. de l'Acad. de Turin V, 413 (1791). (Lindsaea, Isoloma, Synophlebium, Schizoloma.)											
L. linearis ; Swartz, synops. filic. 118 & 318, t. 3 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,719	M.fr.V,119	...	P., N.Z.
L. dimorpha ; Bailey, Queensland Ferns 19 (1874)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,719			
L. cultrata ; Swartz, synops. filic. 119 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,719	...	...	As., Afr.
L. flabellulata ; Dryander in Transact. Linn. Soc. III, 41, t. 8 (1797)	—	—	—	—	—	Q.	N.A.	B.fl.VII,720	M.fr.V,119	...	As.
L. decomposita ; Willdenow, spec. plant. V, 425 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,720	...	...	As., P.
L. trichomanoides ; Dryander in Transact. Linn. Soc. III, 43, t. 11 (1797)	—	—	T.	—	N.S.W.	—	—	B.fl.VII,720	M.fr.V,118	...	N.Z.
L. microphylla ; Swartz, synops. filic. 120 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,721	M.fr.V,119.		
L. incisa ; Prentice in Trim. Journ. of Bot. 295 (1873)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,721	...	...	N.Z.
L. Fraseri ; Hooker, spec. filic. I, 221, t. 70 (1846)	—	—	—	—	—	Q.	—	B.fl.VII,721	M.fr.V,118.		
L. ensifolia ; Swartz, synops. filic. 118 (1806)	—	—	—	—	—	Q.	N.A.	B.fl.VII,721	M.fr.V,118	...	As., Afr., P.
L. lanuginosa ; Wallich, numerical list n. 154 (1828)	—	—	—	—	—	Q.	—	B.fl.VII,721	M.fr.V,118	...	As., Afr.
ADIANTUM ; Tournefort in 543, t. 317 (1700) from Hippocrates, Theophrastos, Dioscorides and Plinius.											
A. lunulatum ; N. Burman, Fl. Ind. 235 (1768)	—	—	—	—	—	Q.	N.A.	B.fl.VII,723	M.fr.V,119	...	As., Afr., Am.
A. Capillus veneris ; Linné, spec. plant. 1096 (1753)	—	—	—	—	—	Q.	—	B.fl.VII,723	...	...	E., As., Am., Afr.
A. Aethiopicum ; Linné, syst. nat. edit. decim. n. 15 (1759)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,724	M.fr.V,119	...	E., As., Afr., Am., P.,
A. formosum ; R. Brown, prodr. 155 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,724	M.fr.V,120; VIII,157.		N.Z. [N.Z.
A. affine ; Willdenow, spec. plant. V, 448 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,724	M.fr.V,119	...	N.Z.
A. diaphanum ; Blume, flora Javae, filic. 215 (1829)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,725	M.fr.VII,121	...	As., N.Z., P.
A. hispidulum ; Swartz, syn. filic. 124 & 321 (1806)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,725	M.fr.V,120	...	As., Afr., Am., P., N.Z.
CHEILANTHES ; Swartz, syn. filic. 126, t. 3 (1806). (Notholaena, Nothochlaena.)											
C. Prenticei ; Luerssen in Uhlw. Bot. Centralblatt IX, 440 (1882)	—	—	—	—	—	Q.	N.A.	—			
C. Clelandi ; F. v. M. & Tate in proceed. R.S. of S.A. X, 81 (1887)	—	S.A.	—	—	—	—	—	—			
C. pumilio ; F. v. M., Cens. 138 (1882) from R. Brown (1810)	—	—	—	—	—	Q.	—	B.fl.VII,773	M.fr.VIII,175.		
C. fragillima ; F. v. M., fragm. V, 123 (1866)	—	—	—	—	—	Q.	N.A.	B.fl.VII,774	M.fr.V,123.		
C. vellea ; F. v. M., fragm. V, 123 (1866)	W.A.	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,773	M.fr.V,123; VIII,176.		E., As., Afr.
C. distans ; A. Braun, index sem. hort. Berol. (1859)	W.A.	S.A.	—	V.	N.S.W.	Q.	—	B.fl.VII,774	M.fr.V,122; VIII,176.		P., N.Z.
C. tenuifolia ; Swartz, syn. filic. 126 (1806)	W.A.	S.A.	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,726	M.fr.V,122	...	As., N.Z., P.
PTERIS ; Linné, syst. nat. 9 (1735) ; gen. plant. 322 (1737). (Pellaea, Cheiloplecton, Platyloma, Litobrochia, Pteridium.)											
P. concolor ; Langsdorf & Fischer, icon. filic. t. 21 (1818)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,728	M.fr.V,124	...	As., Afr., Am., P.
P. paradoxa ; Baker in Benth. Fl. Austr. VII, 729 (1878)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,729			
P. falcata ; R. Brown, prodr. 154 (1810)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,729	M.fr.V,123	...	As., N.Z.
P. rotundifolia ; G. Forster, florul. ins. Austr. prodr. 79 (1786)	—	—	—	—	—	Q.	—	B.fl.VII,730	...	...	N.Z.
P. longifolia ; Linné, spec. plant. 1074 (1753)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,730	M.fr.V,126	...	E., As., Afr., Am., P.

P. ensiformis ; N. Burmann, Fl. Ind. 230 (1768)	—	—	—	—	—	Q.	—	B.fl.VII,730	M.fr.V,123			As., P.		
P. umbrosa ; R. Brown, prodr. 154 (1810)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,730	M.fr.V,126.					
P. biaurita ; Linné, spec. plant. 1076 (1753)	—	—	—	—	—	Q.	—	B.fl.VII,731	M.fr.V,125			As., Afr., Am., P., N.Z.		
P. arguta ; Aiton, hort. Kew. III, 458 (1789)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,731	M.fr.V,126			As., Afr., Am., P., N.Z.		
P. aquilina ; Linné, spec. plant. 1075 (1753)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,731	M.fr.V,126			E., As., Afr., Am., P., N.Z.		
P. incisa ; Thunberg, prodr. pl. Capens. 171 (1800)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,732	M.fr.V,124;	VIII,158.		As., Afr., Am., P.,		
P. marginata ; Bory. voy. dans les isles d'Afr. II, 192 (1804)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,733	M.fr.V,125			As., Afr., N.Z.		[N.Z.
P. comans ; G. Forster, florul. ins. Austr. prodr. 79 (1786)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,733	M.fr.V,125;	VII,121.		Am., P., N.Z.		

LOMARIA ; Willdenow in Magaz. der Ges. naturf. Freunde zu Berlin, III, 160 (1809). (Stegania, Plagiogyria.)

L. Patersoni ; Sprengel, sys. veg. IV, 62 (1827)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,734	M.fr.V,122;	VII,158.		As., P., N.Z.		
L. vulcanica ; Blume, fl. Javae, filic. 202 (1836)	—	—	T.	—	—	Q.	—	B.fl.VII,734	M.fr.V,121		...	As., P., N.Z.		
L. discolor ; Willdenow, spec. plant. V, 293 (1810)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,735	M.fr.V,121		...	N.Z.		
L. lanceolata ; Sprengel, syst. veg. IV, 62 (1827)	—	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,735	M.fr.V,121		...	P., N.Z.		
L. attenuata ; Willdenow, spec. plant. V, 290 (1810)	—	—	—	—	N.S.W.	—	—	B.fl.VII,736	..		...	Afr., Am., P.		
L. alpina ; Sprengel, syst. veg. IV, 62 (1827)...	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,736	M.fr.V,121		...	Am., N.Z.		
L. fluviatilis ; Sprengel, syst. veg. IV, 65 (1827)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,736	M.fr.V,121		...	N.Z.		
L. Fullagari ; F. v. M., fragm. VIII, 157 (1874)	—	—	—	—	N.S.W.	—	—	B.fl.VII,737	M.fr.VIII,157.					
L. Capensis ; Willdenow, spec. plant. V, 291 (1810)...	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,737	M.fr.V,121		...	As., Afr., Am., P., N.Z.		
L. euphlebia ; Kunze in Mohl & Schlecht. Bot. Zeit. VI, 521 (1849)	—	—	—	—	—	Q.	—	B.fl.VII,738	M.fr.VI,124		...	As.		

BLECHNUM ; Linné, spec. plant. 1077 (1753).

B. cartilagineum ; Swartz, syn. filic. 114 & 312 (1806)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,738	M.fr.V,120.					
B. laevigatum ; Cavanilles, descripcion de las plantas 650 (1802)	—	—	—	—	N.S.W.	—	—	B.fl.VII,739	M.fr.V,120.					
B. serrulatum ; C. Richard in Annal. du Mus. d'Hist. nat. Par. I, 114 (1802) ...	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,739	M.fr.V,120		...	As., Am., P		
B. orientale ; Linné, syst. nat. ed. decim. (1759)	—	—	—	—	—	Q.	N.A.	B.fl.VII,740	—		...	As., P.		

MONOGRAMMA ; Commerçon in Schkuhr's Kryptog. Gew. 82, t. 87 (1809). (Monogramme, Vaginularia, Pleurogramme, Diclidopteris.)

M. Junghuhnii ; Hooker, spec. filic. V, 123, t. 289 (1864)	—	—	—	—	—	Q.	—	B.fl.VII,740	M.fr.VII,119		...	As., P.		

WOODWARDIA ; Smith in mém. de l'Acad. de Turin V, 411 (1791). (Doodia.)

W. aspera ; Mettenius, fil. hort. bot. Lips. 65 (1856)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,740	M.fr.V,130.					
W. caudata ; Cavanilles, descripcion de las pl. n. 653 (1802)	—	—	T.	V.	N.S.W.	Q.	N.A.	B.fl.VII,742	M.fr.V,129;	VIII,158.		P., N.Z.		

ASPLENIUM ; Linné, gen. plant. 322 (1737) from C. Bauhin (1651). (Scolopendrium, Allantodea, Diplazium, Callipteris, Anisogonium, Thamnopteris, Neottopteris, Darea, Caenopteris, Athyrium, Diplora, Asplenum.)

A. Nidus ; Linné, spec. plant. 1079 (1753)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,744	M.fr.V,130		...	As., Afr., P.		
A. Robinsonii ; F. v. M. in Britten's Journ. Bot. XXII, 289 (1884)	—	—	—	—	N.S.W.	—	—	—						
A. simplicifrons ; F. v. M., fragm. V, 74 (1865)	—	—	—	—	—	Q.	—	B.fl.VII,744	M.fr.V,74,130;	VIII,158.				
A. attenuatum ; R. Brown, prodr. 150 (1810)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,745	M.fr.V,130.					[N.Z.
A. Trichomanes ; Linné, spec. plant. 1080 (1753)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,745	M.fr.V,131		...	E., As., Afr., Am., P.,		
A. flabellifolium ; Cavanilles, descripcion de las pl. n. 636 (1802)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,745	M.fr.V,131		...	N.Z.		
A. paleaceum ; R. Brown, prodr. 150 (1810)	—	—	—	—	—	Q.	—	B.fl.VII,746	M.fr.V,131;	VIII,158.				
A. falcatum ; Lamarck, Encycl. méth. I, 303 (1783)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,746	M.fr.V,131;	VII,156.		As., Afr., P., N.Z.		
A. Hookeranium ; Colenso in Journ. of Nat. Sc. Tasm. II, 169 (1842)	—	—	—	V.	N.S.W.	—	—	B.fl.VII,747	...		...	N.Z.		
A. furcatum ; Thunberg, prodr. pl. Cap. 172 (1800)	W.A.	S.A.	—	V.	N.S.W.	—	—	B.fl.VII,748	M.fr.V,131		...	As., Afr., Am., P.		
A. laserpitiifolium ; Lamarck, Encycl. méth. I, 310 (1783)...	—	—	—	—	—	Q.	—	B.fl.VII,749	M.fr.V,131		...	As., P.		[N.Z.
A. marinum ; Linné, spec. plant. 1081 (1753)...	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,747	M.fr.V,132		..	E., As., Afr., Am., P.,		
A. bulbiferum ; G. Forster, floral. ins. Austr. prodr. 433 (1786)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,749	M.fr.V,132		...	As., Afr., Am., P., N Z.		
A. pteridoides ; Baker, syn. filic. sec. edit. 488 (1874)	—	—	—	—	N.S.W.	—	—	B.fl.VII,749						
A. umbrosum ; J. Smith in Hook. Lond. Journ. IV, 174 (1842)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,749	M.fr.V,132		...	As., Afr., P., N.Z.		
A. pallidum ; Blume, fl. Javae, filic. 776 (1836)	—	—	—	—	—	Q.	—	—	M.fr.XI,131		..	As., P.		
A. sylvaticum ; Presl, reliq. Haenk. I, 42 (1830)	—	—	—	—	—	Q.	—	B.fl.VII,750	M.fr.V,133		...	As., Afr., P.		

A. maximum ; D. Don, prodr. fl. Nepal. 8 (1825)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,751		...	As.
A. polypodoides ; Mettenius, fil. hort. Lips. 78 (1856)	—	—	—	—	—	Q.	—	B.fl.VII,751	M.fr.V,132	...	As.
A. melanochlamys ; Hooker, spec. filic. III, 259 (1860)	—	—	—	—	N.S.W.	—	—	B.fl.VII,751			
A. decussatum ; Swartz, spec. fil. 76 & 260 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,751	...	...	As., Afr., P.
A. diversifolium ; Cunningham in Loud. hort. Brit. suppl. 581 (1832)	—	—	—	—	N.S.W.	—	—	—			
CYSTOPTERIS ; Bernhardi in Schrader's neuem Journ. II, 26 (1806).											
C. fragilis ; Bernhardi in Schrad. Journ. I, 26 (1806)	—	—	T.	—	—	—	—	B.fl.VII,752	M.fr.V,118	...	E., As., Afr., Am., P. N.Z.
ASPIDIUM ; Swartz in Schrader's Journ. II, 429 (1800). (Nephrodium, Nephrolepis, Polystichum partly, 1797, Lastraea, Sagenia, Oleandra.)											
A. cordifolium ; Swartz, syn. filic. 45 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,754	M.fr.V,136	...	As., Afr., Am., P., N.Z.
A. exaltatum ; Swartz, syn. filic. 45 (1806)	—	—	—	—	—	Q.	N.A.	B.fl.VII,754	M.fr.V,136	...	As., Afr., Am.
A. ramosum ; Palisot, Fl. d'Oware et de Benin II, 53, t. 91 (1807)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,754	M.fr.V,135	...	As., Afr., P.
A. unitum ; Swartz, syn. filic. 47 (1806)	W.A.	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,755	M.fr.V,137; VIII,158.		As., Afr.
A. pteroides ; Swartz, syn. filic. 49 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,755	...	...	As., P.
A. molle ; Swartz, syn. filic. 49 (1806)	—	S.A.	—	V.	N.S.W.	Q.	N.A.	B.fl.VII,756	M.fr.V,135	...	As., Afr., P., N.Z.
A. truncatum ; Gaudichaud in Voy. Freyc. Bot. 332, t. 10 (1826)...	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,756	M.fr.V,135	...	As., P.
A. confluens ; Mettenius in Schlecht. Linnaea XXXVI, 125 (1863)	—	—	—	—	—	Q.	—	B.fl.VII,757	M.fr.V,133; VIII,158.		
A. aculeatum ; Swartz in Schrad. Journ. II, 37 (1800)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,757	M.fr.V,134	...	E., As., Afr., Am., P., N.Z.
A. aristatum ; Swartz, syn. filic. 53 & 253 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,757	M.fr.V,134	...	As., Afr., P., N.Z.
A. coriaceum ; Swartz, syn. filic. 57 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,758	M.fr.V,134	...	As., Afr., Am., N.Z.
A. decompositum ; Sprengel, syst. IV, 109 (1827)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,758	M.fr.V,136	...	P., N.Z.
A. tenerum ; Sprengel, syst. IV, 109 (1827)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,759			
A. uliginosum ; Kunze in Schlecht. Linnaea XX, 6 (1847)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,759	M.fr.V,133	...	As., P.
A. hispidum ; Swartz, syn. filic. 56 (1806)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,760	M.fr.V,133	...	N.Z.
MENISCIUM ; Schreber, gen. plant. II, 757 (1791).											
M. triphyllum ; Swartz, syn. filic. 19 & 206 (1806)	—	—	—	—	—	Q.	—	—	...	...	As.
POLYPODIUM ; Tournefort, inst. 540, t. 316 (1700), from Theophrastos, Dioscorides and Plinius. (Niphobolus, Goniophlebium, Goniopteris, Phlebodium, Phegopteris, Phymatodes, Pleopeltis, Drynaria, Dictyopteris, Arthropteris, Xiphopteris, Meniscium partly ; from Dodoens, l'Ecluse, Bauhin, Morison, Ray, Plumier and particularly Petiver as Polypodium.)											
P. Australe ; Mettenius, filic. hort. bot. Lips. 36 (1856)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,762	M.fr.V,127	...	Am., N.Z.
P. Hookeri ; Brackenridge, Bot. Wilk. Exped. Filic. 4 (1854)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,763	...	...	As., P.
P. contiguum ; Brackenridge, Bot. Wilk. Exped. Filic. 6, t. 2 (1854)	—	—	—	—	—	Q.	—	B.fl.VII,763	M.fr.V,127	...	P.
P. grammitidis ; R. Brown, prodr. 147 (1810)	—	—	T.	V.	N.S.W.	—	—	B.fl.VII,764	M.fr.V,127	...	N.Z.
P. tenellum ; G. Forster, florul. ins. Austr. prodr. 440 (1786)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,764	M.fr.V,126; VII,121.		P., N.Z.
P. proliferum ; Roxburgh in Wall. num. list 11 (1828)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,765	M.fr.V,128	...	As., Afr., P.
P. urophyllum ; Wallich, numerical list 11 (1828)	—	—	—	—	—	Q.	—	B.fl.VII,765	M.fr.IV,165; V,137; VIII,121.		As., P.
P. Hillii ; Baker in Hooker, syn. filic. sec. edit. 505 (1874)...	—	—	—	—	—	Q]	—	B.fl.VII,766			
P. poecilophlebium ; Hooker, spec. filic. V, 14 (1863)	—	—	—	—	—	Q.	—	B.fl.VII,766	M.fr.V,127.		
P. serpens ; G. Forster, florul. insul. Austr. prodr. 81 (1786)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,767	M.fr.V,129	...	P., N.Z.
P. confluens ; R. Brown, prodr. 146 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,767	...	...	P.
P. acrostichoides ; G. Forster, florul. insul. Austr. prodr. 81 (1786)	—	—	—	—	—	Q.	—	B.fl.VII,767	M.fr.V,129	...	As., P.
P. attenuatum ; R. Brown, prodr. 146 (1810)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,768	M.fr.V,128; V,121.		P.
P. simplicissimum ; F. v. M. in Hooker & Baker, syn. filic. sec. edit. 513 (1874)	—	—	—	—	—	Q.	—	B.fl.VII,768	M.fr.VII,120,156.		
P. fusco-pilosum ; F. v. M. & Baker in Britt. journ. bot. XXV, 163 (1887) ...	—	—	—	—	—	Q.	—	—			
P. nigrescens ; Blume, Fl. Javae, filic. 161, t. 70 (1829)	—	—	—	—	—	Q.	—	B.fl.VII,769	...	...	As., Afr., Am., P.
P. phymatodes ; Linné, mant. pl. alt. 306 (1771)	—	—	—	—	—	Q.	N.A.	B.fl.VII,769	...	...	As., Afr., P., N.Z.
P. pustulatum ; G. Forster, florul. ins. Austr. pr. 81 (1786)	—	—	T.	V.	N.S.W.	Q.	—	B.fl.VII,769	M.fr.V,127	...	N.Z.
P. scandens ; G. Forster, florul. ins. Austr. pr. 81 (1786)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,770	M.fr.V,128	...	P., N.Z.
P. verrucosum ; Wallich, numerical list II (1828)	—	—	—	—	—	Q.	—	B.fl.VII,770	...	...	As., P., N.Z.

P. subauriculatum; Blume, Fl. Javae, filic. 177, t. 83 (1829)	—	—	—	—	—	Q.	—	B.fl.VII,770	M.fr.V,128	...	As., P.
P. rigidulum; Swartz, synops. filic. 38 & 320 (1806)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,771	...	...	As., P.
P. quercifolium; Linné, spec. plant. 1087 (1753)	—	—	—	—	—	Q.	N.A.	B.fl.VII,772	M.fr.V,127	...	As., P.
P. irioides; Poiret, Encycl. méth. V, 513 (1804)	—	—	—	—	—	Q.	—	B.fl.VII,771	M.fr.V,127	...	As., Afr., P.
P. punctatum; Thunberg, fl. Japon. 337 (1784)	—	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,764	M.fr.V,129	...	As., Afr., N.Z.
HYPOLEPIS; Bernhardi in Schrader, neu. Journ. I, 34 (1806).											
H. tenuifolia; Bernhardi in Schrad. Journ. II, 24 (1800)	—	—	—	V.	N.S.W.	Q.	—	B.fl.VII,726	M.fr.V,129	...	As., P., N.Z.
GRAMMITIS; Swartz in Schrader, Journ. II, 3 & 17 (1800). (Gymnogramme, Selliguea.)											
G. Reynoldsii; F. v. M. in Benth. Fl. Austr. VII, 775 (1878)	—	S.A.	—	—	—	Q.	—	B.fl.VII,775	M.fr.VIII,175.		
G. Muelleri; Hooker in F. v. M., fragm. V, 138 (1866)	—	—	—	—	—	Q.	—	B.fl.VII,775	M.fr.VI,124; VII,121.		
G. rutifolia; R. Brown, prodr. 146 (1810)	W.A.	S.A.	T.	V.	N.S.W.	Q.	—	B.fl.VII,775	M.fr.V,137	...	E., Am., N.Z.
G. leptophylla; Swartz, syn. filic. 23 & 218, t. 1 (1806)	W.A.	S.A.	T.	V.	N.S.W.	—	—	B.fl.VII,776	M.fr.V,137	...	E., As., Afr., Am., P., N.Z.
G. pinnata; F. v. M., fragm. VI, 124 (1868)	—	—	—	—	—	Q.	—	B.fl.VII,776	M.fr.VI,124	...	As., P.
G. ampla; F. v. M., fragm. V, 188 (1866)	—	—	—	—	—	Q.	—	B.fl.VII,777	M.fr.V,188.		
G. Sayeri; F. v. M. & Baker in Britt. journ. bot. XXV, 163 (1887	—	—	—	—	—	Q.	—	—			
ANTROPHYUM; Kaulfuss, enum. filic. 197 (1824).											
A. reticulatum; Kaulfuss, enum. filic. 197 (1824)	—	—	—	—	—	Q.	—	B.fl.VII,777	M.fr.V,138	...	As., P.
ACROSTICHUM; Linné, gen. plant. 322 (1737), indicative. (Elaphoglossum, Stenochlaena, Lomariopsis, Hymenolepis, Gymnopteris, Chrysodium.)											
A. conforme; Swartz, syn. filic. 10 & 192, t. 1 (1806)	—	—	—	—	—	Q.	—	B.fl.VII,778	M.fr.V,138	...	As., Afr., Am.
A. scandens; J. Smith in Hook. spec. fil. V, 249 (1864)	—	—	—	—	—	Q.	N.A.	B.fl.VII,778	M.fr.VI,124	...	As., P.
A. sorbifolium; Linné, spec. plant. 1069 (1753)	—	—	—	—	—	Q.	—	B.fl.VII,779	M.fr.VII,119.		As., Afr., Am.
A. neglectum; Bailey in proc. Linn. Soc. of N.S.W. V, 32 (1880)	—	—	—	—	—	Q.	—	—	M.fr.XI,131.		
A. repandum; Blume, Fl. Javae 39, t. 14 & 15 (1829)	—	—	—	—	—	Q.	—	B.fl.VII,779	M.fr.V,138	...	As., P.
A. aureum; Linné, spec. plant. 1069 (1753)	—	—	—	—	N.S.W.	Q.	N.A.	B.fl.VII,779	M.fr.V,138; VII,120.		As., Afr., Am., P.
A. spicatum; Linné, fil. suppl. pl. 444 (1781)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,780	M.fr.V,138	...	As., Afr., P.
A. pteroides; R. Brown, prodr. 145 (1810)	—	—	—	—	—	Q.	N.A.	B.fl.VII,780	M.fr.X,76.		
A. Taylori; Bailey in trans. Accl. Soc. of Queensl. (1884)	—	—	—	—	—	Q.	—	—			
PLATYCERIUM; Desvaux in Mém. de la Soc. Linn. de Paris VI, 213 (1827).											
P. alcicorne; Desvaux in Mém. Soc. Linn. Par. VI, 213 (1827)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,780	M.fr.V,139	...	Afr.
P. grande; J. Smith in Hooker, Lond. Journ. III, 402 (1841)	—	—	—	—	N.S.W.	Q.	—	B.fl.VII,781	M.fr.V,139	...	As.

SEQUENCE OF ORDERS ACCORDING TO SYSTEMATIC AFFINITY, WITH INDICATION OF THEIR NUMBERS OF SPECIES.

Dilleniaceae ...	95	Rutaceae ...	190	Phytolacceae ...	11	Olacineae ...	15	Convolvulaceae...	70	Haemodoraceae	66
Ranunculaceae ...	17	Simarubeae ...	7	Nyctagineae ...	6	Balanophoreae ...	1	Hydrophylleae ...	2	Amaryllideae ...	21
Ceratophylleae ...	1	Zygophylleae ...	22	Thymeleae ...	75	Santalaceae ...	43	Solanaceae ...	79	Dioscorideae ...	4
Nymphaeaceae ...	5	Lineae	4	Leguminosae ..	1065	Loranthaceae ...	27	Scrophularinae...	80	Roxburghiaceae	1
Piperaceae ...	10	Geraniaceae ...	8	Connaraceae ...	2	Proteaceae ...	597	Orobancheae ...	1	Liliaceae ...	161
Magnoliaceae ...	4	Malvaceae ...	110	Rosaceae ...	17	Cornaceae ...	1	Lentibularinae ...	25	Palmae	25
Anonaceae ...	18	Sterculiaceae ...	125	Saxifrageae ...	36	Rubiaceae ...	127	Podostemoneae...	1	Nipaceae ...	1
Monimiaceae ...	16	Tiliaceae ...	56	Nepenthaceae ...	2	Caprifoliaceae ...	2	Gesneriaceae ...	4	Pandaneae ...	11
Myristiceae ...	1	Euphorbiaceae ...	224	Aristolochieae ...	5	Passifloreae ...	6	Bignoniaceae ...	7	Aroideae ...	10
Lauraceae ...	37	Urticaceae ...	65	Crassulaceae ...	6	Cucurbitaceae ...	27	Pedalinae ...	3	Typhaceae ...	2
Menispermeae ...	17	Cupuliferae ...	4	Hamamelideae ...	1	Compositae ...	539	Acanthaceae ...	30	Lemnaceae ...	6
Papaveraceae ...	1	Casuarineae ...	24	Onagreae ...	5	Campanulaceae	35	Labiatae ...	125	Fluviales ...	36
Capparideae ...	24	Celastrinae ...	18	Salicarieae ...	19	Candolleaceae ...	97	Verbenaceae ...	82	Alismaceae ...	6
Cruciferae ...	54	Meliaceae ...	36	Stackhousieae ...	13	Goodeniaceae ...	220	Myoporinae ...	76	Pontederiaceae ...	1
Violaceae ...	13	Sapindaceae ...	100	Halorageae ...	58	Gentianeae ...	24	Asperifoliae ...	52	Philhydreae ...	4
Flacourtieae ...	7	Malpighiaceae ...	2	Callitrichinae ...	2	Loganiaceae ...	52	Ericaceae ...	7	Commelineae ...	19
Samydaceae ...	4	Burseraceae ...	3	Rhizophoreae ...	7	Plantagineae ...	4	Epacrideae ...	275	Xyrideae ...	9
Pittosporeae ...	40	Anacardiaceae ...	9	Combretaceae ...	27	Primulaceae ...	6	Coniferae ...	29	Flagellariaceae ...	1
Droseraceae ...	46	Frankeniaceae ...	7	Myrtaceae ...	663	Myrsinaceae ...	12	Cycadeae ...	14	Junceae... ...	16
Elatineae ...	4	Plumbagineae . .	4	Melastomaceae ...	7	Sapotaceae ...	19	Scitamineae ...	11	Eriocauleae ...	18
Hypericinae ...	1	Portulaceae ...	32	Rhamnaceae ...	90	Ebenaceae ...	15	Orchideae ...	271	Restiaceae ...	93
Ternstroemiaceae	1	Caryophylleae ...	26	Viniferae ...	17	Aquifoliaceae ...	2	Apostasiaceae ...	1	Cyperaceae ...	380
Guttiferae ...	3	Amarantaceae ...	102	Loeaceae ...	2	Styraceae ...	2	Burmanniaceae	2	Gramineae ...	345
Polygaleae ...	32	Salsolaceae ...	111	Araliaceae ...	22	Jasmineae ...	21	Irideae	24	Rhizospermae ...	11
Tremandreae ...	17	Ficoideae ...	28	Umbelliferae ...	106	Apocyneae ...	47	Hydrocharideae	9	Lycopodinae ...	21
Ochnaceae ...	1	Polygonaceae ...	25	Elaeagneae ...	1	Asclepiadeae ...	61	Taccaceae ...	1	Filices	212

SEQUENCE OF ORDERS ACCORDING TO PREDOMINANCE, WITH INDICATION OF THEIR NUMBERS OF SPECIES.

Leguminosae	1065	Scrophularinae	80	Ficoideae	28	Viniferae	17	Ericaceae	7	Callitrichinae	2
Myrtaceae	663	Solanaceae	79	Combretaceae	27	Rosaceae	17	Nyctagineae	6	Loeaceae	2
Proteaceae	597	Myoporinae	76	Cucurbitaceae	27	Monimiaceae	16	Crassulaceae	6	Caprifoliaceae	2
Compositae	539	Thymeleae	75	Loranthaceae	27	Junceae	16	Passifloreae	6	Aquifoliaceae	2
Cyperaceae	380	Convolvulaceae	70	Caryophylleae	26	Olacineae	15	Primulaceae	6	Styraceae	2
Gramineae	345	Haemodoraceae	66	Polygonaceae	25	Ebenaceae	15	Lemnaceae	6	Hydrophylleae	2
Epacrideae	275	Urticaceae	65	Lentibularinae	25	Cycadeae	14	Alismaceae	6	Burmanniaceae	2
Orchideae	271	Asclepiadeae	61	Palmae	25	Violaceae	13	Nymphaeaceae	5	Typhaceae	2
Euphorbiaceae	224	Halorageae	58	Capparideae	24	Stackhousieae	13	Aristolochieae	5	Ceratophylleae	1
Goodeniaceae	220	Tiliaceae	56	Casuarineae	24	Myrsinaceae	12	Onagreae	5	Myristiceae	1
Filices	212	Cruciferae	54	Gentianeae	24	Phytolacceae	11	Magnoliaceae	4	Papaveraceae	1
Rutaceae	190	Loganiaceae	52	Irideae	24	Scitamineae	11	Samydaceae	4	Ternstroemiaceae	1
Liliaceae	161	Asperifoliae	52	Zygophylleae	22	Pandaneae	11	Elatineae	4	Hypericinae	1
Rubiaceae	127	Apocyneae	47	Araliaceae	22	Rhizospermae	11	Lineae	4	Ochnaceae	1
Sterculiaceae	125	Droseraceae	46	Jasmineae	21	Piperaceae	10	Cupuliferae	4	Podostemoneae	1
Labiatae	125	Santalaceae	43	Amaryllideae	21	Aroideae	10	Plumbagineae	4	Balanophoreae	1
Salsolaceae	111	Pittosporeae	40	Lycopodinae	21	Anacardiaceae	9	Plantagineae	4	Hamamelideae	1
Malvaceae	110	Lauraceae	37	Salicarieae	19	Hydrocharideae	9	Gesneriaceae	4	Elaeagneae	1
Umbelliferae	106	Meliaceae	36	Sapotaceae	19	Xyrideae	9	Dioscorideae	4	Cornaceae	1
Amarantaceae	102	Saxifrageae	36	Commelineae	19	Geraniaceae	8	Philhydreae	4	Orobancheae	1
Sapindaceae	100	Fluviales	36	Anonaceae	18	Flacourtieae	7	Guttiferae	3	Apostasiaceae	1
Candolleaceae	97	Campanulaceae	35	Celastrinae	18	Simarubeae	7	Burseraceae	3	Taccaceae	1
Dilleniaceae	95	Polygaleae	32	Ericoauleae	18	Frankeniaceae	7	Pedalinae	3	Roxburghiaceae	1
Restiaceae	93	Portulaceae	32	Ranunculaceae	17	Rhizophoreae	7	Malpighiaceae	2	Nipaceae	1
Rhamnaceae	90	Acanthaceae	30	Menispermeae	17	Melastomaceae	7	Connaraceae	2	Pontederiaceae	1
Verbenaceae	82	Coniferae	29	Tremandreae	17	Bignoniaceae	7	Nepenthaceae	2	Flagellariaceae	1

Total Number of Orders of Vasculares, 156; of Genera, 1409; of Species, 8839. Of these occur in W.A., 3560: in S.A., 1892; in T., 1029; in V., 1894; in N.S.W., 3251; in Q., 3753; in N.A., 1956. Admitted from Flora Australiensis, 7814; additional species recorded in Census, 1025. Proportion of distribution through the respective Colonial territories:—In W.A., 40.3 per cent.; in S.A., 21.4 per cent.; in T., 11.6 per cent.; in V., 21.4 per cent.; in N.S.W., 36 8 per cent.; in Q., 42.5 per cent.; in N.A., 22.1 per cent. Of the total number 7501 are endemic in Continental Australia and Tasmania; therefore 1338 or 15.1 per cent. extend also to other countries. Of these 160 are found in Europe; 1032 in Asia; 515 in Africa; 315 in America; 558 in Polynesia; 291 in New Zealand.

Species added since the publication of the first Census 193, irrespective of some systematic changes. Geographic entries added since 1150.

Chronologic quotations of original authors of such species, as are transferred to other genera, will be given in the next supplement for which also many species additional for Australia have already been recorded in manuscript, while these pages were going through the press.

In all instances immigrated plants are excluded.

INDEX OF ORDERS AND GENERA.

Abbottia ... 125
Abelmoschus... 25
Abildgaardia... 211
Abroma ... 27
Abrophyllum .. 82
Abrotanella ... 142
Abrus... ... 72
Abutilaea ... 24
Abutilon ... 24
Acacia ... 74
Acaena ... 82
Acalypha ... 36
Acanthaceae ... 167
Acanthocarpus ... 200
Acanthocladium ... 136
Acanthus ... 167
Achilleopsis ... 27
Achnophora ... 131
Achras ... 154
Achromolaena ... 138
Achrysum ... 140
Achyranthes ... 47
Acianthus ... 191
Acicalyptus ... 102
Aciphylla ... 108
Ackama ... 82
Acmene ... 101
Acomis ... 138
Acrachne ... 225
Acradenia ... 20
Acroclinium ... 136
Acronychia ... 21
Acrophyllum... 82
Acrostichum... 234
Acrotriche ... 177
Actephila ... 34
Actinocarpus... 205
Actinodium ... 87
Actinopappus ... 138
Actinostachys ... 229
Actinostigma ... 18
Actinostrobus ... 184
Actinotus ... 108
Adansonia ... 26
Adeliopsis ... 7
Adenanthera... 74
Adenanthos ... 113
Adenochilus ... 191
Adenosma ... 163
Adenostemma ... 130
Adenostephanus ... 116
Adiantum ... 231
Adrastaea ... 1
Adriana ... 36
Aegialinites ... 45
Aegialitis ... 45
Aegiceras ... 154
Aegopogon ... 223
Aera ... 224
Aeschynomene ... 69
Afzelia ... 74
Agapetes ... 177
Agastachys ... 114
Agathis ... 183
Agati ... 68
Ageratum ... 130
Aglaia ... 16
Aglaiopsis ... 16
Agnostus ... 122
Agonis ... 92
Agrimonia ... 81
Agropyron ... 227
Agrostis ... 224
Agrostocrinum ... 192
Ailantus ... 22
Aillya... 149
Aira ... 224
Aizoon ... 52
Ajuga... 171
Akania ... 42
Alania... 199
Albizzia ... 80
Alchemilla ... 81
Alchornea ... 36
Aldrovanda ... 12
Alectryon ... 42
Alepyrum ... 207
Aleurites ... 36
Alisma ... 205
Alismaceae ... 205
Allania ... 199
Allantodea ... 232
Allophylus ... 41
Alocasia ... 203
Alopecurus ... 222
Alphitonia ... 102
Alpinia ... 185
Alsomitra ... 129
Alsophila ... 230
Alstonia ... 157
Alternanthera ... 47
Althenia ... 204
Altingia ... 183
Alysicarpus ... 70
Alyssopsis ... 9
Alyssum ... 9
Alyxia ... 156
Amanoa ... 35
Amarantaceae ... 47
Amarantus ... 49
Amaryllideae ... 195
Amblysperma ... 143
Ambrina ... 50
Ammannia ... 84
Ammobium ... 138
Amomum ... 185
Amoora ... 16
Amorphophallus ... 203
Amorphospermum ... 154
Amperea ... 33
Amphibolis ... 204
Amphibromus ... 225
Amphicosmia ... 230
Amphipogon... 223
Amphodus ... 70
Anacardiaceae ... 44
Anadenia ... 117
Anagallis ... 153
Ansrthria ... 208
Ancana ... 5
Ancistrostigma ... 52
Ancistrum ... 82
Andersonia ... 182
Andrachne ... 33
Andrewsia ... 175
Andriapetalum ... 116
Andropogon ... 222
Androstemma ... 194
Androstoma ... 177
Aneilema ... 205
Anemone ... 3
Angianthus ... 139
Angiopteris ... 229
Angophora ... 97
Anguillaria ... 198
Anigozanthos ... 195
Anisacantha ... 51
Anisandra ... 170
Aniseia ... 159
Anisogonium... 232
Anisolepis ... 136
Anisomeles ... 169
Anisopogon ... 225
Anisotome ... 108
Anodopetalum ... 82
Anoectochilus .. 188
Anonaceae ... 4
Anopterus ... 82
Anosporum ... 210
Anotis ... 125
Antennaria ... 134
Antheidosorus ... 139
Anthericum ... 198
Anthistiria ... 222
Anthobolus ... 110
Anthocerastes ... 138
Anthocercis ... 162
Anthosachne... 227
Anthotium ... 147
Anthotroche ... 162
Antiaris ... 38
Anticoryne ... 91
Antidesma ... 33
Antirrhoea ... 126
Antrophyum ... 234
Aotus ... 62
Apera ... 224
Aphananthe ... 37
Aphanopetalum ... 82
Aphelia ... 207
Apium ... 108
Aplotaxis ... 143
Apluda ... 222
Apocyneae ... 156
Apodytes ... 109
Apolochlamys ... 138
Aponogeton ... 204
Apophyllum ... 9
Aporosa ... 35
Apostasia ... 192
Apostasiaceae ... 192
Aquifoliaceae ... 155
Arabis ... 9
Araliaceae ... 105
Araucaria ... 183
Archeria ... 181
Archidendron ... 80
Archontophoenix ... 202
Ardisia ... 154, 177
Areca ... 202
Arenaria ... 46
Argophyllum ... 82
Argyrocome ... 136
Argyrodendron ... 27
Argyrophanes ... 136
Argyroglottis ... 136
Aristida ... 223
Aristolochia ... 83
Aristolochieae ... 83
Aristotelia ... 30
Arnocrinum ... 200
Aroideae ... 203
Artanema ... 164
Arthratherum ... 223
Arthraxon ... 221
Arthrochilus ... 191
Arthrocnemum ... 52
Arthropodium ... 199
Arthropteris ... 233
Arthrostylis ... 217
Arthrotaxis ... 183
Arthrotrichum ... 48
Arum ... 203
Arundinella ... 220
Arundo ... 227
Aryteria ... 41
Asclepiadeae ... 157
Ascyrum ... 14
Asparagopsis... 197
Asparagus ... 197
Asperella ... 222
Asperifoliae ... 175
Asperula ... 128
Aspidium ... 233
Asplenium ... 232
Asprella ... 222
Astartea ... 92
Astelia ... 198
Aster ... 131
Asteridea ... 135
Asterochiton ... 29
Asterolasia ... 18
Asteromyrtus ... 94
Asterotrichon ... 24
Astraea ... 90
Astrebla ... 225
Astroloma ... 177
Astrotricha ... 105
Atalantia ... 21
Atalaya ... 41
Atelandra ... 170
Atherocephala ... 182
Atherosperma ... 5
Athrixia ... 135
Athrodactylis ... 202
Athrotaxis ... 183
Athyrium ... 232
Atkinsonia ... 111
Atriplex ... 49
Atylosia ... 71
Australina ... 39
Automachia ... 20
Avicennia ... 173
Aylmeyera ... 47
Azolla... 228
Azorella ... 107

Babbagia ... 51
Babingtonia ... 91
Backhousia ... 100
Bacularia ... 201
Baea ... 166
Baeckea ... 91
Baeobotrys ... 154
Balanophora ... 109
Balanophoreae .. 109
Balanops ... 39
Balantium ... 230
Balaustion ... 92
Balfouria ... 157
Baloghia ... 37
Bambusa ... 227
Banksia ... 55, 122
Barbaraea ... 9
Barklya ... 72
Barringtonia ... 102
Bartlingia ... 62, 200
Basilicum ... 168
Bassia ... 51
Batatas ... 159
Batratherum ... 221
Bauera ... 83
Bauhinia ... 73
Baumea ... 216
Baxteria ... 200
Bea ... 166
Beaufortia ... 96
Bedfordia ... 142
Beilschmiedla ... 6
Bellendena ... 114
Bellis ... 130
Benincasa ... 129
Bergia ... 14
Berrya ... 30
Bertolonia ... 175
Bertya ... 33

Beyeria 32
Beyeriopsis 32
Bidaria 158
Bidens 141
Bignoniaceae... ... 166
Billardiera 12
Billotia ... 92, 96
Biophytum 23
Bischoffia 35
Blackburnia 21
Blackwellia 11
Blainvillea 141
Blancoa 194
Blandfordia 197
Bleasdalea 116
Blechnum 232
Blennodia 9
Blennospora 140
Blepharanthemum ... 24
Blepharocarya ... 44
Blitum 50
Blumea 133
Blyxa 194
Bobea 126
Boea 166
Boehmeria 38
Boerhaavia 54
Bolbophyllum ... 186
Bombax 26
Bonnaya 164
Boronia 17
Borya 200
Bosistoa 20
Bossiaea 65
Bothryodendron ... 106
Botrychium 229
Bouchardatia ... 21
Bowenia 184
Brachychiton ... 26
Brachycome 130
Brachyloma 181
Brachynema... ... 82
Brachypterum ... 72
Brachysema 56
Brachyspatha ... 203
Brachystephium ... 130
Brackenridgea ... 17
Bradleia 34
Bramia 164
Brasenia 4
Brassaia 106
Brathys 14
Breweria 160
Breynia 35
Bridelia 35
Brizopyrum 226
Brizula 207
Brochosiphon ... 168
Brombya 21
Bromidium 224
Bromus 226
Bruncea 22
Bruguiera 86
Brunella 169
Brunonia 146
Bryonia 129
Bryonopsis 129
Buchanania 44
Buchnera 165
Buckinghamia ... 122
Buechnera 165
Bulbine 198
Bulbophyllum ... 186
Bulliarda 83
Bulweria 167
Bunnya 173
Burchardia 198
Burgesia 56
Burmannia 192
Burmanniaceae ... 192
Burnettia 191
Bursaria 12
Burseraceae 43
Burtonia 59
Busbeckea 8
Butomopsis 205
Byblis... 13
Byronia 155

Cabomba 4
Cadaba 8
Cadellia 22
Caenopteris 232
Caesalpina 72
Caesia... 199
Cajanus 71
Cakile... 10
Caladenia 191
Caladium 203
Calamus 201
Calandrinia 45
Calanthe 188
Caldasia 108
Caldesia 205
Caleana 191
Calectasia 200
Caletia 32
Caleya 191
Calliandra 80
Callicarpa 173
Callicoma 82
Callicystus 71
Callipteris 232
Callistachya 57
Callistemon 94
Callitriche 86
Callitrichinae ... 86
Callitris 184
Calocephalus... ... 140
Calochilus 189
Calogyne 149
Calomeria 138
Calonyction 159
Calopetalon 12
Calophyllum... ... 14
Calorophus 209
Calosciadium ... 108
Calostemma 196
Calostrophus... ... 209
Calothamnus... ... 96
Calotis 131
Calpurnia 72
Caltha... 3
Calycomis 82
Calycopeplus... ... 31
Calycothrix 89
Calyptranthes .. 102
Calyptrocalyx ... 202
Calyptrostegia ... 55
Calyptrostigma ... 32
Calystegia 160
Calythrix 89
Campanulaceae ... 143
Camphoromyrtus ... 91
Camptostemon ... 26
Campuleia 165
Campylanthera .. 12
Campynema 195
Cananga 5
Canarium 44
Canavalia 71
Candollea ... 1, 144
Candolleaceae ... 144
Canscora 152
Cansjera 109
Canthium 126
Capparideae 8
Capparis 8
Caprifoliaceae ... 128
Capsella 10
Carallia 86
Carapa 16
Carara 10
Cardamine 9
Cardiospermum ... 41
Cardwellia 122
Carex 217
Careya 102
Cargillia 155
Carissa 156
Carmichaelia... ... 68
Carnarvonia 122
Carpha 214
Carpodontos 83
Carronia 8
Cartonema 206
Carumbium 37
Carunostylis 189
Caryodaphne... ... 6
Caryophylleae ... 46
Caryospermum ... 40
Caryota 202
Casearia 11
Cassia... 73
Cassinia 138
Cassiniola 136
Cassytha 7
Castanospermum ... 72
Castanospora ... 41
Casuarina 39
Casuarineae 39
Catakidozamia ... 184
Cathartocarpus ... 73
Cathormion 80
Catosperma 147
Caturus 38
Caulinia ... 70, 203, 204
Caustis 217
Cedrela 16
Celastrinae 40
Celastrus 40
Celmisia 131
Celosia 49
Celtis 37
Cenarrhenes 114
Cenchrus 220
Centaurea 143
Centella 107
Centipeda 141
Centotheca 226
Centranthera ... 165
Centratherum ... 129
Centrolepis 207
Centropappus ... 142
Centunculus 153
Cephaelis 127
Cephalipterum ... 140
Cephalosorus... ... 139
Cephalotus 83
Ceranthea 156
Ceratogyne 142
Ceratopetalum ... 82
Ceratophylleae ... 3
Ceratophyllum ... 3
Ceratopteris 230
Cerbera 156
Cercodia 85
Ceriops 86
Ceropegia 159
Cesatia 106
Chaetanthus 209
Chaetaria 223
Chaetospora 214
Chalcas 21
Chamaelaucium ... 87
Chamaeraphis ... 218
Chamaescilla... ... 199
Chamaespherion ... 140
Chamaexeros... ... 200
Chapelliera 216
Charlwoodia 197
Cheilanthes 231
Cheiloplecton ... 231
Cheiranthera... ... 12
Cheiroloma 131
Chenolea 51
Chenopodina... ... 52
Chenopodium 50, 52
Cheynia 92
Chilocarpus 156
Chilodia 169
Chiloglottis 192
Chionachne 222
Chionanthus 156
Chithonanthus ... 74
Chlamydia 197
Chlamysporum ... 198
Chloanthes 172
Chloris 225
Chlorophytum ... 199
Choirizotheca ... 32
Chondrachne... ... 213
Choretrum 110
Choriceras 34
Chorilaena 20
Choripetalum ... 154
Chorizandra 213
Chorizema ... 57, 58
Chromochiton ... 138
Chrysobactron ... 198
Chrysocephalum ... 136
Chrysocoryne ... 139
Chrysodiscus... ... 135
Chrysodium 234
Chrysogonium ... 140
Chrysophyllum ... 154
Chrysopogon ... 222
Chrysostemon ... 32
Chryssorrhoea ... 88
Chthonocephalus ... 140
Chuncoa 87
Cibotium 230
Cinna 223
Cinnamomum ... 6
Cirrhopetalum ... 186
Cissampelos 7
Cissodendron ... 105
Cissus 104
Citriobatus 12
Citrus 22
Cladium 216
Cladodes 36
Claoxylon 36
Clausena 21
Claytonia 45
Cleisostoma 187
Cleistanthes 35
Cleistocalyx ... 101, 102
Clematis 3
Cleome 8
Clerodendrum ... 173
Clianthus 68
Clidanthera 69
Clinostigma 202
Clitoria 70
Cloezia 100
Coatesia 21
Coccoloba 53
Cocculus 7
Cochlospermum ... 11
Cocos 202
Codiaeum 37
Codonocarpus ... 54
Coelachne 224
Coelandria 186
Coelebogyne 36
Coelorrhachis ... 221
Coelospermum ... 126
Coffea 126
Coilocarpus 51
Coldenia 176
Coleanthera 181
Coleocoma 134
Coleostylis 146
Coleus 168
Colletia 104
Colmeiroa 82
Colobandra 170
Colobanthus 46
Colocasia 203
Colubrina 102
Columbea 183
Colymbea 183
Combretaceae ... 87
Comesperma 15
Commelina 205
Commelineae ... 205
Commerconia ... 27
Compositae 129
Conanthodium ... 136
Conchium 120
Coniferae 183
Coniogeton 44
Connaraceae 81
Conospermum ... 113
Conostephiopsis ... 181
Conostephium ... 181
Conostylis 194
Conothamnus ... 96
Convolvulaceae ... 159
Convolvulus... ... 160
Conyza 133
Coprosma 127
Corchorus 30
Cordia 177
Cordyline 197
Corethrostylis ... 29
Coridochloa 218
Cornaceae 124
Correa 20
Corybas 190
Corymbis 188
Corymborchis ... 188
Corynotheca 199
Coryospermum ... 40
Corypha 202
Corysanthes 190
Cosmelia 182
Costus 185
Cotula 141
Covellia 37
Crantzia 108
Crantziola 108
Craspedia 140
Crassula 83
Crassulaceae 83
Crepis... 143
Cressa... 161
Crinopodium... ... 84
Crinum 196
Crossolepis 139
Crossotoma 148
Crotalaria 66
Croton 36

Crowea 18
Cruciferae 9
Cryphia 169
Cryptandra 103
Cryptocarya 6
Cryptosema 56
Cryptostemon ... 87
Cryptostylia 189
Ctenosperma 141
Cucumia 129
Cucurbitaceae ... 128
Cudrania 38
Culcoa 197
Cupania 41
Cupuliferae 39
Curculigo 196
Curcuma 185
Cuscuta 161
Cuttsia 82
Cyanostegia 173
Cyanothamnus ... 17
Cyanotis 205
Cyathea 230
Cyathochaete ... 214
Cyathodes 177
Cyathopappus ... 139
Cyathopsis 177
Cyathula 49
Cybele 122
Cycadeae 184
Cycas 184
Cyclogyne 68
Cyclotheca 54
Cycnogeton 204
Cylicodaphne ... 6
Cylindrosorus ... 139
Cymbidium 187
Cymbonotus 143
Cyminosma 21
Cymodocea 204
Cynanchum ... 157, 158
Cynoctonum 157
Cynodon 225
Cynoglossum 175
Cynometra 74
Cyperaceae 210
Cyperus 210
Cyphanthera 162
Cyphokentia 202
Cypselocarpus ... 54
Cyrtostylis 191
Cystanthe 183
Cystopteria 233

Dacrydium 184
Dactyloctenium ... 225
Daemia 158
Dalbergia 72
Dallachya 102
Damasonium ... 193, 205
Dammara 183
Dampiera 146
Danthonia 225
Daphnandra 5
Daphne 56
Daphnobryon ... 56
Darea 232
Darlingia 122
Darwinia 87
Dasymalla 172
Dasypogon 201
Datura 162
Daucus 108
Davallia 231
Davidsonia 83
Daviesia 61
Decadon 85
Decalophium 87
Decaspermum ... 101
Decaspora 181
Decazesia 139
Deeringia 49
Delabechea 26
Delarbrea 105
Dendrobium 186
Denhamia 40
Denisonia 173
Deunstaedtia 230
Dentella 125
Deparia 230
Deplanchea 167
Depremesnilia ... 172
Derris 72
Deschampsia 224
Desmocladus 209
Desmodium 69
Desvauxia 207
Deyeuxia 224
Dianella 197
Diaspasis 147
Diastemanthe ... 220
Dicarpidium 27
Dicera 30
Dicerma 69
Dichelachne 223
Dichoglottis 46
Dichondra 161
Dichopetalum ... 107
Dichopogon 199
Dichosema 59
Dichrostachys ... 74
Dicksonia 230
Dicladanthera ... 168
Diclidopteris 232
Dicliptera 168
Dicrastylis 172
Dictyopteris 233
Didiscus 106
Didymanthus ... 51
Didymeria 20
Didymocarpus ... 166
Didymochaeta ... 224
Didymonema ... 216
Didymoplexis ... 188
Didymotheca ... 54
Digitaria 218
Dilivaria 167
Dillenia 1
Dilleniaceae 1
Dillwynia 64
Dimeria 221
Dimetopia 106
Dimorphocoma ... 142
Dimorpholepis ... 136
Diodontium 141
Dioscorea 196
Dioscorideae 196
Diospyros 155
Diotosperma 142
Diplachne 226
Diplacrum 217
Diplanthera ... 167, 204
Diplarrhena 193
Diplaspis 107
Diplax 222
Diplazium 232
Diploglottis 41
Diplolaena 20
Diplolobium 68
Diplopappus 131
Diplopeltia 42
Diplopogon 223
Diplospora 126
Diplostephium ... 131
Diplora 232
Dipodium 187
Dipogonia 223
Dipteracanthus ... 167
Dipteranthemum ... 48
Disarrhenum 223
Discaria 104
Dischidia 159
Discopodium 214
Discospermum ... 126
Diselma 184
Diaemma 128
Disoon 175
Disiliaria 34
Dissocarpus 51
Distichlis 226
Distichostemon ... 43
Distylis 149
Dithyrocarpus ... 206
Dithyroategia ... 139
Ditoca 46
Ditomostrophe ... 29
Diuris 189
Dodonaea 42
Dolichandra 167
Dolichandrone ... 167
Dolichos 71
Donatia 146
Donia 68
Doodia 232
Dopatrium 164
Doryanthes 196
Doryphora 5
Dracaena 197
Dracophyllum ... 183
Drakaea 191
Drapetes 56
Drimya 4
Drosera 13
Droseraceae 12
Drummondita ... 20
Dryandra 123
Drymaria 46
Drymispermum ... 56
Drymophila 197
Drymophlaeus ... 202
Drynaria 233
Duboisia 163
Dunbaria 71
Duperreya 160
Duttonia ... 136, 174
Dysophylla 168
Dysoxylum 16
Dysphania 50

Eadesia 162
Earlia 168
Ebelingia 22
Ebenaceae 155
Ebermayera 167
Ecdeiocolea 208
Echinocarpus ... 30
Echinochloa 218
Echinocroton ... 36
Echinodorus 205
Echinopogon ... 223
Echinopsilon 51
Echinospermum ... 175
Echinus 36
Eclipta 141
Ectrosia 227
Ebretia 176
Ehrharta 222
Elachanthera ... 197
Elachanthus 142
Elachocroton 37
Elachopappus ... 139
Elachothamnus ... 131
Elaeagneae 109
Elaeagnus 109
Elaeocarpus 30
Elaeodendron ... 40
Elaphoglossum ... 234
Elatine 14
Elatineae 14
Elatostemma ... 38
Elattostachys ... 41
Electrosperma ... 207
Eleocharis 211
Eleochiton 211
Elephanthopus ... 130
Elettaria 185
Eleusine 225
Eleuthranthes ... 127
Elichryaum 136
Elionurus 221
Elodea 194
Elynanthus 214
Elythrophorus ... 227
Emblingia 8
Embothrium 122
Emex 53
Emilia 142
Emmenospermum ... 103
Emphysopus 130
Empleurosma ... 42
Encephalartos ... 184
Enchylaena 52
Enchysia 144
Endiandra 6
Enhalus 193
Enhydra 141
Enneapogon 224
Entada 74
Enterolobium ... 80
Enydra 141
Epacrideae 177
Epacris 177, 181, 183
Epaltes 134
Epiandra 216
Epiblema 189
Epicharis 16
Epilobium 84
Epipogum 188
Epipremnum ... 203
Epithinia 126
Epitriche 139
Eragrostis 226
Eranthemum 168
Erechtites 143
Eremaea 97
Eremodendron ... 174
Eremophila 174
Eremopyxis 90
Eremosyne 83
Eria 187
Eriachne 224
Erianthus 221
Ericaceae 177
Ericomyrtus 91
Erigeron 133
Eriocalia 108
Eriocauleae 207
Eriocaulon 207
Eriochilus 191
Eriochiton 51
Eriochlamys 140
Eriochloa 218
Eriocladium 139
Erioglossum 41
Eriosema 72
Erioatemon 18
Eritrichum 175
Erodiophyllum ... 131
Erodium 23
Erpetion 10
Erycibe 159
Eryngium 108
Erysimum 9
Erythraea 151
Erythrina 71
Erythrophlaeum ... 74
Erythrorrhis 188
Erythroxylum ... 23
Etaeria 188
Ethulia 129
Ethuliopsis 134
Euandra 217
Eucalyptus 97
Eucarya 110
Euchilopsis 64
Euchilus 62
Euchiton 134
Eucryphia 83
Eudesmia 97
Eugenia 101
Eulalia 221
Eulophia 187
Eulophus 187
Euodia 20
Euonymus 40
Euosma 153
Eupatorium 130
Euphorbia 31
Euphorbiaceae ... 31
Euphoria ... 41, 42
Euphrasia 165
Eupomatia 5
Euroschinus 44
Eurostorrhiza ... 217
Eurybia 131
Eurybiopsis 133
Eurycles 196
Euryomyrtus ... 91
Eustrephus 197
Eutacta 183
Eutassa 183
Eutaxia 64
Euthalea 150
Euxolus 49
Evandra 217
Evolvulus 160
Evonymus 40
Exarrhena 176
Excaecaria 37
Exocarpos 110
Exocarya 213
Eyrea 133

Fabricia 93
Fagraea 152
Fagus 39
Faradaya 173
Fatoua 38
Faustula 136
Fawcettia 7
Fenzlia 101
Festuca 226
Fibichia 225
Ficoideae 52
Ficus 37
Fieldia 166
Filices 229
Fimbristylis 211
Figuetia 202
Fischera 107
Fitzalania 5
Fitzgeraldia ... 5, 191
Fitzroya 184
Flacourtieae 11

Flagellaria 206
Flagellariaceae ... 206
Flaveria 141
Fleischera 24
Flemingia 72
Fleurya 39
Flindersia 17
Floriscopa 206
Floscopa 206
Flueggea 35
Fluvialea 203
Fontainea 37
Forstera 146
Forsteropsis 144
Fragosa 107
Franciscia 87
Frankenia 45
Frankeniaceae ... 45
Franklandia 114
Freirea 39
Fremya 100
Frenela 184
Freycinetia 202
Friesia 30
Froebelia 177
Fugosia ... 25, 26
Fuirena 213
Funkia 198
Fusanus 110

Gahnia 216
Gaiadendron 111
Galactia 71
Galeola 188
Galium 128
Gamelythrum ... 223
Gamozygis 139
Ganophyllum ... 44
Gardenia 125
Gardneria 152
Garuga 43
Gastrodia 188
Gastrolobium ... 58
Gaultiera 177
Geijera 21
Geissois 82
Geitonoplesium ... 197
Geleznowia 18
Genetyllis 87
Geniostoma 153
Genoplesium 189
Genosiris 193
Gentiana 152
Gentianeae 151
Geodorum 187
Geophila 127
Geraniaceae 123
Geranium 23
Gerbera 143
Gesneriaceae... ... 166
Geum 81
Gilberta 139
Gilesia 27
Gillbeea 83

Gingidium 108
Glabraria 6
Gleichenia 230
Glinus... 53
Glischrocaryon ... 85
Glochidion 34
Glossocarya 173
Glossodia 192
Glossogyne 141
Glossostigma... ... 164
Glyceria 226
Glycine 70
Glycosmis 21
Glycyrrhiza 69
Gmelina 173
Gnaphalium 134
Gnaphalodes 140
Gnephosia 139
Gomphandra... ... 109
Gompholobium ... 59
Gomphrena 47
Gongronema 158
Goniocarpus 85
Goniophlebium ... 233
Goniopogon 131
Goniopteris 233
Goniotriche 48
Goodenia 149
Goodeniaceae ... 146
Goodenoughia ... 140
Goodia 66
Goodyera 188
Gossypium 26
Gouania 104
Gramineae 218
Grammatotheca ... 143
Grammitis 234
Graptophyllum ... 168
Gratiola 164
Graumuellera ... 204
Greevesia 25
Grevillea 117
Grewia 31
Griesebachia 201
Grumilia 127
Guettarda 126
Guettardella 126
Guichenotia 29
Guilandina 72
Guioa 41
Gunnera 85
Gunnia ... 52, 187
Gunniopsis 52
Guttiferae 14
Gymnagathis ... 94
Gymnanthera ... 157
Gymnaphanes ... 134
Gymnema 158
Gymnochaete ... 214
Gymnococca 55
Gymnogramme ... 234
Gymnogyne 141
Gymnopteris... ... 234
Gymnoschoenus ... 214

Gymnosporia... ... 40
Gymnostachys ... 203
Gymnostyles... ... 141
Gymnothrix 220
Gynopogon 156
Gynothrix 24
Gynura 142
Gypsophila 46
Gyrocarpus 87
Gyrostachis 188
Gyrostemon 54
Gyrostephium ... 140

Habenaria 192
Haeckeria 138
Haemodoraceae ... 194
Haemodorum ... 194
Hakea 120
Halfordia 21
Halgania 176
Halocnemum... ... 52
Halodule 204
Halophila 193
Halorageae 85
Haloragis 85
Halothamnus ... 24
Hamamelideae ... 84
Hamelinia 198
Hannafordia 28
Haplocaryon 126
Haplopappus... ... 133
Haplotaxis 143
Hardenbergia ... 70
Harmogia 91
Harpullia 42
Harrisonia 22
Hartigshea 16
Haussmannia ... 167
Hearnia 16
Hedaroma 87
Hedera 105
Hedraianthera ... 40
Hedycarya 6
Hedyotis 125
Hedyscepe 201
Heleocharis 211
Helichrysum... ... 136
Helicia 116
Helicteres 27
Heliophytum... ... 176
Heliotropium ... 176
Helipterum 136
Hellenia 185
Helmholtzia 205
Helmia 196
Helminthostachys ... 229
Helophyllum... ... 146
Helopus 218
Helosciadium ... 108
Helothrix 214
Hemarthria 221
Hemiandra 170
Hemiarrhena ... 164
Hemicarpus 106

Hemichroa 49
Hemiclidia 123
Hemicyclia 35
Hemigenia 170
Hemiphora 173
Hemiphues 108
Hemisteirus 48
Hemistemma ... 1
Hemistephus... ... 1
Hemitelia 230
Heptapleurum ... 105
Heritiera 27
Hermannia 27
Hernandia 7
Herniaria 46
Herpestis 164
Herpolirium 199
Hetaeria ... 188, 205
Heterachne 227
Heterodendron ... 42
Heterolaena 55
Heteropogon... ... 222
Hewardia 197
Hexatheca 204
Hibbertia 1
Hibiscus 25
Hicksbeachia ... 116
Hierochloe 223
Hippocratea 41
Hippocrepandra ... 32
Hodgkinsonia ... 126
Hodgsonia 199
Hodgsoniola 199
Holcus 222
Hollandaea 122
Hologamium 221
Holostigma 143
Holotome 108
Homalanthus ... 37
Homalium 11
Homalocalyx... ... 90
Homalospermum ... 93
Homalostoma ... 182
Homoranthus ... 87
Hormogyne 154
Hovea... 66
Howea 201
Howittia 25
Hoya 159
Huanaca 107
Huegelia ... 88, 106
Huenefeldia 131
Hugonia 23
Humata 231
Humea 138
Husemannia 8
Hutchinsia 10
Huttia 1
Hyalochlamys ... 139
Hyalolepis 138
Hyalosperma ... 136
Hybanthera 158
Hybanthus 10
Hydnophytum ... 127

Hydriastele 201
Hydrilla 194
Hydrocharideae ... 193
Hydrocharis 193
Hydrocotyle 106
Hydroglossum ... 229
Hydrolea 161
Hydropeltis 4
Hydrophylleae ... 161
Hygrophila 167
Hylococcus 34
Hylogyne 122
Hymenanthera ... 11
Hymenolepis... ... 234
Hymenophyllum ... 230
Hymenosporum ... 12
Hymenotheca 54, 193
Hypaelyptum ... 213
Hypelytrum 213
Hypericinae 14
Hypericum 14
Hypocalymma ... 92
Hypoestes 168
Hypolaena 209
Hypolepis 234
Hypolytrum 213
Hypoporum 217
Hypoxis 195
Hypserpa 7
Hypsophila 40
Hyptiandra 22
Hyrtanandra... ... 39

Iberis... 10
Ichnocarpus 157
Ilex 155
Ilyogeton 164
Ilysanthes 164
Imbricaria 91
Imperata 221
Indigofera 67
Ionidium 10
Iphigenia 198
Ipomoea 159
Iridcae 193
Iris 193
Irvingia 105
Isachne 218
Isandra ... 163, 198
Ischaemum 221
Iseilema 222
Isoetes 228
Isoetopsis 142
Isolepis 213
Isolobus 143
Isoloma 231
Isopogon 112
Isoschoenus 214
Isotoma 144
Isotropis 58
Ixauchenus 130
Ixiochlamys 133
Ixiolaena 135
Ixiosporum 12

Ixodia 138
Ixora 126

Jacksonia 60
Jamboa 101
Jansonia 56
Jasmineae 155
Jasminum 155
Johnsonia 200
Josephia 123
Josephina 167
Junceae 206
Juncella 207
Juncus 206
Jungia 91
Jussieua 84
Justicia 167

Kaleniczenkia ... 56
Kamphusia 148
Kamptzia 100
Kayea 14
Kelleria 56
Kennedya 70
Kentia 201
Kentiopsis 201
Kentropsis 51
Keraudrenia ... 28
Kermadecia 116
Kernera 204
Kibara 5
Killingia 210
Kingia 201
Kippistia 131
Kissodendron ... 105
Klanderia 169
Knoxia 128
Kochia 50
Koehnea 84
Kreysigia 198
Kunzea 93
Kyllingia 210

Labiatae 168
Labichea 73
Laboucheria 74
Laccospadix 202
Lachnagrostis .. 224
Lachnocephalus ... 172
Lachnostachys ... 171
Lachnothalamus ... 140
Lactaria 156
Lagenaria 129
Lagenophora ... 130
Lagerstroemia ... 84
Lagrezia 49
Lagunaria 26
Lalage 65
Lamarchea 97
Lambertia 116
Lambertya 33
Lampocarya 216
Lamprochlaena ... 139
Lamprolobium ... 67

Laportea 39
Lappago 220
Lappula 175
Lasianthus 127
Lasiopetalum ... 29
Lastraea 233
Latourea 147
Latrobea 64
Lauraceae 6
Laurencia 144
Lavatera 23
Lawrencella 136
Lawrencia 24
Lawsonia 85
Laxmannia 200
Learosa 5
Lebediera 35
Ledgeria 188
Leea 105
Leeaceae 105
Leersia 222
Leewenhoekia ... 146
Legnephora 7
Leguminosae 56
Leichhardtia ... 7, 158, 184
Lemna 203
Lemnaceae 203
Lentibularinae ... 165
Leontopodium ... 134
Lepia 10
Lepidium 10
Lepidobolus 209
Lepidocoma 138
Lepidosperma ... 215
Lepidospora 215
Lepidozamia 184
Lepigonum 46
Lepilaena 204
Lepironia 213
Lepistemon 159
Leptaspis 220
Leptatherum... ... 221
Leptinella 141
Leptocarpus 209
Leptoceras 191
Leptochloa 225
Leptocyamus ... 70
Leptocytisus 64
Leptolobium 70
Leptomeria 110
Leptopteris 230
Leptorrhynchos ... 135
Leptosema 56
Leptospermum ... 93
Leptotriche 139
Lepturus 221
Lepyrodia 208
Leschenaultia ... 147
Lespedeza 70
Lestibudesia 49
Leucas 169
Leucocarpum ... 40
Leucolaena 107
Leucophyta 140
Leucopogon 177
Leucorchis 188
Leucothamnus ... 28
Leuzea 143
Levenhookia 146
Lhotzkya 90
Libertia 193
Licuala 202
Ligusticum 108
Ligustrum 156
Liliaceae 197
Limnanthemum ... 151
Limnophila 163
Limnostachya ... 205
Limosella 164
Lindernia 164
Lindsaya 231
Lineae 23
Linkia... 114
Linociera 156
Linospadix 201
Linschottenia ... 146
Linum 23
Liparis 185
Liparophyllum ... 151
Lipocarpha 213
Lippaya 125
Lippia... 171
Lissanthe ... 177, 181
Litobrochia 231
Litsea 6
Livistona 202
Lobelia ... 143, 144
Lobophyllum ... 176
Lobopogon 181
Logania 153
Loganiaceae 152
Lomandra 200
Lomaria 232
Lomariopsis 234
Lomatia 122
Lonchocarpus ... 72
Lopadocalyx .. 109
Lophoclinium ... 134
Lophostemon ... 100
Lopidium 229
Loranthaceae ... 111
Loranthophyllum ... 152
Loranthus 111
Lotus 66
Loudonia 85
Lourea 70
Loxocarya 209
Lucuma 154
Ludwigia 84
Luffa 129
Luisia 187
Lumnitzera 87
Luzula 206
Luzuriago 197
Lycium 162
Lycopodinae 228
Lycopodium ... 228, 229
Lycopus 168
Lyginia 208
Lygodictyon 229
Lygodium 229
Lyonsia 157
Lyperanthus 191
Lysicarpus 100
Lysimachia 153
Lysinema 182
Lysiosepalum ... 29
Lyssanthe 117
Lythrum 84
Maba 155
Macadamia 116
Macaranga 36
Macarthuria 53
Maccoya 175
Macdonaldia ... 188
Macgregoria 44
Mackinlaya 106
Macintyria 15
Macropidia 195
Macropiper 4
Macropodia 195
Macropteranthes ... 87
Macrostegia 55
Macrostigma ... 81
Macrozamia 184
Maesa 154
Magnoliaceae ... 4
Mahernia 27
Maireana 50
Malacochaete ... 213
Malaisia 38
Malcolmia 9
Mallophora 172
Mallotus 36
Malpighiaceae ... 43
Malva 24
Malvaceae 23
Malvastrum 24
Manglesia 117
Manisuris 221
Mapania 213
Mappa 36
Marattia 229
Marianthus 12
Mariscus 210
Marlea 124
Marquisia 127
Marsdenia 158
Marsilea 228
Maundia 204
Mayepa 156
Mazentoxeron ... 20
Mazus 163
Medicosma 20
Medinilla 102
Megalotheca 208
Meionectes 85
Melachne 216
Melaleuca 94
Melanthesa 35
Melanthesiopsis ... 35
Melanthium 198
Melastoma 102
Melastomaceae ... 102
Melhania 27
Melia 16
Meliaceae 16
Melichrus 177
Melicope 20
Melicytus 11
Melochia 27
Melodinus 156
Melodorum 5
Melothria 129
Memecylon 102
Memorialis 39
Meniocus 9
Meniascium 233
Menispermeae ... 7
Menkea 10
Mentha 168
Merkusia 148
Mertensia 230
Meryta 106
Mesembrianthemum 52
Mesomelaena ... 214
Mesotriche 177
Methorium 27
Metrosideros... ... 100
Mezoneuron 72
Michiea 181
Micraira 224
Micrantheum ... 32
Microcachrys ... 184
Microcarpaea ... 164
Microchloa 225
Microclisia 8
Microcorys 170
Microcybe 18
Microgyne 133
Microlaena 222
Microlepidium ... 10
Microlepis 231
Micromelum 21
Micromyrtus... ... 90
Micropyxis 153
Microsciadium ... 107
Microseris 143
Microstachya ... 37
Microstemma ... 159
Microstylis 188
Microtis 190
Miliusia 5
Milletia 68
Milligania ... 85, 198
Millotia 138
Milnea 16
Mimosa 74
Mimulus 163
Mimusops 154
Minuria 131
Mirbelia 59
Mischocarpus ... 41
Mitragyne 152
Mitrasacme 152
Mitreola 152
Mitrephora 5
Mniarum 46
Modecca 128
Molkenboeria ... 148
Mollinedia 5
Molloya 116
Mollugo 53
Momordica 129
Monacather 225
Monencyanthes ... 136
Monenteles 134
Monimiaceae... ... 5
Monochoria 205
Monococcus 54
Monogramma ... 232
Monoploca 10
Monotaxis 32
Monotoca 177
Montia ... 45, 46
Moonia 140
Moraea 193
Morelotia 216
Morgania 163
Moriuda 126
Morisia 214
Morna 135
Moschosma 168
Motherwellia ... 105
Mucuna 71
Muehlenbeckia ... 53
Muehlenbergia ... 223
Mukia 129
Murraya 21
Musa 185
Myoporinae 174
Myoporum 175
Myosotis 176
Myosurus 3
Myriocephalus ... 139
Myriogyne 141
Myriophyllum ... 86
Myristica 6
Myristiceae 6
Myrmecodia 127
Myrsinaceae 154
Myrsine 154
Myrtaceae 87
Myrtus 101

Nablonium 140
Nageia 184
Najas 203
Nania... 100
Nasturtium 9
Needhamia 181
Negria 166
Nelitris 101
Nelsonia 167
Nelumbium 4
Nelumbo 4
Nematoceras... ... 190
Nematolepis 20
Nematophyllum ... 66
Nematopus 139
Nematostigma ... 193
Nemedra 16
Neoroepera 35
Neottopteris 232
Nepenthaceae ... 83
Nepenthes 83
Nephelium 42
Nephrodium 233
Nephrolepis 233
Neptunia 74
Nertera 127
Nesaea 85
Nesodaphne 6
Nesogenes 172
Nettoa 31
Neurachne 220
Newcastlia 172
Nicotiana 162
Niemeyera ... 154, 192
Nipa 202
Nipaceae 202
Niphobolus 233
Nitraria 22
Notelaea 156
Nothochlaena ... 231
Notholaena 231
Nothopanax 105
Notomerium 157
Notothixos 111
Nuytsia 111
Nyctagineae 54
Nymphaea 4
Nymphaeaceae ... 4
Nyssanthes 47

Oberonia 185
Obione 49
Ochna... 17
Ochnaceae 17
Ochrolasia 1
Ochrosia 156
Ocimum 168
Octoclinis 184
Ocymum 168
Oenanthe 108
Oenothera 84
Ogcerostylus... ... 139
Olacineae 109
Olax 109
Oldenlandia 125
Olea 155
Oleandra 233
Olearia 131
Oliganthemum ... 133
Oligarrhena 181
Omalanthus 37
Omphacomeria ... 110
Onagreae 84
Onchosepalum ... 209
Oncinocalyx 171
Oncosporum 12
Oncostylia 211
Opercularia 127
Ophioderma 229

Ophioglossum ... 229
Ophiorrhiza ... 125
Ophiuros ... 221
Opilia ... 109
Oplismenus ... 220
Orchideae ... 185
Oreobolus ... 214
Oreocallis ... 122
Oreomyrrhis ... 108
Oreostylidium ... 146
Orites ... 116
Oritina ... 116
Ormocarpum ... 69
Ornithochilus ... 187
Orobanche ... 165
Orobancheae ... 165
Orthoceras ... 189
Orthopogon ... 220
Orthosiphon ... 168
Orthostemon ... 152
Orthotropis ... 57
Orthrosanthus ... 193
Oryza ... 222
Osbeckia ... 102
Osbornia ... 100
Oschatzia ... 107
Osmunda ... 230
Osteocarpum ... 51
Otanthera ... 102
Ottelia ... 193
Ourisia ... 164
Owenia ... 16
Oxalis ... 23
Oxleya ... 17
Oxycladium ... 59
Oxylobium ... 57
Oxymyrrhine ... 91
Oxystelma ... 157
Ozothamnus ... 136

Pachycornia ... 52
Pachygone ... 8
Pachynema ... 3
Pachysura ... 140
Pacquerina ... 130
Pagetia ... 21
Palmae ... 201
Palmeria ... 6
Panactia ... 135
Panax ... 105
Pancovia ... 41
Pandaneae ... 202
Pandanophyllum ... 213
Pandanus ... 202
Panicum ... 218
Panopsis ... 116
Papaver ... 8
Papaveraceae ... 8
Pappophorum ... 224
Paractaenum ... 218
Paratropia ... 105
Parduyna ... 198
Parietaria ... 39
Parinarium ... 81

Paritium ... 25
Parkeria ... 230
Parsonsia ... 157
Paryphantha ... 90
Paspalum ... 218
Passiflora ... 128
Passifloreae ... 128
Patania ... 230
Patersonia ... 193
Pavetta ... 126
Pavonia ... 25
Pedalinae ... 167
Pedilonum ... 186
Pelargonium ... 23
Pellaea ... 231
Pelonastes ... 86
Peltophorum ... 72
Pemphis ... 85
Pennantia ... 109
Pennisetum ... 220
Pentaceras ... 22
Pentachondra ... 181
Pentacraspedion ... 223
Pentadactylon ... 114
Pentadynamis ... 66
Pentagonaster ... 93
Pentalepis ... 140
Pentapanax ... 105
Pentapeltis ... 107
Pentapogon ... 223
Pentaptelion ... 177
Pentataphrus ... 177
Pentatropis ... 158
Peperomia ... 4
Peplidium ... 164
Pericalymma ... 93
Pericampylus ... 7
Pernettya ... 177
Peros ... 177
Perojoa ... 177
Perotis ... 220
Perrottetia ... 40
Persoonia ... 114
Petalogyne ... 73
Petalolepis ... 136
Petalostigma ... 34
Petalostylis ... 73
Petermannia ... 196
Petrolobium ... 72
Petrophila ... 111
Phacelothrix ... 139
Phaenopoda ... 134
Phajus ... 188
Phalangium ... 199
Phaleria ... 56
Phanerandra ... 177
Pharbitis ... 159
Phaseolus ... 71
Phebalium ... 18
Phegopteris ... 238
Pherosphaera ... 184
Philhydreae ... 205
Philhydrum ... 205
Philotheca ... 20

Philoxerus ... 47
Philydrella ... 205
Philydrum ... 205
Phlebocalymma ... 109
Phlebocarya ... 194
Phlebodium ... 233
Phoberos ... 11
Pholidia ... 174
Pholidiopsis ... 174
Pholidota ... 185
Phormium ... 197
Phragmites ... 227
Phreatia ... 186
Phyllachne ... 146
Phyllanthus ... 34
Phyllocalymma ... 139
Phyllocladus ... 184
Phyllodium ... 69
Phylloglossum ... 229
Phyllopappus ... 143
Phyllota ... 62
Phymatocarpus ... 96
Phymatodes ... 233
Physalis ... 161
Physolobium ... 70
Physopsis ... 172
Phytolacceae ... 54
Picrophyta ... 149
Pigea ... 10
Pileanthus ... 89
Pilitis ... 183
Pilularia ... 228
Pimelea ... 55
Piper ... 4
Piperaceae ... 4
Piptandra ... 91
Piptocalyx ... 6
Piptomeria ... 60
Piptostemma ... 139
Pipturus ... 39
Pisonia ... 54
Pithecolobium ... 80
Pithocarpa ... 138
Pithyrodia ... 172
Pittosporeae ... 11
Pittosporum ... 11
Plagianthus ... 24
Plagiogyria ... 232
Plagiolobium ... 66
Plagiosetum ... 220
Plantagineae ... 153
Plantago ... 153
Platycarpidium ... 107
Platycerium ... 234
Platychilum ... 66
Platylobium ... 64
Platyloma ... 231
Platyptelea ... 82
Platysace ... 107
Platytheca ... 15
Platyzoma ... 230
Plectranthus ... 168
Plectronia ... 126
Pleea ... 198

Pleiococca ... 21
Pleiogyne ... 141
Pleiogynium ... 44
Pleistichostemon ... 43
Pleogyne ... 8
Pleopeltis ... 233
Pleurandra ... 1
Pleurandropsis ... 18
Pleurocarpaea ... 129
Pleurogramme ... 232
Pleuropappus ... 139
Plexaure ... 186
Plinia ... 101
Plinthacanthus ... 225
Plokiostigma ... 44
Pluchea ... 133
Plumbagineae ... 45
Plumbago ... 45
Poa ... 226
Podocarpus ... 184
Podocoma ... 133
Podolepis ... 135
Podolobium ... 57
Podopetalum ... 72
Podosperma ... 134
Podostemoneae ... 166
Podotheca ... 134
Poecilodermis ... 26
Pogonatherum ... 221
Pogonetes ... 148
Pogonia ... 175, 188
Pogonolepis ... 139
Pogonolobus ... 126
Pogostemon ... 168
Poiretia ... 66, 182
Polanisia ... 8
Pollia ... 206
Pollichia ... 176
Pollinia ... 221
Polyalthia ... 5
Polycalymma ... 139
Polycarpaea ... 47
Polycarpon ... 47
Polycnemon ... 49
Polygala ... 14
Polygaleae ... 14
Polygonaceae ... 53
Polygonum ... 53
Polymeria ... 160
Polyosma ... 82
Polyphragmon ... 126
Polypodium ... 233
Polypogon ... 220
Polypompholyx ... 166
Polystichum ... 233
Polyzone ... 87
Pomaderris ... 103
Pomatoderris ... 103
Pomax ... 127
Ponceletia ... 182
Pongamia ... 72
Pontederiaceae ... 205
Popowia ... 5
Porana ... 160

Poranthera ... 32
Porospermum ... 105
Portenschlagia ... 40
Portulaca ... 45
Portulaceae ... 45
Posidonia ... 204
Potamogeton ... 204
Potamophila ... 222
Potentilla ... 81
Pothos ... 203
Pouzolzia ... 39
Pozoa ... 107
Pozoopsis ... 107
Prasophyllum ... 189
Pratia ... 143
Premna ... 173
Primulaceae ... 153, 154
Prionosepalum ... 209
Prionotes ... 182
Pritzelia ... 106, 205
Procris ... 38
Pronaya ... 12
Prostanthera ... 169
Proteaceae ... 111
Prunella ... 169
Pseudalangium ... 124
Pseudanthus ... 32
Pseudatalaya ... 41
Pseudomorus ... 38
Pseudopholidia ... 174
Psilotrichum ... 48
Psilotum ... 228
Psoralea ... 66
Psychotria ... 127
Psydrax ... 126
Pteridium ... 231
Pterigeron ... 133
Pteris ... 231
Pterocaulon ... 134
Pterochaete ... 135
Pterolobium ... 72
Pteroloma ... 69
Pteropogon ... 136
Pterostigma ... 163
Pterostylis ... 190
Pterygopappus ... 134
Ptilotus ... 48
Ptychosema ... 67
Ptychosperma ... 202
Pultenaea ... 62
Pumilo ... 138
Punicella ... 92
Pycnarrhena ... 8
Pycnolachne ... 171
Pycnosorus ... 140
Pycnospora ... 70
Pycreus ... 210
Pygmaea ... 164
Pyxidaria ... 164

Quinetia ... 139
Quintinia ... 82
Quoya ... 172

Randia ... 125
Ranunculaceae ... 3
Ranunculus ... 3
Raoulia ... 134, 136
Rapuntium ... 143
Ratonia ... 41
Reedia ... 217
Regelia ... 96
Remirea ... 214
Renealmia ... 193
Restiaceae ... 207
Restio ... 208
Rhabdothamnus ... 166
Rhagodia ... 50
Rhamnaceae ... 102
Rhamnus ... 102
Rhamphicarpa ... 165
Rhamphidia ... 188
Rhaphidophora ... 203
Rhipogonum ... 197
Rhizophora ... 86
Rhizophoreae ... 86
Rhizospermae ... 228
Rhodamnia ... 101
Rhodanthe ... 136
Rhododendron ... 177
Rhodomyrtus ... 101
Rhodosphaera ... 44
Rhopala ... 116
Rhus ... 44
Rhyncharrhena ... 158
Rhynchosia ... 71
Rhynchospora ... 214
Rhynchostemon ... 28
Rhytidandra ... 124
Rhytidanthe ... 135
Rhytidosporum ... 12
Richea ... 140, 183
Ricinocarpus ... 33
Riedleya ... 27
Rinzia ... 91
Ripogonum ... 197
Rochelia ... 175
Roea ... 60
Roettlera ... 166
Roepera ... 22
Roeperia ... 8, 33
Rosaceae ... 81
Rostellaria ... 167
Rostellularia ... 67
Rotala ... 84
Rottboellia ... 221
Rottlera ... 136
Roupala ... 116
Rourea ... 81
Roxburghia ... 196
Roxburghiaceae ... 196
Rubiaceae ... 125
Rubus ... 81
Ruelingia ... 27
Ruellia ... 167
Rumex ... 53
Ruppia ... 204
Rutaceae ... 17

Rutidochlamys ... 135
Rutidosis 138
Ryssopterys 43

Saccharum 221
Saccolabium 187
Saccopetalum ... 5
Sagenia 233
Sagina 46
Salacia 41
Salicarieae 84
Salicornia 52
Salisia... 93
Salomonia 14
Salsola 52
Salsolaceae 49
Salvia... 169
Salvinia 228
Samara 154
Sambucus 128
Samolus 153
Samydaceae 11
Sandfordia 18
Santalaceae 110
Santalum 110
Sapindaceae 41
Sapindus 41
Saponaria 46
Sapota 154
Sapotaceae 154
Sarauia 14
Sarcocephalus ... 125
Sarcochilus 187
Sarcogonum 53
Sarcopetalum ... 7
Sarcostemma... ... 158
Sarcozygium 22
Sarotes 29
Saussurea 143
Saxifrageae 82
Scaevola 148
Scalia 135
Scalliopsis 135
Schelhammera ... 198
Schedonorus 226
Schenodorus 226
Schidiomyrtus ... 91
Schizachyrium ... 222
Schizaea 229
Schizeilema 107
Schizoloma 231
Schizomeria 82
Schizopleura 96
Schleidenia 176
Schmidelia 41
Schoberia 52
Schoenia 136
Schoenodum ... 208, 209
Schoenolaena ... 107
Schoenus 214
Scholtzia 91
Schuermannia ... 87
Schwaegrichenia ... 195
Scirpidium 211

Scirpodendron ... 213
Scirpus 213
Scitamineae 185
Scleranthus 46
Scleria 217
Sclerochlamys ... 50
Sclerolaena 51
Scleroleima 142
Sclerothamnus ... 64
Scolopendrium ... 232
Scolopia 11
Scoparia 164
Scorzonera 143
Scottia 65
Scrophularinae ... 163
Scutellaria 169
Scyphiphora 126
Scyphocoronis ... 138
Seaforthia 202
Sebaea 151
Sebastiania 37
Secamone 157
Securinega 35
Selaginella 229
Selliera 149
Selliguea 234
Selwynia 7
Semecarpus 44
Senebiera 10
Senecio 142
Sentis... 174
Seqnoia 183
Serianthes 80
Seringea 28
Sersalisia 154
Sesbania 68
Seseli 108
Sesuvium 52
Setaria 220
Sheffieldia 153
Sicyos 129
Sida 24
Sideroxylon 154
Siebera 107
Siegesbeckia 141
Sieglingia 226
Siemssenia 135
Sieversia 81
Siloxerus 139
Silphiospermum ... 130
Simarubeae 22
Simsia 113
Siona 199
Siphonodon 41
Sisymbrium 9
Sisyrinchium ... 193
Sium 108
Skinnera 159
Skirrophorus... ... 139
Sloanea 30
Smilax 197
Smithia 69
Solanaceae 161
Solanum 161

Soleniscia 177
Solenogyne 130
Solenostigma... ... 37
Soliva 141
Sollya 12
Sondera 13
Sonneratia 102
Sonzaya 44
Sophora 72
Sopubia 165
Sorghum 222
Sorostachys 210
Sowerbaea 199
Spadostyles 62
Spanoghea 42
Sparganium 203
Spartothamnus ... 171
Spathodea 167
Spathoglottis ... 188
Spergularia 46
Spermacoce 128
Spermalepis 100
Spermaxyrum ... 109
Sphaeranthos ... 134
Sphaerolobium ... 60
Sphaeromorphaea 134, 141
Sphaeropus 217
Sphenoclea 144
Sphenotoma 183
Sphincterostoma ... 182
Spiculaea 191
Spilanthes 141
Spinifex 220
Spiraeanthemum ... 83
Spiranthera 12
Spiranthes 188
Spirodela 203
Spiropodium 133
Spodiopogon ... 221, 222
Spondias 44
Sponia 37
Sporobolus 224
Sprengelia 182
Spyridium 103
Stachystemon ... 32
Stackhousia 44
Stackhousieae ... 44
Stadmannia 41
Statice 45
Stawellia 200
Steetzia 131
Stegania 232
Steiroglossa 130
Stekhovia 149
Stellaria 46
Stemodia 163
Stenanthemum ... 103
Stenanthera 177
Stenocarpus 122
Stenochilus 174
Stenochlaena... ... 234
Stenodiscus 103
Stenopetalum ... 9
Stenotaphrum ... 220

Stephania 7
Sterculia 26
Sterculiaceae... ... 26
Stipa 223
Stiphelia 177
Stirlingia 113
Stomarrhena... ... 177
Strangea 116
Stravadium 102
Streblorrhiza ... 68
Streptachne 223
Streptocarpus ... 166
Streptoglossa ... 133
Streptothamnus ... 11
Striga... 165
Strobilanthes ... 167
Stromatopteris ... 230
Strongylodon ... 71
Strongylospermum ... 141
Strychnos 152
Strzeleckia ... 17, 144
Stuartina 134
Sturmia 185
Sturtia 26
Stylideae 144
Stylidium ... 124, 144
Stylobasium 81
Stylocoryne 126
Stylolepis 135
Stylocerus 139
Stylurus 117
Stypandra 199
Styphelia 177
Styraceae 155
Suaeda 52
Suriana 22
Swainsona 68
Swammerdamia ... 136
Sycios 129
Symphiomera ... 141
Symphionema ... 114
Symphyomyrtus ... 97
Symphyonema ... 114
Symphyopetalum ... 20
Symplocos 155
Synaphea 113
Synaptantha... ... 125
Syncarpia 100
Synophlebium ... 231
Synostemon 34
Synoum 16
Syzygium 101

Tabernaemontana ... 157
Tacca 194
Taccaceae 194
Taeniophyllum ... 187
Talinum 45
Tamarindus 73
Tapeinocheilos ... 185
Tarenna 126
Tarrietia 27
Tasmannia 4
Tatea 173

Taxanthema 45
Tecoma 166
Tecticornia 52
Telanthera 47
Telmatophace ... 203
Telopea 122
Temminckia 148
Templetonia 65
Tenagocharis ... 205
Tephrosia 67
Terminalia 87
Ternstroemiaceae ... 14
Tersonia 54
Tetracarpaea... ... 83
Tetracera 3
Tetracheilus 74
Tetragonella 52
Tetragonia 52
Tétralepis 214
Tetralobus 166
Tetranthera 6
Tetrapasma 104
Tetraphylax 149
Tetrapora 91
Tetrarrhena 222
Tetratelia 8
Tetratheca 15
Teucrium 171
Thalamia 184
Thamnopteris ... 232
Thecanthes 55
Theleophyton ... 49
Thelychiton 186
Thelymitra 188
Themeda 222
Therogeron 131
Thesium 111
Thespesia 26
Thespidium 134
Thlaspi 10
Thomasia 28
Thouarsea 220
Thozetia 159
Threlkeldia 52
Thrixspermum ... 187
Thryptomene ... 90
Thuarea 220
Thunbergia 167
Thymeleae 55
Thyrsacanthus ... 168
Thysanella 198
Thysanotus 198
Tiaridium 176
Tiliaceae 30
Tillaea 83
Timonius 126
Tinospora 7
Titania 185
Tittmannsia 164
Tmesipteris 228
Todea... 230
Torenia 164
Tournefortia 176
Toxanthus 138

Trachycaryon ... 36
Trachymene 107
Trachypogon... ... 222
Tragia... 36
Tragus 220
Trema... 37
Tremandra 16
Tremandreae... ... 15
Trianthema 52
Tribonanthes ... 194
Tribulopsis 22
Tribulus 22
Trichantodium ... 139
Trichelostylis ... 211
Trichinium 48
Trichobasis 96
Trichobolus 81
Trichocline 143
Trichodesma 176
Trichodium 224
Tricholoma 164
Trichomanes 229
Trichosanthes ... 128
Trichosiphon... ... 26
Trichostegia 135
Tricondylus 122
Tricoryne 199
Tricostularia... ... 214
Trigsstrotheca ... 53
Triglochin 204
Trigonella 66
Trineuron 142
Triodia 226
Tripetelus 128
Triphelia 87
Tripladenia 198
Tripterococcus ... 44
Triptilodiscus ... 136
Trirhaphis 227
Trisetum 224
Tristania 100
Tristaniopsis... ... 100
Tristellateia 43
Tristichocalyx ... 7
Trithuria 207
Triticum 227
Triumfetta 30
Trochocarpa 181
Trollius 3
Trymalium 103
Turraea 16
Turritis 9
Tylophora 158
Typha... 203
Typhaceae 203
Typhonium 203

Umbelliferae... ... 106
Uncinia 217
Ungeria 27
Uniola 226
Urachne 223
Uragoga 127
Uralepis 226

Uraria 70
Urena... 25
Urocarpus 18
Urodon 62
Urostigma 37
Urtica 39
Urticaceae 37
Utricularia 165
Uvaria 4
Uvedalia 163

Vaccinium 177
Vachellia 74
Vaginularia 232
Vallisneria 194
Vandellia 164
Veitchia 201
Velleya 150
Ventenata 144
Ventenatia 177
Ventilago 102
Verbena 171
Verbenaceae... ... 171
Vernonia 130
Veronica 164
Verreauxia 148
Verticordia 88
Vigna 71
Vilfa 224
Villaresia 109
Villarsia 151
Viminaria 61
Vincetoxicum ... 157
Viniferae 104
Viola 10
Violaceae 10
Viraya 135
Viscum 111
Vitex 173
Vitis 104
Vittadinia 133
Vittaria 231
Vlamingia 10
Vulpia ... 226, 227

Wahlenbergia ... 144
Waitzia 135
Walcottia 171
Waltheria 27
Warburtonia... ... 1
Wartmannia... ... 37
Webera 126
Wedelia 141
Wehlia 91
Weinmannia... ... 82
Westonia 66
Westringia 171
Wichurea 103
Wickstroemia ... 56
Wilckia 9
Wilkiea 5
Wilsonia 161
Wistaria 68
Wittsteinia 177
Wolffia 203
Wollastonia 141
Woodwardia... ... 232
Woollsia 182
Wormia 1
Wrightia 157
Wrixonia 170
Wurmbsea 198

Xanthochrysum ... 136
Xanthophyllum ... 15
Xanthorrhoea ... 201
Xanthosia 107
Xanthostemon ... 100
Xanthoxylum ... 21
Xerochloa 220
Xerosollya 12
Xerotes 200
Ximenia 109
Xiphopteris 233
Xylocarpus 16
Xylomelum 116
Xylosma 11
Xyridanthe 136
Xyrideae 206
Xyris 256
Xystidium 220

Youngia 143
Zaleya 52
Zamia... 184
Zannichellia 204
Zanonia 129
Zanthoxylum ... 21
Zapania 171
Zehneria 129
Zichya 70
Zieria 17
Zizyphus 102
Zornia... 69
Zostera 204
Zosterostylis... ... 189
Zoysia 221
Zygia 80
Zygomenes 205
Zygophylleae ... 22
Zygophyllum ... 22

OMISSIONS.

Senebiera (Coronopus) 10
Sisymbrium (Geococcus) 9
Ventilago (Berchemia) 102

McCarron, Bird and Co., Printers, 479 Collins-street, Melbourne.

THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW

AN INITIAL FINE OF 25 CENTS

WILL BE ASSESSED FOR FAILURE TO RETURN THIS BOOK ON THE DATE DUE. THE PENALTY WILL INCREASE TO 50 CENTS ON THE FOURTH DAY AND TO $1.00 ON THE SEVENTH DAY OVERDUE.

JUN 6 1960

JAN 24 1963

FEB 13 1963

LIBRARY USE

NOV 21 1963

DEC 12 1969

Book Slip—10m-8,'58(5916s4)458

174669
Mueller, F.
Second systematic
census of Australian

Call Number:
QK431
M93

Mueller
QK431
M93

174669